

Assessment of Iodine Nutritional Status of School-Age Children in Kolkata District of West Bengal State in Post-Iodation Scenario

Udayan Bhattacharya and Amar K Chandra

Correspondence: Amar K Chandra, Department of Physiology, Endocrinology and Reproductive Physiology Laboratory, University College of Science and Technology, University of Calcutta, 92, A.P.C. Road, Kolkata 700009, West Bengal, India. Tel: +91 94331 61840.
E-mail <physiology.ac@gmail.com>.

ABSTRACT

To evaluate the state of iodine nutrition in post-iodation scenario, 3500 children were examined clinically for endemic goitre. Iodine and thiocyanate were measured in 240 urine samples; iodine content in 210 salt samples was measured. Total goitre prevalence was 6.1%. Median urinary iodine level was 21.80 µg/dl, and mean (\pm SD) urinary thiocyanate was 0.89 ± 0.49 mg/dl. Iodine content of only 11.9% salt samples was below recommended level of 15 ppm, 25.2% was between 15 and 30 ppm and 62.9% was >30 ppm. Iodine deficiency disorders are thus clinically mild public health problem of the studied population; however, they have no biochemical iodine deficiency. Studied population found exposed to thiocyanate load that might be the possible cause for persistence of endemic goitre. People of Kolkata should be advised to eat commonly consumed goitrogenic foods after boiling and decanting the water. Periodical monitoring and evaluation of iodine status should be mandatory.

KEYWORDS: iodine deficiency disorders, goitre, salt iodation, urinary iodine, urinary thiocyanate

BACKGROUND

Iodine is required for the production of thyroid hormone by thyroid. Requirement of iodine per day is 90 µg for preschool children (0–59 months), 120 µg for school children (6–12 years), 150 µg for adolescents (>12 years) and adults and 250 µg for pregnant and lactating mothers [1]. In iodine deficiency, thyroid does not produce thyroid hormone, which may lead to a spectrum of diseases named as iodine deficiency disorders (IDDs). IDDs are manifested by goitre, stillbirths, spontaneous abortion and congenital anomalies, including endemic cretinism and irreversible brain damage [2, 3].

WHO estimated nearly 2 billion people are at risk of IDDs, including one-third of school children [4].

Total goitre prevalence (TGP) was lowest in America (10.6%), whereas it was highest in Europe (52.0%). Australia is mildly iodine deficient, and in Switzerland, median urinary iodine (MUI) excretion is 14 µg/dl [5]. In Portugal, MUI is 12.9 µg/dl [6]. In Ireland, 26% population is iodine deficient [7]. TGP of Southern Ethiopia is 48.9% [8] and north-west Ethiopia is 29.1% [9]. Israel is iodine deficient (MUI 8.3 µg/dl) [10].

In India, 350 million peoples are at risk of IDDs, as they consume inadequately iodized salt [11]. Of 587 districts, 282 surveyed for IDD and 241 have been found goitre endemic [12]. TGP of Maharashtra is 11.9% [11]. Tripura, Manipur, and Uttar Pradesh are affected by IDDs [13–17]. TGP of

Nainital is 15.9% (MUI 12.5 µg/dl) [12]. In Kullu, TGP is 23.4% (MUI 17.5 µg/dl) [18]. In Udaipur, TGP is 8.4% (MUI 20 µg/dl) [19]. In Kottayam, TGP is 7.05% (MUI 17.5 µg/dl) [20].

In 1962, Government of India (GOI) launched National Goitre Control Programme (NGCP). In August 1992, NGCP was renamed as National Iodine Deficiency Disorders Control Programme [21]. In 1997, GOI banned sale of non-iodized salt. In September 2000, GOI lifted ban stating 'On point of principle, compulsions in the matter of individual choice are undesirable' that results nationwide 12% decrease in coverage of iodized salt; in May 2006, GOI reimposed the ban [5, 22].

Iodine nutritional status of school children in Kolkata during post-iodation scenario is not available. Present investigation has been undertaken to evaluate iodine status of selected population with possible involvement of dietary goitrogens (if any) in Kolkata.

MATERIALS AND METHODS

Data collection

This school-based cross-sectional study was undertaken in North, East, and West Kolkata. List of schools and their enrolment number in each area were previously collected from School Education Department. Probability proportion to size methodology was adopted for selection of schools [1]. Total 30 schools (clusters) were selected. From each school, all children in the age group of 6–12 years of both sexes were clinically examined for goitre. Sample size was calculated based on assumed goitre prevalence 5%, confidence interval 95% and permissible error of margin 20%. Sample size of 1900 was obtained [15, 16, 23]. To increase precision of study, 3500 school children were examined.

Clinical examination of goitre

IDD is manifested by goitre [24]. Pupils were examined clinically for enlargement of thyroid. Grading of goitre was done as Grade 0, no palpable or visible goitre; Grade 1, goitre that was palpable but not visible when the neck was in normal position; and Grade 2, goitre that was visible when the neck was in normal position. TGP (Grade 1 + Grade 2 goitre) in

school children in the age group of 6–12 years is one of the most acceptable markers of IDD. Prevalence rate of 0–4.9% was considered as none, 5–19.9% as mild, 20.0–29.9% as moderate and >30.0% as a severe public health problem [1].

Measurement of urinary iodine

Iodine content in 30 urine samples collected at random from pupils in the age group of 6–12 years gives proper idea about iodine nutritional status in a region [1]. From each locality, 80 urine samples were randomly collected from children in the age group of 6–12 years of both sexes irrespective of their thyroid status. Total 240 urine samples were measured for iodine content by arsenite method [25]. Internal quality control was adopted during analysis [12, 26]. MUI <9.9 µg/dl was considered as insufficient, 10–19.9 µg/dl as adequate, 20–29.9 µg/dl as above requirements and ≥30 µg/dl as excessive iodine intake [1].

Measurement of urinary thiocyanate

The amount of thiocyanate excreted in urine is an excellent indicator for consumption of most common goitrogenic plant foods [27]. Dietary supplies of cyanogenic glycosides and glucosinolates might be estimated from concentration of thiocyanate in urine [28]. Thiocyanate was estimated by the method of Aldridge [29] and modified by Michajlovskij and Langer [30].

Measurement of salt iodation status

In India, iodine is supplemented through edible salts because salt is inexpensive and consumed by all classes of people throughout the year in almost uniform quantity. Iodine content of at least 35 salt samples collected at random provides a valid estimate of iodation status [1]. Students were asked to carry salt samples from their household next day. From each locality, 70 salt samples were collected. Total 210 salt samples were measured for iodine by iodometric titration [31].

Data analysis and management

As per WHO/UNICEF/ICCIDD, mean, median, mode, SD, skewness (*Sk*), kurtosis (*K*) and percentile values (P_{20} and P_{80}) were computed for urinary iodine [1]. For urinary thiocyanate, mean ± SD was

computed. Salt iodine values were categorized <15, 15–30 and >30 ppm. Pearson's product moment correlation was computed between urinary iodine and iodine content of salt samples [12, 13]. All descriptive statistical analysis was done in Microsoft Office Excel 2007 [1].

RESULTS

Overall, TGP of studied population was 6.1%. In North, East and West Kolkata TGP was 5.8, 7.0 and 5.5%, respectively (Table 1).

Overall, MUI excretion of Kolkata was 21.80 µg/dl (mean ± SD = 23.65 ± 11.74 µg/dl, mode = 20.75, confidence level 95% was between 22.16 and 25.14 µg/dl). In North, East and West Kolkata, MUI was 23.10, 21.70 and 18.60 µg/dl, respectively (Table 2, Fig. 1). Frequency distribution of MUI indicated that 6.25% population was iodine deficient, 32.50% population's iodine intake was adequate, 39.17% population's iodine intake was more than adequate and 22.08% population's iodine intake was excessive (Table 3). P_{20} was 13.45 µg/dl, and P_{80} was 30.75 µg/dl. Mean was much higher than median, indicating that the distribution was heavily skewed to

right ($Sk = +0.934262$) and histogram was platykurtic ($K = 0.773096$) (Table 2, Fig. 2).

Mean urinary thiocyanate (MUSCN) of total studied population was 0.89 ± 0.49 mg/dl. In North, East and West Kolkata, it was 0.84 ± 0.54 , 0.94 ± 0.49 and 0.88 ± 0.44 mg/dl, respectively (Table 4, Fig. 3).

Overall, 11.9% of collected salt samples were <15 ppm, and 62.9% salt samples were >30 ppm (Fig. 4). In North Kolkata, 15.7% salt samples were <15 ppm, and 70% salt samples were >30 ppm. In East Kolkata, 10% salt samples were <15 ppm, and 62.9% salt samples were >30 ppm. In West Kolkata, 10% salt samples were <15 ppm, and 55.7% salt samples were >30 ppm (Table 5). A significant correlation was found in between urinary iodine excretion and iodine content of edible salt samples ($p < 0.01$).

DISCUSSION

IDDs are not restricted in classical goitre endemic belt of India. Goitre is endemic in many newly introduced pockets in post-iodation scenario [11–20]. IDD is mild to moderately prevalent in certain non-conventional plain and coastal region of West Bengal (WB) [1]. Average TGP of WB is 9% [2].

Table 1. Goitre prevalence of the studied population of Kolkata district

| Studied localities of Kolkata | Population studied | Goitre prevalence (%) | | | | Severity as a public health problem |
|-------------------------------|--------------------|-----------------------|-----------|----------|-------------|-------------------------------------|
| | | Grade 0 | Grade 1 | Grade 2 | TGP (1 + 2) | |
| North | 1559 | 1468 (94.2) | 82 (5.2) | 9 (0.6) | 91 (5.8) | Mild |
| East | 1030 | 958 (93.0) | 63 (6.1) | 9 (0.9) | 72 (7.0) | Mild |
| West | 911 | 861 (94.5) | 46 (5.1) | 4 (0.4) | 50 (5.5) | Mild |
| Total | 3500 | 3287 (93.9) | 191 (5.4) | 22 (0.6) | 213 (6.1) | Mild |

Grade 0: no goitre, Grade 1: palpable goitre, Grade 2: visible goitre, TGP: sum of Grade 1 and Grade 2 goitre.

Table 2. Urinary iodine excretion pattern of the studied population of Kolkata district

| Studied localities | Number of urine samples assayed | Mean (±SD) (µg/dl) | Median (µg/dl) | Iodine intake | Iodine nutritional status |
|--------------------|---------------------------------|--------------------|----------------|--------------------|---------------------------|
| North | 80 | 26.43 ± 12.67 | 23.10 | Above requirements | More than adequate |
| East | 80 | 22.57 ± 10.20 | 21.70 | Above requirements | More than adequate |
| West | 80 | 21.94 ± 11.85 | 18.60 | Adequate | Optimal |
| Total | 240 | 23.65 ± 11.74 | 21.80 | Above requirements | More than adequate |




Fig. 1. Diagrammatic representation of urinary iodine excretion pattern in iodation scenario of Kolkata.

Table 3. Frequency distribution of the values of urinary iodine in Kolkata district

| Urinary iodine excretion pattern | Frequency | Total | Percentage |
|----------------------------------|-----------|-------|------------|
| Insufficient iodine intake | | | |
| 0–4.9 | 6 | 15 | 6.25 |
| 5.0–9.9 | 9 | | |
| Adequate iodine intake | | | |
| 10.0–14.9 | 42 | 78 | 32.5 |
| 15.0–19.9 | 36 | | |
| More than adequate iodine intake | | | |
| 20.0–24.9 (location of median) | 60 | 94 | 39.17 |
| 25.0–29.9 | 34 | | |
| Excessive iodine intake | | | |
| 30.0–34.9 | 15 | 53 | 22.08 |
| 35.0–39.9 | 15 | | |
| 40.0–44.9 | 3 | | |
| 45.0–49.9 | 8 | | |
| 50.0–54.9 | 10 | | |
| 55.0–59.9 | 0 | | |
| 60.0–64.9 | 2 | | |
| 65.0–69.9 | 0 | | |
| Total | 240 | 240 | 100 |

Of 19 districts in Gangetic WB, 8 have been surveyed for IDD and found goitre endemic. In certain districts of WB, IDDs are still prevalent, such as Howrah

(TGP 38%) [32], Sundarban Delta incorporating North 24 Parganas (33.1%) [33] and South 24 Parganas (38.2%) [34], Maldah (11.3%) [35], Birbhum (12.6%) [36], Purulia (25.9%) [37], Dakshin Dinajpur (18.6%) [38] and Darjeeling (8.7%) [39]. TGP of Kolkata is 6.1%. As per WHO/UNICEF/ICCIDD, severity of IDDs in Kolkata is clinically a mild public health problem in post-iodation scenario [1].

The indicator of IDDs eradication is a median value for urinary iodine concentration of 10 µg/dl, iodine content of 50% of urine samples should be >10 µg/dl and not <5 µg/dl of 20% population [1]. MUI of the studied region is well above the recommended level, and thus, in general, deficient iodine intake is not the criteria for the persistence of mild goitre endemicity (Table 1). MUI of the studied region found more than adequate, and they are susceptible to iodine-induced hyperthyroidism (IIH) in long run. Increasing dose of iodine in microgram range may cause hyper or hypothyroidism in those with past or present thyroid abnormalities [5]. In children, excess dietary iodine has been associated with goitre and thyroid dysfunction [40]. In present study, 6.25% population found iodine deficient, 32.50% population's iodine intake was adequate, 39.17% population's iodine intake was more than adequate and they are susceptible to IIH and 22.08% population's iodine intake is excessive and they are susceptible to risk of adverse health consequences like IIH and autoimmune thyroid diseases [1, 5]. MUI is reported >20 µg/dl in several states and districts of India [32, 41–43]. In Delhi and Jammu and Kashmir, MUI was >30 µg/dl [44, 45]. In present study, TGP is 6.1%, and 6.25% population is iodine deficient. It may be concluded that persisting goitre prevalence may be because of iodine deficiency or insufficient iodine intake.

Dietary goitrogens are substances that can interfere directly on iodine metabolism in thyroid. Goitrogens are classified as thiocyanate, isothiocyanate, cyanogenic glycosides, glucosinolates, thiourea, thionamides, flavonoids, etc. [46–50]. MUSCN is an excellent marker to study consumption of dietary goitrogens. Thiocyanate is exogenous in origin and a potent goitrogenic substance as anion with same molecular size as iodine. It inhibits thyroidal




Fig. 2. Histogram to show distributions of urinary iodine values after iodation in Kolkata.

Table 4. Urinary thiocyanate excretion pattern of the studied population of Kolkata district

| Studied localities | Number of urine samples | Mean (\pm SD) (mg/dl) |
|--------------------|-------------------------|--------------------------|
| North | 80 | 0.84 ± 0.54 |
| East | 80 | 0.94 ± 0.49 |
| West | 80 | 0.88 ± 0.44 |
| Total | 240 | 0.89 ± 0.49 |




Fig. 3. Diagrammatic representation of urinary thiocyanate excretion pattern of Kolkata.

accumulation of iodide. At higher dose, it competes with iodine during protein binding [46–50]. It serves as a substrate for thyroid peroxidase (TPO) to inhibit iodination of tyrosyl residues of thyroglobulin molecule [51]. Within normal thyroid, iodine serves a dual role, as a substrate for iodination of thyroglobulin and as a ligand for coupling reaction [52]. Thiocyanate binds with same regulatory site as




Fig. 4. Diagrammatic representation of overall salt iodation status of Kolkata district.

iodine but has a slightly different affinity [51]. Boiled extracts of goitrogenic plant foods showed maximum inhibition of TPO activity followed by cooked and raw extracts [53]. Regular consumption of goitrogenic plant foods affects thyroid physiology and may develop endemic goitre [54]. A large number of goitrogenic plant foods are commonly consumed by Indian population, and IDD persists in many such newly introduced pockets, despite recommended iodine intake [13–18, 32–34]. Earlier workers mentioned that mean value of urinary thiocyanate excretion obtained from non-goitrous population is 0.71 ± 0.34 mg/dl [55]. We have considered that value as a control. MUSCN excretion of studied region is significantly higher than control ($p < 0.01$). Therefore, the studied population of Kolkata metropolitan city in the age group of 6–12 years is exposed to excess thiocyanate load (Table 4).

Table 5. Salt iodation status of the studied population of Kolkata district

| Studied localities | Number of salt samples tasted | Iodine content of edible salt samples of Kolkata | | |
|--------------------|-------------------------------|--|------------|-------------|
| | | < 15 ppm | 15–30 ppm | > 30 ppm |
| North | 70 | 11 (15.7%) | 10 (14.3%) | 49 (70.0%) |
| East | 70 | 7 (10.0%) | 19 (27.1%) | 44 (62.9%) |
| West | 70 | 7 (10.0%) | 24 (34.3%) | 39 (55.7%) |
| Total | 210 | 25 (11.9%) | 53 (25.2%) | 132 (62.9%) |

According to Prevention of Food Adulteration act, iodine content of the salt samples shall not be <15 ppm at household level [56]. As per National Family Health Survey-III, 51% households were using adequately iodized salt whose iodine content was ≥ 15 ppm. Another 25% were using salt that had iodine content below recommended level of 15 ppm, and 24% of households were found to be using non-iodized salt [57]. In Kolkata, 11.9% of salt samples have iodine <15 ppm, 25.2% of salt samples have iodine from 15 to 30 ppm and 62.9% of families are consuming iodized salt >30 ppm of iodine (Table 5). The goal of salt iodation programme is that >90% of household should use iodized salt [1]. In Kolkata, 11.9% families' dietary need for iodine is not properly fulfilled by salt iodation programme, but their iodine status is optimum as evidenced by MUI. Kolkata is not located in iodine-deficient zone, so their extra need for iodine is managed by drinking water and other consumable foods. Iodine status of overall studied population is more than adequate, as studied population excretes more iodine because of intake of edible salts whose iodine content is more than recommended level of 30 ppm. Distribution of values of MUI around median indicates that there was no iodine deficiency, and salt iodation therefore had the required impact. Central tendency (median) of distribution lies between ranges 20 and 24.9 $\mu\text{g}/\text{dl}$, which indicates that iodine intake of studied population is more than adequate, and they are susceptible to iodine-induced thyroid diseases. After iodine prophylaxis, an increased iodine intake is associated with increased thyroiditis [58].

Findings of present investigation indicate that studied population of Kolkata is clinically mild iodine deficient. They are susceptible to adverse health consequences like IIH and autoimmune thyroid diseases

in long run. Their mild goitre endemicity may be because of iodine deficiency and relatively high consumption of goitrogenic plant foods.

People of Kolkata are ignorant about iodine nutrition, and they should be encouraged to avoid goitrogenic plant foods for eating or they should eat those foods after boiling and decanting water. The Central Health Directorate and salt manufacturing companies should be aware to maintain iodine concentration 30 ppm in edible salt samples at manufacturers' level. Periodical monitoring and evaluation of iodine status is necessary.

ACKNOWLEDGEMENT

The authors duly acknowledge all the students who were solely participated in the study.

FUNDING

This work was supported by University of Calcutta research grants BI (92) for Endocrinology and Reproductive Physiology Laboratory, Department of Physiology.

REFERENCES

1. WHO/UNICEF/ICCIDD. Assessment of Iodine Deficiency Disorders and Monitoring Their Elimination: A Guide for Programme Managers. Geneva: WHO Publication, 2007, 1–97.
2. Hetzel BS. An overview of the prevention and control of iodine deficiency disorders. In: Hetzel BS, Dunn JT, Stanbury JB (eds). The Prevention and Control of Iodine Deficiency Disorders. Amsterdam: Elsevier Science, 1987, 7–31.
3. Maberly GF, Trowbridge FL, Yip R, *et al.* Programs against micronutrient malnutrition: ending hidden hunger. *Annu Rev Public Health* 1994;15:277–01.
4. deBenoist B, McLean E, Andersson M, *et al.* Iodine deficiency in 2007: global progress since 2003. *Food Nutr Bull* 2008;29:195–02.

5. Zimmermann MB. Iodine deficiency. *Endocr Rev* 2009;30: 376–08.
6. Costa Leite J, Keating E, Pestana D, *et al.* Iodine status and iodised salt consumption in Portuguese school-aged children: the iogeneration Study. *Nutrients* 2017;9:1–14.
7. McNulty BA, Nugent AP, Walton J, *et al.* Iodine intakes and status in Irish adults: is there cause for concern? *Br J Nutr* 2017;117:422–31.
8. Workie SB, Abebe YG, Gelaye AA, *et al.* Assessing the status of iodine deficiency disorder (IDD) and associated factors in Wolaita and Dawro Zones School Adolescents, Southern Ethiopia. *BMC Res Notes* 2017;10:1–8.
9. Abebe Z, Gebeye E, Tariku A. Poor dietary diversity, wealth status and use of un-iodised salt are associated with goitre among school children: a cross-sectional study in Ethiopia. *BMC Public Health* 2017;17:1–11.
10. Hebrew University of Jerusalem. National survey in Israel reports one of the lowest iodine intakes in the world. *IDD Newsl* 2017;45:13.
11. Pandav CS, Yadav K, Srivastava R, *et al.* Iodine deficiency disorders (IDD) control in India. *Indian J Med Res* 2013; 138:418–33.
12. Kapil U, Pandey RM, Prakash S, *et al.* Assessment of iodine deficiency in school age children in Nainital District, Uttarakhand State. *Asia Pac J Clin Nutr* 2014; 23:278–81.
13. Chandra AK, Ray I. Evaluation and effectiveness of salt iodation status in Tripura, north east India. *Indian J Med Res* 2002;115:22–7.
14. Chandra AK, Singh LS, Tripathy S, *et al.* Iodine nutritional status of children in north east India. *Indian J Pediatr* 2006;73:795–8.
15. Chandra AK, Singh LH, Debnath A, *et al.* Dietary supplies of iodine and thiocyanate in the etiology of endemic goitre in Imphal east district of Manipur, north east India. *Indian J Med Res* 2008;128:601–5.
16. Chandra AK, Bhattacharjee A, Malik T, *et al.* Goitre prevalence and iodine nutritional status of school children in sub-Himalayan Tarai region of eastern Uttar Pradesh. *Indian Pediatr* 2008;45:469–74.
17. Chandra AK, Bhattacharjee A, Malik T, *et al.* Etiological factor for the persistence of endemic goitre in selected areas of Siddharthanagar district in eastern Uttar Pradesh, India. *J Pediatr Endocrinol Metab* 2009;22: 317–25.
18. Chander S, Kapil U, Jain V, *et al.* Iodine deficiency disorders in school age children in Kullu District, Himachal Pradesh. *Indian Pediatr* 2013;50:883–4.
19. Pradhan R, Choudhry M. Assessment of iodine deficiency disorders in urban areas of Udaipur District, Rajasthan. *Indian Pediatr* 2003;40:406–9.
20. Kapil U, Jayakumar PR, Singh P, *et al.* Assessment of iodine deficiency in Kottayam district, Kerala State: a pilot study. *Asia Pac J Clin Nutr* 2002;11:33–5.
21. Tiwari BK, Kundu AK, Bansal RD. National iodine deficiency disorders control programme in India. *Indian J Pub Health* 1995;39:148–51.
22. Sankar R, Pandav CS. Ban on sale of non-iodised salt for human consumption: a step in the right direction. *Natl Med J India* 2005;18:169–71.
23. SundarRao PSS, Richard J. Experimental studies. In: *An Introduction to Biostatistics: A Manual for Students in Health Sciences*. New Delhi: Prentice-Hall of India Private Limited, 2003, 187–202.
24. Dunn JT, Van der Harr F. Detection of iodine deficiency. In: *A Practical Guide to the Correction of Iodine Deficiency*. Technical Manual Number 3. Netherland: WHO/UNICEF/ICCIDD Publication, 1990, 13.
25. Karmarkar MG, Pandav CS, Krishnamachari KAVR, Principle and procedure for iodine estimation. In: *A Laboratory Manual*. New Delhi: Indian Council of Medical Research Publication, 1986, 1.
26. Westgard JO, Barry PL, Hunt MR. A multi-rule Shewhart chart for quality control in clinical chemistry. *Clin Chem* 1981;27:493–501.
27. Gaitan E. Environmental goitrogens other than iodine deficiency. *IDD Newsl* 1986;2:11–13.
28. Oginsky EL, Stein AE, Greer MA. Myrosinase activity in bacteria as demonstrated by the conversion of progoitrin to goitrin. *Proc Soc Exp Biol Med* 1965;119:360–4.
29. Aldridge WN. The estimation of micro quantities of cyanide and thiocyanate. *Analyst* 1945;70:474–5.
30. Michajlovskij N, Langer P. Studienuber Benziehungenzwischen Rhodanbildung, kropfbildender Eigenschaftvon Nahrungsmitteln. *J Physiol Chem* 1958;312:26–30.
31. De Maeyer EM, Lowenstein PW, Thilly CH. Titration methods for salt iodine analysis. In: Sullivan KM, Houston R, Gorstein J, *et al.* (eds). *Monitoring Universal Salt Iodation Programme*. Geneva: PAMM/MI/ICCIDD Publication, 1995, 86–96.
32. Chandra AK, Tripathy S, Lahiri D, *et al.* Iodine nutritional status of the school children in a rural area of Howrah district in Gangetic West Bengal. *Indian J Physiol Pharmacol* 2004;48:219–24.
33. Chandra AK, Tripathy S, Ghosh D, *et al.* Goitre prevalence and the state of iodine nutrition in Sundarban delta of North 24-Parganas in West Bengal. *Asia Pac J Clin Nutr* 2006;15:357–61.
34. Chandra AK, Tripathy S, Ghosh D, *et al.* Iodine nutritional status & prevalence of goitre in Sundarban delta of South 24-Parganas, West Bengal. *Indian J Med Res* 2005;122: 419–24.
35. Biswas AB, Chakraborty I, Das DK, *et al.* Iodine deficiency disorders among school children of Maldah, West Bengal, India. *J Hlth Pop Nutr* 2002;20:180–3.
36. Biswas AB, Chakraborty I, Das D, *et al.* Iodine deficiency disorders among the school children of Birbhum, West Bengal. *Current Science* 2004;87:78–80.

37. Biswas AB, Chakraborty I, Das DK, *et al.* Assessment of iodine deficiency disorders in Purulia District, West Bengal, India. *J Trop Pediatr* 2006;52:288–92.
38. Das DK, Chakraborty I, Biswas AB, *et al.* Iodine deficiency disorders among school children of Dakshin Dinajpur District, West Bengal. *Indian J Public Health* 2005;49: 68–72.
39. Biswas AB, Das DK, Chakraborty I, *et al.* Goitre prevalence, urinary iodine, and salt iodation level in sub-Himalayan Darjeeling district of West Bengal, India. *Indian J Public Health* 2014;58:129–33.
40. Suzuki H, Higuchi T, Sawa K, *et al.* “Endemic coast goitre” in Hokkaido, Japan. *Acta Endocrinol* 1965;50:161–76.
41. Kapil U, Singh P. Current status of urinary iodine excretion levels in 116 districts of India. *J Trop Pediatr* 2004;50: 245–7.
42. Kapil U, Sharma TD, Singh P, *et al.* Thirty years of a ban on the sale of noniodised salt: impact on iodine nutrition in children in Himachal Pradesh, India. *Food Nutr Bull* 2005;26:255–8.
43. Kapil U. Urinary iodine excretion levels amongst school-children in Haryana. *Indian Pediatr* 2009;46:57–97.
44. Marwaha RK, Tandon N, Desai A, *et al.* Iodine nutrition in upper socioeconomic school children of Delhi. *Indian Pediatr* 2010;47:335–58.
45. National Institute of Nutrition, Indian Council of Medical Research. Current Status of IDD in Select Districts of Different Regions of Country. Hyderabad: NIN Press, 2003, 11–28.
46. Langer P, Michajlovskij N. Effect of naturally occurring goitrogens on thyroid peroxidase and influence of some compounds on iodide formation during the estimation. *Endocrinol Exp* 1972;6:97–103.
47. Chandra AK. Goitrogens in food: cyanogenic and flavonoids containing plant foods in the development of goitre. In: Watson RR, Preedy VR (eds). *Bioactive Foods in Promoting Health: Fruits and Vegetables*. USA: Academic Press (imprint of Elsevier), 2010, 691–716.
48. Chandra AK. Iodine, thiocyanate and the thyroid. *Biochem Pharmacol* 2015;4:1–6.
49. VanEtten CH. Goitrogens. In: Liener IE (ed). *Toxic Constituents of Plant Foodstuffs*. New York and London: Academic Press, 1969, 103–34.
50. Wollman SH. Inhibition by thiocyanate of accumulation of radioiodine by thyroid gland. *Am J Physiol* 1962;203: 517–23.
51. Michot JL, Osty J, Nunez J. Regulatory effects of iodide and thiocyanate on tyrosine oxidation catalyzed by thyroid peroxidase. *Eur J Biochem* 1980;107:297–01.
52. Virion A, Deme D, Pommier J, *et al.* Opposite effect of thiocyanate on tyrosine iodination and thyroid hormone synthesis. *Eur J Biochem* 1980;112:1–7.
53. Chandra AK, Mukhopadhyay S, Lahiri D, *et al.* Goitrogenic content of Indian cyanogenic plant foods and their *in vitro* anti-thyroidal activity. *Indian J Med Res* 2004;119:180–5.
54. Delange F, Thilly C, Bourdoux P, *et al.* Influence of dietary goitrogens during pregnancy in humans on thyroid function of the newborn. In: Delange F, Itke FB, Ermans AM (eds). *Nutritional Factors Involved in the Goitrogenic Action of Cassava*. Ottawa: International Development Research Centre publication, 1982, 40–50.
55. Marwaha RK, Tandon N, Gupta N, *et al.* Residual goitre in the postiodation phase: iodine status, thiocyanate exposure and autoimmunity. *Clin Endocrinol* 2003;59:672–81.
56. Malik S. Together with prevention of food adulteration rules, notifications and commodity index. In: Malik S (ed). *Food Adulteration: Handbook of Food Adulteration and Safety*. Lucknow: Laws Eastern Book Company, 2012, 31–40.
57. National Family Health Survey III (NFHS-III). Deonar, Mumbai: International Institute for Population Sciences, 2006, 295–98.
58. Harach HR, Escalante DA, Onativia A, *et al.* Thyroid carcinoma and thyroiditis in a endemic goitre region before and after iodine prophylaxis. *Acta Endocrinol* 1985;108: 55–60.