

BRIEF REPORT

Vitamin B₁₂ Deficiency in Infancy: The Case for Screening

Gurpreet Singh, MD,^{1,2} Doan Le, MD, FAAP, FRCPC,^{2,3} Kareena Schnabl, MSc, PhD, FCACB,^{4,5} Michael T. Leaker, MD, FRCPC,^{2,3} MacGregor Steele, MD, FRCPC,^{2,3} and Rebecca L. Sparkes, MD, FRCPC, FCCMG^{6,7}

The classic principles put forth by Wilson and Jungner are often applied to determine the suitability of a condition for universal newborn screening. The three cases described here portray the harmful effects of vitamin B₁₂ deficiency in infancy. The challenges and opportunities of early recognition and treatment are highlighted. Screening newborns would allow early detection and prevention

of severe neurological damage in vitamin B₁₂-deficient infants and enable diagnosis of unrecognized maternal pernicious anemia in asymptomatic mothers. However, lack of standardized methodology and screening cutoffs present challenges to the use of current tandem mass spectrometry technologies for screening. *Pediatr Blood Cancer* 2016;63:740–742. © 2016 Wiley Periodicals, Inc.

Key words: early detection; infants; neurological effects; newborn screening; vitamin B₁₂ deficiency

INTRODUCTION

Vitamin B₁₂ (cobalamin) has important physiologic roles in hematopoiesis, intermediary metabolism, growth, and early brain development.[1,2] Clinical features of B₁₂ deficiency include macrocytic anemia lethargy, hypotonia, psychomotor delay, tremor, seizures, encephalopathy, irritability, weakness, diarrhea, stomatitis, glossitis, and failure to thrive.[1–4]

In children, B₁₂ deficiency is most commonly caused by inadequate dietary intake or impaired absorption. Rare causes include the inborn errors of intracellular cobalamin transport or metabolism.[1,2] B₁₂ deficiency due to decreased fetal stores and insufficient intake can occur in exclusively breastfed infants of mothers with undiagnosed B₁₂ deficiency. Women with vegetarian or vegan dietary practices or unrecognized pernicious anemia are at risk.[2,5]

The precise incidence of neonatal B₁₂ deficiency resulting from maternal deficiency is unknown.[2] In a retrospective American survey, Hinton et al. estimated it to occur in 0.88/100,000 births [95% confidence interval (CI) 0.60–1.26].[6] A higher incidence of 1:5,000 was detected by a newborn-screening pilot project in the Italian population.[7]

Because infants can present with severe neurological sequelae, the importance of early recognition has been emphasized in the literature.[3,4,8] However, B₁₂ deficiency may be overlooked due to its nonspecific clinical features.[3,4]

CASE DESCRIPTIONS

Case 1 was an exclusively breastfed 6-month-old male who presented with profound hypotonia, hyperreflexia, and regression in psychomotor development. Magnetic resonance imaging of the brain and electroencephalography were normal. Laboratory studies revealed anemia, an elevated mean corpuscular volume (MCV) and low serum B₁₂ level (Table I). His mother had low serum B₁₂ (49 pmol/l; reference range 155–700 pmol/l), high plasma total homocysteine (90.1 μmol/l; reference range 4.1–9.9 μmol/l), high plasma methylmalonic acid (MMA) (21.2 μmol/l; reference range 0.05–0.27 μmol/l), and positive anti-intrinsic factor (anti-IF) antibody. The treatment resulted in rapid and sustained improvement in B₁₂ levels. He regained the majority of his developmental milestones but remained mildly hypotonic at 10 months of age.

Case 2 was a 10-month-old breastfed female who presented with the parental concern of excessive somnolence. She had been meeting her developmental milestones at age-appropriate times. Complete physical and neurological examinations were unremarkable. An elevated MCV on complete blood count prompted the treating hematologist to order a serum B₁₂ level that was very low (Table I). Her mother had low B₁₂ (90 pmol/l), high plasma homocysteine (13.0 μmol/l), and positive anti-IF and antiparietal cell antibodies. The treatment was initiated, but she was lost to follow-up after the family relocated.

Case 3 was an exclusively breastfed 10-month-old male referred by his primary care physician for pallor and lethargy. His psychomotor development was delayed; he was unable to sit without support. Physical examination was unremarkable with the exception of axial hypotonia. Anemia and an elevated MCV prompted measurement of serum B₁₂ level that was very low (Table I). His mother had low B₁₂ (94 pmol/l), high plasma

Abbreviations: DBS, dried blood spot; IF, intrinsic factor; MCV, mean corpuscular volume; MMA, methylmalonic acid; NBS, newborn screening

¹Sections of Paediatric Haematology and Paediatric Oncology, Department of Paediatrics, Alberta Children's Hospital, Calgary, Alberta, Canada; ²Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada; ³Section of Paediatric Haematology, Department of Paediatrics, Alberta Children's Hospital, Calgary, Alberta, Canada; ⁴Newborn Screening and Biochemical Genetics Laboratory, University of Alberta Hospital, Edmonton, Alberta, Canada; ⁵Department of Laboratory Medicine and Pathology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada; ⁶Section of Clinical Genetics, Department of Paediatrics, Alberta Children's Hospital, Calgary, Alberta, Canada; ⁷Department of Medical Genetics, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

Conflict of interest: Nothing to declare.

Correspondence to: Gurpreet Singh, Sections of Paediatric Haematology and Paediatric Oncology, Department of Paediatrics, Alberta Children's Hospital, 2888 Shaganappi Trail NW, Calgary, Alberta T3B 6A8, Canada. E-mail: gurpreet.singh@albertahealthservices.ca

Received 29 May 2015; Accepted 10 October 2015

TABLE I. Laboratory Parameters of Index Cases at Presentation

Parameter (units) [reference range]	Case 1 (age 6 months)	Case 2 (age 10 months)	Case 3 (age 10 months)
Hb (g/l) [110–147]	100	72	112
MCV (fl) [71–90]	103	113	111
ANC (10 ⁹ /l) [0.6–5.1]	0.5	0.30	1.5
Platelets (10 ⁹ /l) [150–400]	152	339	547
Serum B ₁₂ (pmol/l) [155–700]	31	22	31
Plasma total homocysteine (μmol/l) [4.7–10.8]	156.5	130.7	72.5
Plasma MMA (μmol/l) [0.05–0.27]	19.5	15.6	37.8
C3/C2 on newborn DBS [<0.22]	0.26	0.14	0.22
C3 (μmol/l) on newborn DBS [<7.3]	6.20	3.63	5.21
C3–C16 on newborn DBS [No]	1.7	1.7	1.7

Hb, hemoglobin; MCV, mean corpuscular volume; ANC, absolute neutrophil count; MMA, methylmalonic acid; C3, propionylcarnitine; C2, acetylcarnitine; DBS, dried blood spot.

homocysteine (19.0 μmol/l), and positive antiparietal cell antibody. With the treatment, his psychomotor development had improved, but not normalized on follow-up at 16 months of age. His B₁₂ levels remained normal after treatment completion due to an improved diet.

DISCUSSION

The three cases of infantile-nutritional B₁₂ deficiency described herein presented clinically between 6 and 10 months of age, which is similar to other cases described in the literature.[3] All were breastfed by B₁₂-deficient mothers, and the two 10-month-olds had not been receiving age-appropriate complementary feeding. Although the Canadian Pediatric Society's current recommendations are to delay introduction of solid foods until 6 months of age, earlier introduction (around 4 months) of foods rich in B₁₂ may be considered to prevent deficiency.[9,10]

Nonspecific clinical findings of B₁₂ deficiency are often overlooked, leading to a delay in the diagnosis. Although the treatment reverses the hematological abnormalities quickly, the neurological manifestations may not completely resolve.[1,3,6,8,11, 12] Two of the three described cases had mild to moderate psychomotor delay at presentation. In both cases, follow-up

confirmed neurological improvement, but not normalization, after correction of the B₁₂ deficiency.

B₁₂-deficient adults may be clinically and hematologically normal.[11,13,14] In all three described cases, the mothers were asymptomatic but were found to have low B₁₂ levels and evidence of pernicious anemia on laboratory evaluation. Elevated MMA in blood and urine and, less commonly, total plasma homocysteine are sensitive, but nonspecific, markers of B₁₂ deficiency.[7,14]

Population-based screening may provide an opportunity for early recognition of, and intervention for, infants with B₁₂ deficiency.[7,11,15,16] Targeted screening of infants of vegan and vegetarian mothers is reasonable, but would only detect a minority of affected infants as many cases are due to impaired absorption of B₁₂ from dietary sources, regardless of maternal nutritional practices. An approach of B₁₂ screening in pregnancy by serum levels of B₁₂ and/or MMA may be very cost effective, easy to implement, and helpful in preventing severe pregnancy-related complications in addition to benefits for the newborn.[3,17]

Universal newborn screening (NBS) programs are jurisdiction specific (typically overseen by state/provincial or national public health departments) and hold potential for maximal case ascertainment. NBS also offers the additional benefit of

TABLE II. Suitability of Population-Based Screening for Vitamin B₁₂ Deficiency Based on Wilson and Jungner's Principles

Wilson and Jungner principle [20]	References supporting screening
1. The condition sought should be an important health problem.	[2–5,7–9]
2. There should be an accepted treatment for patients with recognized disease.	[2–5]
3. Facilities for diagnosis and treatment should be available.	[2–5]
4. There should be a recognizable latent or early symptomatic stage.	[2–5,7,8]
5. There should be a suitable test or examination.	[2–5]
6. The test should be acceptable to the population.	Not studied
7. The natural history of the condition, including development from latent to declared disease, should be adequately understood.	[2,7,8]
8. There should be an agreed-upon policy on whom to treat as patients.	[2–5]
9. The cost of case finding (including diagnosis and treatment of patients diagnosed) should be economically balanced in relation to possible expenditure on medical care as a whole.	Not investigated
10. Case finding should be a continuing process and not a “once and for all” project.	Not applicable

discovering asymptomatic, B₁₂-deficient mothers. Presymptomatic treatment could prevent the development of hematologic and neurologic dysfunction in these women.[14,17]

Considerable advances in NBS have occurred since the 1990s with the widespread adoption of tandem mass spectrometry.[18] With this technology, simultaneous quantification of multiple amino acid and acylcarnitine metabolites extracted from a dried blood spot (DBS) enables the diagnosis of certain inherited disorders of amino and fatty acid metabolism as a part of expanded NBS programs.

Propionylcarnitine (C3 acylcarnitine) accumulates in the setting of an inherited deficiency of the propionyl-CoA carboxylase or methylmalonyl-CoA mutase enzymes; both of which are targeted by NBS via C3 elevation. The mutase is adenosylcobalamin dependent; therefore, B₁₂ deficiency results in a functional deficiency of this enzyme and C3 elevation. Campbell et al. have argued that the C3 level alone is not sensitive or specific enough to detect all newborns with B₁₂ deficiency.[15] C3 levels may not be sufficiently high during the first few days of life when the DBS is collected.[15]

There is some evidence that the C3 to acetylcarnitine (C2) ratio may be better than C3 levels alone for detecting or identifying B₁₂ deficiency.[19] Sarafoglou et al. were successful in detecting B₁₂ deficiency in 11 cases by using lower cutoff levels for C3 (from >9.2 to >5.2 $\mu\text{mol/l}$), a C3–C2 (≥ 0.1) and C3 to palmitoylcarnitine (C16) ratios (≥ 2) followed by second-tier testing of MMA ($\geq 5 \mu\text{mol/l}$), and total homocysteine ($\geq 15 \mu\text{mol/l}$).[16] None went on to develop any symptoms or signs of B₁₂ deficiency with cyanocobalamin treatment. Mothers of eight of the 11 newborns were diagnosed with B₁₂ deficiency after their children were identified by NBS.

All three cases described in this report could have been identified if Sarafoglou et al.'s suggested cutoff for C3–C2 was used. The screening cutoffs suggested by Sarafoglou et al. are a good starting point, but refinement of the screening algorithm would require analysis of additional samples and careful consideration of the need for both maximal case ascertainment and minimization of false positives.

According to Wilson and Jungner's principles of screening, B₁₂ deficiency may be a suitable candidate for inclusion in universal NBS panels (Table II).[20] Challenges, including the

paucity of epidemiological studies defining the incidence of B₁₂ deficiency in newborns, and lack of consensus on laboratory methods to maximize sensitivity and specificity exist.

ACKNOWLEDGMENT

We gratefully acknowledge the patients described herein, and their parents.

REFERENCES

1. Black MM. Effects of vitamin B₁₂ and folate deficiency on brain development in children. *Food Nutr Bull* 2008;29:S126–S131.
2. Rasmussen SA, Fernhoff PM, Scanlon KS. Vitamin B₁₂ deficiency in children and adolescents. *J Pediatr* 2001;138:10–17.
3. Honzik T, Adamovicova M, Smolka V, Martin M, Eva H, Jiri Z. Clinical presentation and metabolic consequences in 40 breastfed infants with nutritional vitamin B₁₂ deficiency: What have we learned? *Eur J Paediatr Neurol* 2010;14:488–495.
4. Dobrozsi S, Flood VH, Panepinto J, Scott JP, Brandow A. Vitamin B₁₂ deficiency: The great masquerader. *Pediatr Blood Cancer* 2014;61:753–755.
5. Monagle PT, Tauro GP. Infantile megaloblastosis secondary to maternal vitamin B₁₂ deficiency. *Clin Lab Haematol* 1997;19:23–25.
6. Hinton CF, Ojodu JA, Fernhoff PM, Rasmussen SA, Scanlon KS, Hannon WH. Maternal and neonatal vitamin B₁₂ deficiency detected through expanded newborn screening: United States, 2003–2007. *J Pediatr* 2010;157:162–163.
7. Scolamiero E, Villani GR, Ingenito L, Pecce R, Albano L, Caterino M, di Gerolamo MG, Di Stefano C, Franzese L, Gallo G, Ruoppolo M. Maternal vitamin B₁₂ deficiency detected in expanded newborn screening. *Clin Biochem* 2014;47:312–317.
8. Demir N, Koc A, Üstöl, Peker E, Abuhandan M. Clinical and neurological findings of severe vitamin B₁₂ deficiency in infancy and importance of early diagnosis and treatment. *J Paediatr Child Health* 2013;49:820–824.
9. Critch JN. Canadian Paediatric Society; Nutrition and Gastroenterology Committee. Nutrition for healthy term infants, birth to six months: An overview. *Paediatr Child Health* 2013;8:206–207.
10. Greer FR, Sicherer SH, Burks AW. Effects of early nutritional interventions on the development of atopic disease in infants and children: The role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. *Pediatrics* 2008;121:183–191.
11. Michaud JL, Lemieux B, Ogier H, Lambert MA. Nutritional vitamin B₁₂ deficiency: Two cases detected by routine newborn urinary screening. *Eur J Pediatr* 1992;151:218–220.
12. Korenke GC, Hunneman DH, Eber S, Hanefeld F. Severe encephalopathy with epilepsy in an infant caused by subclinical maternal pernicious anaemia: Case report and review of the literature. *Eur J Pediatr* 2004;163:196–201.
13. Jain R, Kapil M, Gupta GN. MCV should not be the only criteria to order vitamin B₁₂ for anemia under evaluation. *Open J Gastroenterol* 2012;2:187–190.
14. Carmel R, Green R, Rosenblatt DS, Watkins D. Update on cobalamin, folate, and homocysteine. *Hematology Am Soc Hematol Educ Program* 2003;1:62–81.
15. Campbell CD, Ganesh J, Ficicioglu C. Two newborns with nutritional vitamin B₁₂ deficiency: Challenges in newborn screening for vitamin B₁₂ deficiency. *Haematologica* 2005;90:e119–e121.
16. Sarafoglou K, Rodgers J, Hietala A, Matern D, Kristi B. Expanded newborn screening for detection of vitamin B₁₂ deficiency. *JAMA* 2011;305:1198–1200.
17. Sande H, Jacquemyn Y, Karepouan N, Ajaji M. Vitamin B12 in pregnancy: Maternal and fetal/neonatal effects—A review. *Open J Obstet Gynecol* 2013;3:599–602.
18. Wilcken B, Wiley V. Fifty years of newborn screening. *J Paediatr Child Health* 2015;51:103–107.
19. Armour CA, Brebner A, Watkins D, Geraghty MT, Chan A, Rosenblatt DS. A patient with an inborn error of vitamin B₁₂ metabolism (cblF) detected by newborn screening. *Pediatrics* 2013;132:e257–e261.
20. Wilson JMG, Jungner G. Principles and practice of screening for disease. Geneva: World Health Organization; 1968.