
Relationship of Vitamin D Status with Muscle Mass and Muscle Strength in Young Indian Adults – Evidence from Andhra Pradesh Children and Parents Study Cohort





European Journal of Nutrition & Food Safety 5(5): 918-919, 2015, Article no.EJNFS.2015.355 ISSN: 2347-5641



SCIENCEDOMAIN international

www.sciencedomain.org

Relationship of Vitamin D Status with Muscle Mass and Muscle Strength in Young Indian Adults – Evidence from Andhra Pradesh Children and Parents Study Cohort

Bharati Kulkarni^{1*}, Hannah Kuper², Sanjay Kinra², M Sesha Charyulu¹, Yoav Ben-Shlomo³, George Davey Smith³, Shah Ebrahim² and K.V. Radhakrishna¹

¹National Institute of Nutrition, Hyderabad, India. ²London School of Hygiene and Tropical Medicine, London, UK. ³University of Bristol, Bristol, UK.

Article Information

DOI: 10.9734/EJNFS/2015/21163

Special Editors:

Lucie Bohac, Micronutrient Forum Secretariat, Canada. Klaus Kraemer, Director, Sight and Life, Basel, Switzerland.

Chief Edito

Prof. Hans Verhagen, Senior Scientific Advisor 'Nutrition and Food Safety', National Institute for Public Health and the Environment (RIVM), P.O.Box 1, 3720 BA, Bilthoven, The Netherlands.

Conference Abstract

Received 5th February 2015 Accepted 1st March 2015 Published 14th August 2015

ABSTRACT

Objectives: Positive relationship of vitamin D status with muscle mass and strength has been observed in studies from the developed countries but evidence from the developing countries is sparse. This study assessed the relationship of vitamin D status with muscle mass and muscle strength in rural young adults from Hyderabad, India.

Methods: The study participants (n=956; age 18-20 years; 42% women) were a part of Andhra Pradesh Children and Parents Study cohort which was established to assess the long term impact of early nutrition supplementation provided through a government programme. Their serum 25-hydroxyvitamin D was assessed using HPLC, appendicular skeletal muscle mass (ASM) was assessed using dual energy X-ray absorptiometry and grip strength was assessed using grip dynamometer.

Results: The participants were lean with average body mass index of 19.5 kg/m 2 . Prevalence of vitamin D deficiency (serum 25(OH) vitamin D3 < 20 ng/ml) was 33.6% in men and 51.4% in women. Vitamin D deficiency was associated with lower ASM (β (95% CI): - 0.38 (-0.72 to -0.05)

kg; p = 0.02) with a trend of lower muscle strength in unadjusted analyses. After adjustment for relevant confounders, the relationship of vitamin D deficiency with lower ASM (β (95% CI): -0.21 (-0.37 to -0.05) kg; p =0.01) persisted but not with lower grip strength.

Conclusions: Prevalence of vitamin D deficiency was high in these rural young adults. Vitamin D deficiency was associated with lower muscle mass but not with lower muscle strength in this cohort. Alleviation of vitamin D deficiency may improve muscle mass.

^{© 2015} Kulkarni et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.