

## Identification of vitamin B<sub>12</sub> deficiency in vegetarian Indians

Sadanand Naik\*, Namita Mahalle and Vijayshri Bhide

Pathology Laboratory, Deenanath Mangeshkar Hospital & Research Centre, Pune 411004, Maharashtra, India

(Submitted 4 May 2017 – Final revision received 13 August 2017 – Accepted 6 October 2017)

### Abstract

The prevalence of a sub-clinical vitamin B<sub>12</sub> deficiency in the vegetarians is high. Total serum vitamin B<sub>12</sub> concentration alone does not reliably reflect vitamin B<sub>12</sub> status. Holotranscobalamin (holo-TC) II is a bioactive B<sub>12</sub> fraction promoting specific uptake of B<sub>12</sub> by cells and the circulating concentration reflects the intake of B<sub>12</sub>, whereas total homocysteine (tHcy) indicates the metabolic ability. In this study, we investigated the diagnostic value of circulating holo-TC, B<sub>12</sub>, folate and homocysteine in vegetarians who were at risk of B<sub>12</sub> deficiency. B<sub>12</sub>-related biomarkers were measured in 119 young, healthy graduate vegetarians. None was folate deficient. As per reported definition, half were B<sub>12</sub> deficient; 70% of males and 50% of females had low plasma holo-TC concentrations; and 92% of males and half of females had hyperhomocysteinaemia. None had any clinical signs of B<sub>12</sub> deficiency. Receiver operating characteristic curve analysis demonstrated similar AUC at the B<sub>12</sub> concentration of 100 and 150 pmol/l when holo-TC (0.777 and 0.784) and homocysteine (0.924 and 0.928) were used as variables. Cut-off value of 100 pmol/l resulted in the highest sensitivity of 77.78% and specificity of 71.05% with a predictive value of 19.6 pmol/l for holo-TC and a sensitivity of 82.72% and specificity of 89.7% with a predictive value of 21.7 µmol/l for homocysteine. The combination of B<sub>12</sub>, holo-TC and tHcy improves the diagnostic accuracy at these cut-offs, and we suggest that for the young Indian vegetarians the cut-off for plasma B<sub>12</sub> and holotranscobalamin is 100 pmol/l and 19.6 pmol/l, respectively, and for homocysteine it is 17.6 (females) and 27 µmol/l (males) for identifying B<sub>12</sub> deficiency.

**Key words:** Vitamin B<sub>12</sub>: Holotranscobalamin: Folate: Homocysteine

Vitamin B<sub>12</sub> is essential for 1-C metabolism and cell division. Foods derived from animals are the main sources of vitamin B<sub>12</sub>. Strict vegetarians have limited sources of vitamin B<sub>12</sub> in their diet and therefore likely to have vitamin B<sub>12</sub> deficiency<sup>(1–3)</sup>. Reduced consumption of cobalamin from food or impaired intestinal absorption leads to severe deficiency when tissue stores of the vitamin are depleted. The clinical consequence of vitamin B<sub>12</sub> deficiency includes megaloblastic anaemia and progressive neurologic disease of the central and peripheral nervous system<sup>(4)</sup> and hyperhomocysteinaemia, a risk factor for CVD<sup>(5)</sup>. Early diagnosis of vitamin B<sub>12</sub> deficiency is useful to prevent irreversible neurological damage by cobalamin supplementation<sup>(6–8)</sup>. It has been observed that asymptomatic Indian lacto-vegetarians, who make up for more than half of the Indian population, had distinctly lower vitamin B<sub>12</sub> concentrations than non-vegetarians<sup>(9)</sup> and was confirmed by studies from different geographic regions of India. However, total plasma vitamin B<sub>12</sub> concentration may not reliably reflect vitamin B<sub>12</sub> status. To obtain more specificity and sensitivity in diagnosing vitamin B<sub>12</sub> deficiency, the concept of measuring holotranscobalamin (holo-TC) II, a transport protein, has aroused great interest. holo-TC is a biologically active vitamin B<sub>12</sub> fraction that promotes the aspecific uptake of vitamin B<sub>12</sub> by cells<sup>(10)</sup>.

Vitamin B<sub>12</sub> deficiency has been divided into four stages<sup>(11)</sup>. In stages I and II, indicated by a low plasma level of holo-TC, the plasma and cell stores become depleted. Stage III is characterised by increased plasma levels of total homocysteine (tHcy) and methylmalonic acid (MMA) in addition to lowered holo-TC. In stage IV, clinical signs become recognisable such as macroovalocytosis, elevated mean corpuscular volume (MCV) or lower Hb levels. Stage III of vitamin B<sub>12</sub> deficiency has been found in over 60% of vegetarians<sup>(12)</sup>. Thus, it is important to monitor vitamin B<sub>12</sub> status in this dietary group. Measurement of plasma vitamin B<sub>12</sub>, holo-TC, tHcy and MMA has been suggested for optimal monitoring of vitamin B<sub>12</sub> status in vegetarians<sup>(13)</sup>. In this study, we investigated the diagnostic cut-off values of plasma vitamin B<sub>12</sub>, holo-TC and functional marker, tHcy, of vitamin B<sub>12</sub> metabolism in Indian vegetarians.

### Methods

The study was conducted at Deenanath Mangeshkar Hospital and Research Centre. Young, healthy, postgraduates and staff members of the hospital and their relatives were invited to participate in the study. Self-explained information regarding the participants was captured by an interview, which included

**Abbreviations:** Hcy, homocysteine; holo-TC, holotranscobalamin; MMA, methylmalonic acid; tHcy, total homocysteine.

\* **Corresponding author:** Dr S. S. Naik, email sadu\_naik@rediffmail.com

age, vitamin supplementation, food habits and routine lifestyle. Study subjects were enrolled after the purpose and requirements of the study were explained. They were clinically examined for gross clinical signs of protein-energy under-nutrition and vitamin deficiencies (vitamin A, B-complex, C and D). Non-vegetarians and pregnant women were not included. Subjects with diabetes, cancer and those taking drugs known to influence vitamin B<sub>12</sub> absorption were also excluded. A total of 119 eligible participants (forty-six male and seventy-three female) were enrolled for the study. The study protocol was approved by the Hospital Ethical Committee and all the participants gave written informed consent (2015\_APR/SN/169).

### Experimental procedure

A volume of 10 ml of fasting blood sample was collected in EDTA vacutainers. Haematological parameters were measured on a five-part differential cell counter (Sysmex) and the remaining blood was centrifuged at 1500 *g* for 20 min. Separated plasma was stored for biochemical investigations at -20°C. Plasma holo-TC was measured using microparticle enzyme immunoassay. Microparticle enzyme intrinsic factor assay was used for the quantitative determination of plasma vitamin B<sub>12</sub>. Plasma folate and tHcy were measured by the fluorescence polarisation immunoassay technique. All four biomarkers were analysed on an AxSYM immunoassay analyzer (Abbott Laboratories)<sup>(17)</sup>. Plasma creatinine was measured using the alkaline picrate method with a Daytona analyser (Randox)<sup>(14)</sup>.

### Dietary intake data

A 24-h dietary recall questionnaire was administered to capture detailed information about all food, beverages and dietary supplements consumed in the past 24 h from midnight to midnight the previous day by an experienced nutritionist. Energy, protein, fat and vitamin B<sub>12</sub> intake was calculated using Dietsoft<sup>(15,16)</sup>.

**Definitions.** Folate and vitamin B<sub>12</sub> deficiency was defined as concentrations <2 ng/ml and <148 pmol/l, respectively; hyperhomocysteinaemia as plasma tHcy concentrations <15 µmol/l<sup>(17)</sup>; low holo-TC concentration as <35 pmol/l; anaemia as Hb concentration <120 g/l in females and <130 g/l in males; and macrocytosis as MCV > 100 fl<sup>(18)</sup>.

### Statistical analysis

The data are presented as medians and 25th and 75th percentiles. Between-group comparisons were calculated using Mann-Whitney *U* test and associations were tested using Pearson's correlation coefficient.

**Sample size justification.** Prevalence of vitamin B<sub>12</sub> deficiency in vegetarian Indians is found to be 70%<sup>(9)</sup>. Sample size was calculated using Buderer's method considering prevalence rate<sup>(19)</sup>. We selected sensitivity and specificity to be 80 and 90%, respectively. We chose clinically acceptable width of 95% CI for

sensitivity and specificity to be no more than 10%, significance ( $\alpha=0.05$ ). The sample size calculated was 88 for the expected sensitivity of 80% and 116 for the expected sensitivity of 90%. Taking maximum of both, we arrived at the estimate of 116.

We enrolled 119 subjects. Receiver operating characteristic (ROC) curves and AUC (with 95% CI) were used to measure the diagnostic accuracy of (cut-offs) vitamin B<sub>12</sub>, holo-TC and tHcy. ROC decision plots depicting sensitivity and specificity for 100 and 150 pmol/l concentration of vitamin B<sub>12</sub> using holo-TC (<35 pmol/l) and tHcy (>15 µmol/l) variables were determined to identify the cut-off values.

### Results

The participants had no clinical symptoms of vitamin B<sub>12</sub> deficiency (neurological symptoms such as paresthesia, weakness, gait abnormality, tingling of hands and feet or anaemia symptoms such as skin pallor, fatigue, shortness of breath and so on) and were not taking vitamin B<sub>12</sub> supplementation or any drugs known to influence vitamin B<sub>12</sub> absorption.

All subjects had normal plasma folate and creatinine concentrations (5.9, 6.6 ng/ml and 1.2, 0.95 mg/dl in males and females, respectively). Median plasma vitamin B<sub>12</sub>, holo-TC and tHcy concentrations were 146.5 and 164 pmol/l, 22.3 and 26.5 pmol/l, and 22.6 and 14.2 µmol/l in males and females, respectively (Table 1). In all, 39 and 30% had very low concentration (<100 pmol/l), 11 and 20% had low concentration (100–148 pmol/l), and 50 and 52% had normal plasma vitamin B<sub>12</sub> concentration (>148 pmol/l) in males and females, respectively. holo-TC was lower in three-fourth of the participants (22 and 24 pmol/l). A significant sex difference was found in plasma tHcy concentrations (22.6 µmol/l in males and 14.2 µmol/l in females,  $P=0.0001$ ), with no sex difference in holo-TC and vitamin B<sub>12</sub> concentrations (Table 1). Group-wise distribution of vitamin B<sub>12</sub> (<100, 100–150 pmol/l) was strongly related to plasma holo-TC and inversely to tHcy concentrations in both males and females (Table 2,  $P<0.0001$ ). At normal level of vitamin B<sub>12</sub> (>148 pmol/l), the median concentrations of plasma holo-TC concentrations were 27.7 and 29.8 pmol/l in vegetarian males and females, which are lower than those found in non-vegetarians (Enexo). Between plasma vitamin B<sub>12</sub> concentrations of 113 and 122 pmol/l, nineteen participants had normal holo-TC (34–52 pmol/l) with higher tHcy (34 µmol/l) concentrations, which is unexplainable with existing cut-offs (Table 3). ROC analysis demonstrated similar AUC at the vitamin B<sub>12</sub> concentration of 100 and 150 pmol/l when holo-TC (0.784 and 0.777, respectively) and homocysteine (Hcy) (0.928 and 0.924, respectively) were used as variables. Cut-off value of 100 pmol/l resulted in the highest sensitivity (77.78%) with acceptable specificity (71.05%) with a predictive value of 19.6 pmol/l for holo-TC and a sensitivity of 82.72% and specificity of 89.47% with a predictive value of 21.2 µmol/l for tHcy (Fig. 1 and 2).

### Discussion

Our data support the concept that the measurement of plasma holo-TC and tHcy along with vitamin B<sub>12</sub> provides a better index of cobalamin status than the measurement of vitamin B<sub>12</sub> alone<sup>(20)</sup>. Plasma holo-TC and tHcy are both sensitive markers

**Table 1.** Baseline characteristics and biochemistry of the participants (male and female) (Medians and 25th, 75th percentiles)

	Male (n 46)		Female (n 73)		P*
	Median	25th, 75th percentiles	Median	25th, 75th percentiles	
Age (years)	29.0	25.5, 33.0	28.2	27, 32.5	NS
Energy intake (kJ/d)	10006	9372, 10334	9455	8786, 10446	0.042
Protein intake (g/d)	68	60, 74	62	58, 70	NS
Fat intake (g/d)	48	40, 55	47	40, 57	NS
Folate intake (µg/d)	268	220, 310	275	230, 300	NS
Vitamin B <sub>12</sub> intake (µg/d)	1.65	1.3, 1.90	1.80	1.4, 2.0	0.04
Hb (g/l)	140	134, 148	124	117, 132	0.000
Anaemia (%)		4.0		26	
Mean corpuscular volume (fl)	85.8	82.0, 89.5	82.5	78.0, 85.0	0.001
Macrocytic (%)		12		15	
Creatinine (mg/dl)	1.2	0.9, 1.3	0.95	0.8, 1.15	0.001
Vitamin B <sub>12</sub> (pmol/l)	146	84, 244	164	100, 288	NS
Vitamin B <sub>12</sub> deficiency (%)		77		50	
holo-TC (pmol/l)	22.3	12.3, 31.4	24.4	15.5, 41.6	NS
Low holo-TC (%)		78		75	
Folate (ng/ml)	5.9	4.2, 14.9	6.6	4.0, 15.2	NS
Folate deficiency (%)		0		0	
tHcy (µmol/l)	22.6	13.9, 52.0	14.2	9.9, 24.4	0.0001
Hyperhomocysteinaemia (%)		92		50	

holo-TC, holotranscobalamin; tHcy, total homocysteine.

\* Difference between male and female.

**Table 2.** Sex difference in total homocysteine (tHcy) at different vitamin B<sub>12</sub> concentrations (Medians and 25th–75th percentiles)


Plasma vitamin B <sub>12</sub> (pmol/l)	Sex	n	tHcy (µmol/l)		P (sex difference)	holo-TC (pmol/l)	
			Median	25th, 75th percentiles			
≤100	Male	18	50	46, 55	0.0001	13.2	NS
	Female	20	26.4	21.3, 40.4		14.6	
100–150	Male	5	23.8	21.4, 27	0.366	21.4	NS
	Female	15	20.9	15.6, 25.8		23.4	
>150	Male	23	13.9	12.0, 18.2	0.0068	31.3	NS
	Female	38	10.1	8.9, 12.6		35.3	

holo-TC, holotranscobalamin.

**Table 3.** Plasma vitamin B<sub>12</sub> concentrations and corresponding concentrations of plasma holotranscobalamin (holo-TC) and total homocysteine (tHcy)

Vitamin B <sub>12</sub> (pmol/l)	Median		Participants			Vitamin B <sub>12</sub> status
	holo-TC (pmol/l)	tHcy (µmol/l)	Male (n 46)	Female (n 73)	Total (n 119)	
314	46.75	9.64	4	14	18	Normal (B <sub>12</sub> > 148 pmol/l)
266	36.3	11.2	5	9	14	
175	43.0	11.4	2	5	7	
174	24.9	12.49	3	5	8	
170	20.7	17.7	4	5	9	
122	41.65	34.06	2	5	7	Metabolic deficiency tHcy >15 µmol/l, holo-TC > 35 pmol/l)
113	52.7	34.06	3	10	13	
102	15.6	25.7	14	9	23	Deficiency B <sub>12</sub> <200 pg/ml, tHcy > 15 µmol/l, holo-TC > 35 pmol/l
50	10.5	52.0	9	11	20	

holo-TC, holotranscobalamin; tHcy, total homocysteine.




**Fig. 1.** Plasma vitamin B<sub>12</sub> at concentrations of 150 and 100 pmol/l were used for analysis. Metabolic deficiency was defined as plasma holotranscobalamin (holo-TC) < 35 pmol/l in 119 vegetarian Indians. Predictive values of plasma holo-TC were 24.58 and 19.6 µmol/l, respectively, with similar sensitivity and specificity at both 150 and 100 pmol/l of vitamin B<sub>12</sub>. ROC, receiver operating characteristic.

for cobalamin status<sup>(21)</sup>. The lowest median levels of holo-TC were observed in the low vitamin B<sub>12</sub> concentration group, followed by the 100–150 pmol/l group and then by the normal group (>148 pmol/l): 13.2, 21.4 and 31.3 pmol/l, respectively (*P* = 0.0001). At similar plasma cobalamin concentrations, men had different metabolic effects compared with women. The use of holo-TC and tHcy results enables the differentiation between storage, depletion and functional vitamin B<sub>12</sub> deficiency.

Fedosov<sup>(22)</sup> reported that a combination of vitamin B<sub>12</sub>, holo-TC, MMA and Hcy (cB<sub>12</sub>) biomarkers is a reliable diagnostic tool. Fedosov *et al.*<sup>(23)</sup> derived equations that combined two, three or four biomarkers into one diagnostic indicator and provided a guidance for treatment. They suggested that adults having plasma vitamin B<sub>12</sub> levels between 116 and 119 pmol/l, holo-TC between 8.4 and 20 pmol/l, and Hcy between 19.2 and 51 µmol/l be grouped as low vitamin B<sub>12</sub> status and adults with B<sub>12</sub> between 119 and 186 pmol/l, holo-TC between 20 and 37 pmol/l, and Hcy between 13.6 and 19.2 µmol/l as transitional vitamin B<sub>12</sub> deficiency. A Brito *et al.*<sup>(24)</sup> stated that cB<sub>12</sub> could only detect improved neurophysiological function in asymptomatic Chilean elderly with poor vitamin B<sub>12</sub> status and also

suggested to identify functional indicators of sub-clinical vitamin B<sub>12</sub> deficiency.

This is the first study to investigate the diagnostic value of circulating plasma vitamin B<sub>12</sub>, holo-TC and tHcy (a functional marker) concentrations associated with 1-C metabolism in Indian vegetarians. We found that at similar circulating concentrations of plasma vitamin B<sub>12</sub> and holo-TC, there is a sex difference in plasma tHcy concentrations, with men having higher tHcy concentrations than women (*P* < 0.0001). The marked sex difference<sup>(17)</sup> in plasma tHcy concentrations is confirmed in this study. The difference suggests a higher threshold for supplementation of vitamin B<sub>12</sub> to improve reproductive and cardiovascular outcomes<sup>(25)</sup>. The sex difference is not significant in participants whose plasma vitamin B<sub>12</sub> concentrations are between 105 and 148 pmol/l. The strong inverse relation between plasma vitamin B<sub>12</sub> and tHcy and direct relation with holo-TC<sup>(17)</sup> concentrations are confirmed in this study. Although age and sex reference intervals of tHcy have been established<sup>(25,26)</sup>, they are usually ignored in reporting tHcy levels, because of dual biochemical origin (B<sub>12</sub> and folate). Elevated Hcy in plasma has been very often used as



ROC analysis using tHcy as a variable (>15 μmol/l); see Fig. 2

Vitamin B <sub>12</sub> cut-off (pmol/l)	AUC	Sensitivity (%)	Specificity (%)	Predictive value (tHcy, μmol/l)
100	0.928 (95 % CI: 0.865, 0.967) <i>P</i> <0.0001	82.72	89.47	21.2
150	0.924 (95 % CI: 0.860, 0.964) <i>P</i> <0.0001	91.8	79.31	18.56

**Fig. 2.** Plasma vitamin B<sub>12</sub> at 150 and 100 pmol/l were used as variables. Metabolic deficiency was defined as total homocysteine (tHcy) >15 μmol/l in 119 vegetarian Indians. Predictive values of plasma tHcy were 18.56 and 21.2 μmol/l, respectively, with better specificity (89.47 %) at 100 pmol/l of vitamin B<sub>12</sub>. ROC, receiver operating characteristic.

a biomarker, but its relationship to the molecular mechanisms of disease has not been established<sup>(27)</sup>.

In this study, we aimed to meet two criteria: (a) vitamin B<sub>12</sub> absorption capacity in Indian vegetarians and (b) optimum plasma vitamin B<sub>12</sub> concentration required for methylation of Hcy. At normal plasma vitamin B<sub>12</sub> levels (>150 pmol/l), median holo-TC concentrations were 31.3 and 35.3 pmol/l in men and women, respectively, thereby attaining normal tHcy concentrations (13.9 and 10.17 μmol/l, respectively).

A cohort of 100 known patients with CVD and sixty-three normal healthy subjects (median age 44 years) were examined for their vitamin B<sub>12</sub> status in a case-control study in Pune. Median plasma vitamin B<sub>12</sub>, holo-TC and tHcy concentrations were 160 pmol/l, 24 pmol/l and 19.7 μmol/l, respectively, in normal subjects. They stated that hyperhomocysteinaemia and elevated MMA were due to vegetarianism<sup>(18)</sup>. Similarly, another report from Pune, wherein subjects with low plasma vitamin B<sub>12</sub> levels were studied, showed very low plasma holo-TC concentrations (7.7 (sd 4.2) and 9.8 (sd 8.7) pmol/l in men and women, respectively), with tHcy concentrations of 29.2 (sd 19.2) and 15.3 (sd 8.3) μmol/l<sup>(28)</sup>. Low vitamin B<sub>12</sub> concentration (median 110 pmol/l) and hyperhomocysteinaemia (>15 μmol/l) have been reported to be common in Indian men, particularly

in vegetarians and urban middle-class residents. Most of the participants from these studies were vegetarians. In the study by Naik *et al.*<sup>(17)</sup>, young Indian vegetarian subjects with low plasma vitamin B<sub>12</sub> status (<148 pmol/l) were found to have holo-TC and tHcy concentrations of 14.4 pmol/l and 31.9 μmol/l, respectively, and the subjects with normal status (>200 pmol/l) were found to have holo-TC and tHcy concentrations of 27.7 pmol/l and 11.9 μmol/l, respectively. Most of the Indian studies did not use plasma holo-TC measurements. A Dutch study reported 49 (8–388) pmol/l of plasma holo-TC concentrations in healthy subjects with corresponding vitamin B<sub>12</sub> concentrations of 217 (119–1210) pmol/l<sup>(29)</sup>. A study from USA reported plasma holo-TC concentration of 85 (sd 48) pmol/l for 495 (sd 119) pmol/l of vitamin B<sub>12</sub> concentrations in healthy volunteers<sup>(30)</sup>. In both these studies the participants were non-vegetarians with high plasma holo-TC concentrations and the referred cut-offs may not be appropriate for vegetarians.

In healthy individuals, all four biomarkers (plasma vitamin B<sub>12</sub>, holo-TC, tHcy and MMA) had a strong relation to vitamin B<sub>12</sub> intake, with steady-state concentrations at a daily intake of 4–7 μg vitamin B<sub>12</sub><sup>(31)</sup>. These studies suggest that all four markers may be useful for monitoring a population's vitamin B<sub>12</sub> status over time.



Carmel<sup>(32)</sup> categorised available biomarkers as those that directly measured plasma vitamin B<sub>12</sub> and those that measured metabolites that accumulated with inadequate amounts of vitamin B<sub>12</sub>. Plasma holo-TC and vitamin B<sub>12</sub> measured circulating vitamin B<sub>12</sub> concentrations. These two therefore reflected the broad vitamin B<sub>12</sub> status from high risk of severe deficiency to adequacy. Miller *et al.*<sup>(33)</sup> have stated that holo-TC and total vitamin B<sub>12</sub> have equal diagnostic accuracy in screening for metabolic vitamin B<sub>12</sub> deficiency. Measurement of both holo-TC and total vitamin B<sub>12</sub> provided better screen for vitamin B<sub>12</sub> deficiency than either assay alone. According to Green<sup>(34)</sup>, low vitamin B<sub>12</sub> status was indicated by values lower than the reference range (for vitamin B<sub>12</sub> <148 pmol/l; for holo-TC <35 pmol/l), whereas for indirect measures of metabolites (MMA or tHcy) low vitamin B<sub>12</sub> status measures were indicated by a level above the upper limit of the reference range (for MMA >260 nmol/l; for tHcy >12 µmol/l). However, Valente *et al.*<sup>(35)</sup> suggested a diagnostic strategy using holo-TC as the front-line test. The cut-offs for deficiency were defined as 20 pmol/l for holo-TC and 123 pmol/l for serum vitamin B<sub>12</sub> after studying employees and medical students of a local hospital at Dundee, UK.

### Conclusions

holo-TC levels may prove most useful if the aim is to monitor a population with a borderline sub-optimal vitamin B<sub>12</sub> supply. In contrast, total vitamin B<sub>12</sub> may be superior if the goal is to monitor a possible surplus load of vitamin. We advocate a diagnostic cut-off level of plasma vitamin B<sub>12</sub> (105 pmol/l), holo-TC (22.6 pmol/l) and Hcy (17.6 µmol/l for females and 27.0 µmol/l for males) in vegetarian Indian population. In addition to this, we recommend that vegetarians do take a supplement of vitamin B<sub>12</sub> to ensure adequate supply of the micronutrient. This would be of particular importance for females in reproductive age, to prevent the risks associated with maternal–fetal vitamin B<sub>12</sub> deficiency.

The limitation of the study is that plasma MMA has not been measured. However, if none of the participants is folate deficient, the measurement of Hcy shall indicate vitamin B<sub>12</sub> status.

### Acknowledgements

This research received no specific grant from any funding agency, commercial or not-for-profit sectors. Dr Dhanajay Kelkar, Medical Director, extended all the help. Our thanks are due to Swati, Swapnali and Sandeep and Abbott Laboratories for free kits.

S. N. designed the study. V. B. and N. M. prepared the manuscript draft. S. N. prepared the final draft.

The authors declare that there are no conflicts of interest.

### References

- Antony AC (2003) Vegetarianism and vitamin B<sub>12</sub> deficiency. *Am J Clin Nutr* **78**, 3–6.
- Dhopeshwarkar GA, Trivedi JC, Kulkarni BS, *et al.* (1956) The effect of vegetarianism and antibiotics upon proteins and vitamin B<sub>12</sub> in the blood. *Br J Nutr* **10**, 105–110.
- Jathar VS, Inamdar-Deshmukh AB, Rege DV, *et al.* (1975) Vitamin B<sub>12</sub> and vegetarianism in India. *Acta Hematol* **53**, 90–99.
- Obeid R, Giesel J, Schorr T, *et al.* (2002) The impact of vegetarianism on some hematological parameters. *Eu J Hematol* **69**, 275–279.
- Refsum H, Smith AD, Ueland PM, *et al.* (2001) Facts and recommendations about total homocysteine determinations: an expert opinion. *Br J Nutr* **85**, Suppl. 2, 5109–5113.
- Herrmann W, Obeid R, Schorr H, *et al.* (2003) Functional vitamin B<sub>12</sub> deficiency and determination of holo-transcobalamin in population at risk. *Clin Chem Lab Med* **41**, 1478–1488.
- Bhate V, Deshpande S, Bhat D, *et al.* (2008) Vitamin B<sub>12</sub> status of pregnant Indian women and cognitive function in their 9 yr old children. *Food Nutr Bull* **29**, 249–254.
- Weir DC & Scott JM (1999) Brain function in the elderly: role of vitamin B<sub>12</sub> and folate. *Br Med Bull* **55**, 669–682.
- Yajnik CS, Deshpande SS, Lubree HG, *et al.* (2006) Vitamin B<sub>12</sub> deficiency and hyperhomocysteinemia in rural and urban Indians. *J Assoc Physicians India* **54**, 775–782.
- Nexo E & Hoffman-Lücke E (2011) Holotranscobalamin, a marker of vitamin B-12 status: analytical aspects and clinical utility. *Am J Clin Nutr* **94**, 359S–365S.
- Herbert V (1994) Staging vitamin B-12 (cobalamin) status in vegetarians. *Am J Clin Nutr* **59**, 1213S–1232S.
- Herrmann W & Giesel J (2002) Vegetarian lifestyle and monitoring vitamin B<sub>12</sub> status. *Clin Chim Acta* **326**, 47–59.
- Lindenbaum J, Savage DG, Stabler SP, *et al.* (1990) Diagnosis of cobalamin deficiency: II. Relative sensitivities of serum cobalamin, methylmalonic acid and total homocysteine concentrations. *Am J Hematol* **34**, 99–107.
- Tulip Group. Creatinine kit (alkaline picrate method). [http://www.tulipgroup.com/Coral\\_New2/html/pack\\_inserts/Creatinine%20Kit%20Alkaline%20Picrate%20Method.pdf](http://www.tulipgroup.com/Coral_New2/html/pack_inserts/Creatinine%20Kit%20Alkaline%20Picrate%20Method.pdf)
- Kanade AN, Rao S, Kelkar RS, *et al.* (2008) Maternal nutrition and birth size among urban affluent and rural women in India. *Am J Clin Nutr* **27**, 137–145.
- Gopalan C, Rama Sastri BV & Balasubramanian SC (2007) Dietsoft software version 1.1.7. Developed by invincible, iDeAS, *Nutritive Value of Indian Foods*, revised ed. Hyderabad: NIN, ICMR. [www.invincibleiDeas.com](http://www.invincibleiDeas.com)
- Naik S, Bhide V, Babhulkar A, *et al.* (2013) Daily milk intake improves vitamin B<sub>12</sub> status in young vegetarian Indians: an intervention trial. *Nutr J* **12**, 136–144.
- Refsum H, Yajnik CS, Gadkari M, *et al.* (2001) Hyperhomocysteinemia and elevated methylmalonic acid indicate a high prevalence of cobalamin deficiency in Asian Indians. *Am J Clin Nutr* **74**, 233–241.
- Buderer NM (1996) Statistical methodology 1. Incorporating the procedure of disease into the sample size calculation for sensitivity and specificity. *Acad Emerg Med* **3**, 895–900.
- Lysne V, Strand E, Svingen GFT, *et al.* (2016) Peroxisome proliferator-activated receptor activation is associated with altered plasma one-carbon metabolites and B-vitamin status in rats. *Nutrients* **8**, 26.
- Solomon LR (2015) Functional cobalamin (vitamin B<sub>12</sub>) deficiency: role of advanced age and disorders associated with increased oxidative stress. *Eur J Clin Nutr* **69**, 687–692.
- Fedosov SN (2013) Biochemical markers of vitamin B<sub>12</sub> deficiency combined in one diagnostic parameter: the age dependence and associations with cognitive function and blood haemoglobin. *Clin Chim Acta* **422**, 47–53.
- Fedosov SN, Brito A, Miller JW, *et al.* (2015) Combined indicator of vitamin B<sub>12</sub> status: modifications for missing

- biomarkers and folate status and recommendations for revised cut-points. *Clin Chim Lab Med* **53**, 1215–1225.
24. Brito A, Verdugo R, Hertrampf E, *et al.* (2016) Vitamin B<sub>12</sub> treatment of asymptomatic, deficient, elderly Chileans improves conductivity in myelinated peripheral nerves but high serum folate impairs vitamin B<sub>12</sub> response assessed by combined indicator of vitamin B<sub>12</sub> status. *Am J Clin Nutr* **103**, 250–257.
  25. Naik S, Joglekar C, Bhat D, *et al.* (2011) Marked gender difference in plasma total homocysteine concentrations in Indian adults with low vitamin B<sub>12</sub>. *Int J Vitam Nutr Res* **81**, 306–316.
  26. Hannibal L & Blom HJ (2016) Homocysteine and disease: causal associations or epiphenomenons? *Mol Aspects Med* **53**, 36–42.
  27. Hannibal L, Lysne V, Monsen ALB, *et al.* (2016) Biomarkers and algorithms for the diagnosis of vitamin B<sub>12</sub> deficiency. *Front Mol Biosci* **3**, 27–42.
  28. Bhat DS, Thuse NV, Lubree HG, *et al.* (2009) Increased plasma holotranscobalamin can be used to assess vitamin B<sub>12</sub> absorption in individuals with low plasma vitamin B<sub>12</sub>. *J Nutr* **23**, 2119–2123.
  29. Bor MV, Lydeking-Olsen E, Moller J, *et al.* (2009) A daily intake of 6 ug vitamin B<sub>12</sub> appears to saturate all the vitamin B<sub>12</sub> related variables in Danish postmenopausal women. *Am J Clin Nutr* **89**, 1078–1087.
  30. Wokes F, Badenoch J & Sinclair HM (1955) Human dietary deficiency of vitamin B<sub>12</sub>. *Am J Clin Nutr* **3**, 375–382.
  31. Bor MV, von Castel-Roberts KM, Kauwell GPA, *et al.* (2010) Daily intake of 4–7 ug dietary vitamin B<sub>12</sub> is associated with a steady concentrations of vitamin B<sub>12</sub> related biomarkers in a healthy young population. *Am J Clin Nutr* **91**, 571–577.
  32. Carmel R (2011) Biomarkers of cobalamin status in the epidemiologic applications: a critical over view of context, applications and performance characteristics of cobalamin, methylmalonic acid and holotranscobalamin. *Am J Clin Nutr* **94**, 348S–358S.
  33. Miller W, Garrod MG, Rockwood AL, *et al.* (2006) Measurement of both total vitamin B<sub>12</sub> and holotranscobalamin singly and in combination. *Clin Chem* **52**, 278–285.
  34. Green R (2011) Indicators for assessing folate and vitamin B<sub>12</sub> status and for monitoring the efficiency of intervention strategies. *Am J Clin Nutr* **94**, 667S–672S.
  35. Valente E, Scott JM, Cunningham C, *et al.* (2011) Diagnostic accuracy of holotranscobalamin, methylmalonic acid, serum cobalamin, and other indicators of tissue vitamin B<sub>12</sub> status in the elderly. *Clin Chem* **57**, 860–863.