

Supporting Information

**Impact of roadside tree lines on indoor
concentrations of traffic-derived particulate
matter**

Barbara A. Maher*, Imad A.M. Ahmed, Brian Davison, Vassil
Karloukovski and Robert Clarke

Centre for Environmental Magnetism & Palaeomagnetism,
Lancaster Environment Centre, Lancaster University, Lancaster,
LA1 4YQ, United Kingdom

*Corresponding Author: E-mail: b.maher@lancaster.ac.uk; Tel; +44 (0) 1524 510268; Fax: +44
(0)1524 593985.

This document contains 6 pages and 4 Figures

Instrumentation details, magnetic measurements.

The ‘saturation’ remanence ($\text{SIRM}_{0.3\text{T}}$) was acquired by subjecting the screen swabs and birch tree leaf samples to an applied dc magnetic field of 0.3 T generated using a Molspin pulse magnetizer. All leaf remanence values were measured using a Molspin Minispin magnetometer (sensitivity level $\sim 0.1 \times 10^{-8} \text{ Am}^2$) and normalized for upper leaf surface area. The magnetometer was calibrated routinely (i.e. after \sim ten sample measurements) against a magnetically-stable, independently-calibrated laboratory rock specimen. All magnetic measurements were made at the Centre for Environmental Magnetism & Palaeomagnetism, Lancaster University.




Figure S1. Meteorological data (wind speed, temperature, rain) from Hazelrigg Weather Station, and PM_{10} ($\mu\text{g m}^{-3}$) measured inside the two monitored houses 31 and 25, from May 9 – 23 2013.




Figure S2. Traffic count data and indoor PM₁₀ concentrations (hourly sums) from a) house 25 and b) house 31.




Figure S3. PM₁₀ concentration inside house 31 from March 27 – 31 2013, showing the weekday PM₁₀ peaks associated with rush hour traffic flows, in contrast to the very low PM₁₀ concentrations during Good Friday (29/3/2013, a public holiday in the U.K.) and the following Easter Saturday and Sunday.

Leaf sampling protocol for SEM/EDX.

At the end of the experiment, at least 20 leaves were collected from each tree from the road-distal and -proximal sides for both magnetic and SEM analyses. We selected 4 leaves from each tree organised in two groups: two leaves collected from the side facing the main road and two from the side facing the houses. Then we randomly selected 4 leaves from each group for analysis by SEM.




Figure S4. Compilation of SEM-EDXA data from analyses (18) of particle aggregates deposited on the birch leaves. The weight% represents the weight per cent concentration of the element in the analysed region of the sample.