



Unravelling sex-specific BPA toxicokinetics in children using a pediatric PBPK model

Deepika Deepika^a, Raju Prasad Sharma^a, Marta Schuhmacher^a, Amrit Kaur Sakhi^b, Cathrine Thomsen^b, Leda Chatzi^c, Marina Vafeiadi^c, Joane Quentin^d, Remy Slama^d, Regina Grazuleviciene^e, Sandra Andrusaitytė^e, Dagmar Waiblinger^f, John Wright^f, Tiffany C. Yang^f, Jose Urquiza^{g,h,i}, Martine Vrijheid^{g,h,i}, Maribel Casas^{g,h,i}, José L. Domingo^j, Vikas Kumar^{a,k,*}

^a Environmental Engineering Laboratory, Departament D' Enginyeria Quimica, Universitat Rovira I Virgili, Av. Països Catalans 26, 43007, Tarragona, Catalonia, Spain

^b Norwegian Institute of Public Health, Oslo, Norway

^c Department of Social Medicine, Faculty of Medicine, University of Crete, Heraklion, Greece

^d Team of Environmental Epidemiology, IAB, Institute for Advanced Biosciences, Inserm, CNRS, CHU-Grenoble-Alpes, University Grenoble-Alpes, CNRS, Grenoble, France

^e Department of Environmental Sciences, Vytautas Magnus University, Kaunas, Lithuania

^f Bradford Institute for Health Research, Bradford Teaching Hospitals NHS Foundation Trust, Bradford, UK

^g ISGlobal, Barcelona, Spain

^h Universitat Pompeu Fabra (UPF), Barcelona, Spain

ⁱ Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Madrid, Spain

^j Laboratory of Toxicology and Environmental Health, School of Medicine, IISPV, Universitat Rovira I Virgili, Reus, Spain

^k IISPV, Hospital Universitari Sant Joan de Reus, Universitat Rovira I Virgili, Reus, Spain

ARTICLE INFO

Keywords:

Pediatric PBPK
Pharmacokinetic
Sex-specific risk
Children cohort
Endocrine disruptor
Bisphenol A

ABSTRACT

Bisphenol A (BPA) is a widely known endocrine disruptor (ED) found in many children's products such as toys, feeding utensils, and teething rings. Recent epidemiology association studies have shown postnatal BPA exposure resulted in developing various diseases such as diabetes, obesity, and neurodegeneration, etc., later in their lives. However, little is known about its sex-specific metabolism and consequently internal exposure. The aim of this study was to develop a sex-specific pediatric physiologically based pharmacokinetic model (PBPK) for BPA to compare their toxicokinetic differences. First, the published adult PBPK model was re-validated, and then this model was extended by interpolation to incorporate pediatric sex specific physiological and biochemical parameters. We used both the classical body weight and ontogeny-based scaling approach to interpolate the metabolic process. Then, the pharmacokinetic attributes of the models using the two-scaling approach mentioned above were compared with adult model. Further, a sex-specific PBPK model with an ontogeny scaling approach was preferred to evaluate the pharmacokinetic differences. Moreover, this model was used to reconstruct the BPA exposure from two cohorts (Helix and PBAT Cohort) from 7 EU countries. The half-life of BPA was found to be almost the same in boys and girls at the same exposure levels. Our model estimated BPA children's exposure to be about 1500 times higher than the tolerable daily intake (TDI) recently set by European Food Safety Authority (EFSA) i.e., 0.04 ng/kg BW/day. The model demonstrated feasibility of extending the adult PBPK to sex-specific pediatric, thus investigate a gender-specific health risk assessment.

1. Introduction

Humans are being exposed to the bisphenol A (BPA) through plastic

food, beverage containers, thermal receipts, medical devices, canned food, and sealant, with detectable levels occurring in urine worldwide (Harley et al., 2013; Braun and Hauser, 2011; Bushnik et al., 2010). Several human studies have reported BPA exposure is associated with

* Corresponding author. Environmental Engineering Laboratory, Departament d' Enginyeria Quimica, Universitat Rovira i Virgili, Av. Països Catalans 26, 43007, Tarragona, Catalonia, Spain.

E-mail address: vikas.kumar@urv.cat (V. Kumar).

<https://doi.org/10.1016/j.envres.2022.114074>

Received 11 July 2022; Received in revised form 3 August 2022; Accepted 4 August 2022

Available online 19 August 2022

0013-9351/© 2022 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

3. Results

3.1. Age dynamic based adult PBPK model showed good accuracy

The refined adult PBPK model predictions of free BPA and its metabolite concentration in plasma were compared with plasma data from the 12 healthy volunteers after a single oral administration, obtained from Thayer et al. (2015) (see Supp Fig. S8). The predictive accuracy of the model was calculated by the Pearson correlation, which showed that the simulated data fits well with the observed data, within a 2-fold deviation from observed values with r value ≥ 0.95 and p value less than 0.05 (Supp file, Fig. S9).

3.2. An ontogeny scaling based pediatric PBPK model showed the critical differences in pharmacokinetic characteristics vs classical body-weight scaling

Pharmacokinetic parameters were calculated and compared with the adult PBPK and the two different pediatric PBPK models: 1) the classical body-weight scaling and 2) the ontogeny scaling. The plasma concentration of free BPA for the same amount of exposure was comparatively higher in children with ontogeny scaling than in adults, and however, the body-weight-based scaling predicted free BPA for the same amount of exposure almost as much as in adults (Fig. S10 in supplementary file). However, BPAG plasma concentration were almost similar in the adult and two pediatric models but BPAS plasma concentration was

significantly higher in children with ontogeny-based scaling than with body-weight based scaling and the adult PBPK model (Supplementary Fig. S10). Fig. 2 showed pharmacokinetic parameters like AUC, $t_{1/2}$, bioavailability, and MRT for free BPA for different organs. The trend for half-life was adult > pediatric PBPK with ontogeny scaling > pediatric PBPK with body-weight scaling. An increased terminal half-life was seen in gonads compared to other organs for both children and adults ($t_{1/2} = 6-9$ h in gonads, $t_{1/2} = 4.5-6$ h for plasma) (Fig. 2 A). The AUC for ontogeny was highest followed by adult and body-weight scaling in all the organs (Fig. 2B): in case of plasma, the AUC for ontogeny based pediatric model was approximately 17% higher, and the classical body-weight-based model was approximately 13% lower than the adult model (Fig. 2B). Bioavailability was also high in children with ontogeny scaling followed by body-weight and adult model (Fig. 2C), but mean residence time was highest in adults compared to children (Fig. 2C). Overall, the pediatric PBPK model based on ontogeny showed the critical differences in pharmacokinetic of short-acting chemical (BPA), which could not have been explained by simply considering the body-weight-based scaling.

3.3. Impact of sex difference on pharmacokinetics characteristic

Further, the ontogeny-based PBPK model for boys and girl (UGT, SULT, and physiology changes) was taken for sex-specific differences in pharmacokinetics for similar exposure. Plasma concentration of free BPA was higher in girls than boys (Supplementary Fig. S11).




Fig. 2. Comparison for different pharmacokinetic parameters in children (based on body-weight and ontogeny scaling) and adults at the dose of 1 $\mu\text{g/kg BW/day}$. 3 A. represents half-life ($t_{1/2}$) for different organs, B. represents area under curve (AUC) for 7 organs (plasma, liver, brain, fat, kidney, skin and gonads), and C. represents bioavailability (F) and mean residence time (MRT) for free BPA in children and adult. Bioavailability (F) is in % and MRT is in hours in Fig. 2C.

Glucuronidation of BPA was slightly higher in boys, however sulfation was comparatively more in girls (Supplementary Fig. S11). Free BPA half-life was almost similar in boys and girls (girls, $t_{1/2} = 4.38$ h vs. boys, $t_{1/2} = 4.63$ h) in plasma and all other organs (Fig. 3). The trend followed by half-life was gonads > fat > kidney > liver > plasma > brain > skin. The mean residence time was slightly higher in boys, but bioavailability and AUC were found to be slightly more in girls (Supplementary Figs. S12 and S13).

3.4. Daily exposure in boys and girls: reverse dosimetry

The ontogeny-based gender-specific PBPK model was used to calculate daily exposure to total BPA for both the boys and girls. Fig. 4 represents the daily dose calculated from the HELIX and the PBAT cohorts for different countries in Europe. Exposure was found to be a little higher in girls than boys for all countries except Greece and Norway. The median daily intake was lowest in France (girls: 0.045, boys: 0.040 $\mu\text{g/kg BW/day}$) and highest in Austria (girls: 0.254, boys: 0.218 $\mu\text{g/kg BW/day}$). Average daily intake estimates for all the cohorts ranged from 0.0138 to 0.294 (P5:P95) with a median of 0.0723 $\mu\text{g/kg BW/day}$. The P95 value was approximately thirteen times lower than the temporary total daily intake (t-TDI) set by EFSA in 2015, which was 4 $\mu\text{g/kg BW/day}$ (EFSA Panel on Food Contact Materials Flavourings and Processing Aids (CEF) 2015). In 2021, EFSA drastically decreased the TDI for BPA to 0.04 ng/kg BW/day recently opened for comments. Now, median value of daily intake is comparatively 1500 times higher than recent TDI established by EFSA (Bisphenol A: EFSA draft opinion proposes lowering the tolerable daily intake 2021). For the model verification we again checked the model with forward dosimetry taking the predicted reconstructed exposure as an input and simulated the total cumulative urine and compared with the cohort data. Correlation was checked by calculating the r^2 , which was found to be 0.95 (Fig. S15, Supp file).

3.5. Sensitivity analysis

Sensitivity analysis has been carried out for all the biochemical parameters used for developing the pediatric PBPK Model for both boys and girls. Table S4 (Supp file) represents the statistics of parameter sensitivity results along with ranking. Ranking for sensitivity varies in boys and girls for many parameters like conjugation, fat and gonads partition coefficient etc. but the mean value for the parameters was quite similar. Fig. 5 represents the mean sensitivity coefficient for boys and

girls respectively. Liver_plasma partition coefficient is highly sensitive for both sex with a sensitivity coefficient of -1 suggesting that 10% increase in the partition may reduce the plasma concentration by 10%. Fat_plasma partition coefficient is more sensitive in girls than boys. The mean sensitivity coefficient of V_{max} is negative, and K_m is positive for both sexes, implicating both the parameters have opposite impact on the output plasma concentration.

4. Discussion

This study was the first to investigate the pharmacokinetic differences between pediatric population considering sex differences with different scaling approaches and adults using the PBPK model for BPA. The development of pediatric PBPK model based on adult model considering 2 approaches: 1) body-weight scaling and 2) ontogeny Scaling. In the present study, data from two different cohort studies was taken and used to estimate daily exposure for boys and girls. Overall, the pediatric PBPK model with ontogeny scaling greatly improves the understanding of toxicokinetic and risk assessment for short-acting molecules like BPA based on sex and age.

The adult model was validated and then extrapolated for the pediatric population following the classical approach of building PBPK. Overall, free BPA metabolizes and eliminates faster in adults as increased levels of BPAS and BPAG were seen. Individual adult PBPK simulations showed not much difference in plasma time profile of free BPA, indicating physiological parameters have little impact on BPA elimination. However, for two subjects, the data did not fit well which could be explained by enzymatic polymorphism leading to high variability of the glucuronidation and sulfation metabolic processes (Skledar et al., 2015; Hanioka et al., 2008). It can be overcome by optimizing the excretion rate for metabolites for those individuals, but this was outside the scope of this paper.

In the pediatric PBPK model, AUC for plasma and all other organs was higher in ontogeny-based scaling representing the underestimation of risk from short-acting chemicals in the case of children with body weight-based scaling. A higher AUC of the free BPA indicates higher systemic exposure to the chemical and comparatively lower metabolism. As mentioned already in the methodology that glucuronidation is lower in children, and sulfation is comparatively higher in children than adults, but the important point is glucuronidation is the major metabolic pathway for BPA (Karrer et al., 2018), which may be responsible for higher AUC in children. Overall, the model indicated that the




Fig. 3. Half-life for boys and girls based on ontogeny scaling PBPK Model at the dose of 1 $\mu\text{g/kg BW/day}$.




Fig. 4. Reconstructed exposure for boys and girls for two different cohorts (HBM4EU and Helix Cohort). Data from Austria is from PBAT cohort from HBM4EU and for rest other countries, it is from Helix cohort. Upper whisker represents P95 and lower whisker represents P5 confidence interval. For Helix cohort, the percentiles are from combining the individual output results. *represents the aggregated data downloaded from HBM4EU website and the distribution represent changes in creatinine levels. X-axis represents country and y-axis represents daily intake levels ($\mu\text{g/kg BW/day}$) for boys and girls.

body-weight scaling approach might lead to an underestimation of the risk.

The estimated half-life was found to be higher in the gonads followed by adipose tissue and other organs, for both pediatric and adult model. This estimate could be correlated with other research finding that gonads are highly sensitive to the BPA exposure, which is known to be an endocrine disruptor (Berger et al., 2018; Braun and Hauser, 2011; Ferguson et al., 2014; Forner-Piquer et al., 2020; Santangeli et al., 2016). This suggests that BPA has potential to affect the developing female and male reproductive systems in children. This finding is in accordance with literature in which Ferguson et al. demonstrated an association between BPA exposure and increased sex hormone-binding globulin (SBHG) levels and reduced free and total testosterone levels in boys (8–14 years) along with several other studies finding links between BPA exposure and varying onset of puberty (Berger et al., 2018; Braun and Hauser, 2011; Ferguson et al., 2014; Forner-Piquer et al., 2020; Santangeli et al., 2016). In all the organs, shorter half-life in children than in adults was in correlation with MRT, emphasizing on the importance of calculating the PK parameters. Nevertheless, bioavailability and AUC

were higher in children with ontogeny scaling than in adults. A higher AUC and bioavailability mean that a greater amount of chemical is present in systemic circulation and hence more risk. The possible reason for increased bioavailability may be due to decreased extent of metabolic capabilities in the liver and intestine in children than adults which was also seen in case of the infant PBPK model for BPA (Karrer et al., 2018). This points towards the question: Should we set a separate TDI for children and adults as children may be more susceptible to risk? However, further research is needed in this area to reach a conclusion.

Sex differences in growing children were represented by PK parameters with half-life ($t_{1/2}$), bioavailability, AUC, MRT and plasma concentration-time curve. The model simulated that bioavailability, AUC and plasma concentration were higher in girls than in boys based on ontogeny-scaling PBPK model. All these parameters are correlated and the reason behind higher value in girls may be the relatively slower conversion of free BPA to BPAG. The elimination half-life was found to be similar in boys and girls in almost all organs. However, further experimental studies are needed to understand in detail the human sex toxicokinetic differences. The model predicted a comparatively higher




Fig. 5. Sensitivity Coefficient for boys and girls. A) Sensitivity plot for boys, B) Sensitivity plot for girls. All the biochemical parameter used for developing PBPK Model has been used for the sensitivity analysis. Analysis has been done with FME Package.

half-life in the gonads than in other organs, which is consistent with literature findings demonstrating BPA exposure associated with variation in puberty onset in girls (Berger et al., 2018), highlighting the advantage of such a model in tissue dosimetry based risk assessment.

Two biomonitoring studies reported higher urine BPA levels in school age children than adults (Abraham and Chakraborty, 2019; Covaci et al., 2015). Study by Covaci et al. has shown 5 out of 6 countries in Europe with having more than 87% population exposed to BPA above the limit of quantification (LOQ) and children with higher exposure than adults (Covaci et al., 2015). Exposure reconstruction through model for these two cohorts showed that exposure is slightly higher in girls than boys for most countries. One possible reason behind this higher exposure can be that girls are getting more exposed than boys through everyday

use products. A recent study by Robles-Aguilera et al., 2021 showed sex-related differences in exposure concluding that a higher risk of dietary exposure in boys was not from BPA but other analogues, but the girls aged under 14 years were at higher risk for increased BPA dietary exposure, especially overweight and obese girls. A critical finding from the pediatric PBPK model is that even with similar exposure in boys and girls, the risk can be slightly higher in girls due to less metabolism in younger girls than boys. The study by Sonnenberg et al. has reported pre-pubertal exposure with phthalates and BPA associated with pubertal timing in girls (mean age: 8.8 years) observed in 472 subjects with no association for boys (Kasper-Sonnenberg et al., 2017). An important outcome of developing ontogeny and sex-specific pediatric models is to better understand the toxicity, metabolic pathway and exposure in

children of growing age.

At last, the reconstructed exposure by the PBPK model for children was compared with the t-TDI established by EFSA in 2015 for BPA (4 µg/kg BW/day) and it was found to be lower than the t-TDI (EFSA Panel on Food Contact Materials Flavourings and Processing Aids (CEF) 2015). It points towards that there may not be any risk or adverse effects involved with the exposure of BPA in human beings. This may not be true. Several epidemiological studies published over the years showed positive associations between BPA exposure and health outcome in children (Healy et al., 2015). But, recent drastic lowering of TDI by EFSA in 2021 (0.04 ng/kg BW/day) which is open for comments, reconstructed exposure by PBPK is approximately 1500 times higher in children pointing towards multiple health concerns ranging from immune effects to neurotoxicity (Bisphenol A: EFSA draft opinion proposes lowering the tolerable daily intake 2021). Further, pediatric PBPK Model can be combined with toxicodynamic to understand the signaling mechanism involving estrogen, aryl hydrocarbon and thyroid hormone receptors (Sharma et al., 2017). This could help in improving the understanding towards adverse effects involved with BPA exposure taking into account toxicokinetic for children.

Interpolation of the pediatric PBPK model from an adult is the most widely used and preferred approach by scientists and regulatory bodies, based on body-weight scaling. In this model, we tried to incorporate ontogeny-specific metabolic parameters to improve the model. This is the first model to consider sex-specific differences in children for BPA. We did not see major differences between genders but, more metabolic and absorption data on sex-specific differences is required to further confirm this. This model also aids in calculating the external exposure which can be from various sources, which is often difficult to calculate during experimental studies. Simulating daily exposure through urine data is another strength of the pediatric PBPK model replacing blood collection for short acting chemicals. PBPK models can predict the aggregate exposure which may be from multiple sources. However, the limitation of these models is the inability to predict route of exposure (oral, dermal, inhalational or sublingual exposure). Further refinement of the model by including separate exposure sources and dermal compartments can help overcome some of the limitations. Overall, the pediatric PBPK model provides a framework to integrate the sex differences and consider pediatric population for the risk assessment in case of such short-acting chemicals. Also, toxicologists should take into account sex-related differences while developing research protocols for environmental chemicals since boys and girls may have quite different effects on the health even with the same exposure.

5. Conclusion

In summary, the BPA pediatric PBPK model was developed for dosimetry assessment and predicting the tissue concentration-time profile in plasma and other organs. At the same exposure, the ontogeny-based pediatric PBPK model showed children might be at higher risk than adults (higher AUC and higher bioavailability in all the organs), which could not be explained by classical body-weight scaling. AUC and bioavailability were slightly higher in girls than boys suggesting free BPA is more present in girls for the same exposure. Further, the reconstructed daily exposure was found to be higher than the TDI recently established by EFSA and reconstructed daily exposure from a large cohort was found to be slightly higher in girls than boys. Overall, this study presents an example of how ontogeny based pediatric PBPK model can show a larger difference in kinetics than traditional approach which further warrants using this kind of model would improve the risk assessment. Also, it is one step forward towards incorporating sex-specific differences in the PBPK model improving the translation of risk.

Credit roles

Deepika Deepika: Conceptualization, Methodology, Formal analysis

and simulation, and writing; Raju Prasad Sharma: Supervision, Conceptualization, Methodology, writing, review and revision; Marta Schuhmacher: Supervision, Review and revision; Amrit Kaur Sakhi: Provided Cohort Data and reviewing; Cathrine Thomsen: Provided Cohort Data and reviewing; Leda Chatzi: Provided Cohort Data and reviewing; Marina Vafeiadi: Provided Cohort Data and reviewing; Remy Slama: Provided Cohort Data and reviewing; Joane Quentin: Provided Cohort Data and reviewing; Sandra Andrusaitytė: : Provided Cohort Data and reviewing; Regina Grazuleviciene: Provided Cohort Data and reviewing; Tiffany C Yang: Provided Cohort Data and reviewing; John Wright: Provided Cohort Data and reviewing; Dagmar Waiblinger: Provided Cohort Data and reviewing; Martine Vrijheid: Provided Cohort Data and reviewing; Jose Urquiza: Provided Cohort Data and reviewing; Maribel Casas: Provided Cohort Data, Writing, Review and revision; José L. Domingo: Supervision, Review and revision; Vikas Kumar: Supervision, Conceptualization, Methodology, Formal analysis and simulation, writing, review and revision.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgment

This study was financially supported by Marie Skłodowska-Curie “Neurosome Project” under the grant agreement No. 766251, the European Community funded H2020 HBM4EU project under Grant Agreements no. 733032, and the Instituto de Salud Carlos III (PI17/01194), and the European Community’s Seventh Framework Programme (FP7/2007–2013) under grant agreement no 308333 (HELIX project). Maribel Casas holds a Miguel Servet fellowship (CP16/00128) funded by Instituto de Salud Carlos III and co-funded by European Social Fund “Investing in your future”. We acknowledge support from the Spanish Ministry of Science and Innovation through the “Centro de Excelencia Severo Ochoa 2019–2023” Program (2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program. This publication reflects only the authors’ views. The Community and other funding organizations are not liable for any use made of the information contained therein. Born in Bradford is only possible because of the enthusiasm and commitment of the children and parents in Born in Bradford. We are grateful to all participants, health professionals and researchers who have made Born in Bradford happen. BiB receives core infrastructure funding from the Wellcome Trust (WT101597MA) and a joint grant from the UK Medical Research Council (MRC) and Economic and Social Science Research Council (ESRC) (MR/N024397/1). This study has received support from European Research Council under the European Union’s Seventh Framework Programme (FP7/2007–2013)/ERC grant agreement no 669545 and National Institute for Health Research Applied Research Collaboration Yorkshire and Humber (NIHR200166). The Norwegian Mother, Father and Child Cohort Study is supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and Research. We are grateful to all the participating families in Norway who take part in this on-going cohort study. The views expressed are those of the author(s), and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. Jose Urquiza is supported by Spanish regional program PERIS (Ref.: SLT017/20/000119), granted by Departament de Salut de la Generalitat de Catalunya.

- Sharma, R.P., Schuhmacher, M., Kumar, V., 2018a. Development of a human physiologically based pharmacokinetic (PBPK) model for phthalate (DEHP) and its metabolites: a bottom up modeling approach. *Toxicol. Lett.* 296, 152–162. <https://doi.org/10.1016/j.toxlet.2018.06.1217>.
- Sharma, R.P., Schuhmacher, M., Kumar, V., 2018b. The development of a pregnancy PBPK Model for Bisphenol A and its evaluation with the available biomonitoring data. *Sci. Total Environ.* 624, 55–68. <https://doi.org/10.1016/j.scitotenv.2017.12.023>.
- Shelby, M.D., 2008. NTP-CERHR Monograph on the Potential Human Reproductive and Developmental Effects of Bisphenol A. NTP CERHR MON v, pp. 1–64 passim vols. ii–ix.
- Skledar, D.G., Troberg, J., Lavdas, J., Mašič, L.P., Finel, M., 2015. Differences in the glucuronidation of bisphenols F and S between two homologous human UGT enzymes, 1A9 and 1A10. *Xenobiotica* 45, 511–519. <https://doi.org/10.3109/00498254.2014.999140>.
- Soetaert, K., Petzoldt, T., 2010. Inverse modelling, sensitivity and Monte Carlo analysis in R using package FME. *J. Stat. Software* 33, 1–28. <https://doi.org/10.18637/jss.v033.i03>.
- Stahlhut, R.W., Welshons, W.V., Swan, S.H., 2009. Bisphenol A data in nhanes suggest longer than expected half-life, substantial nonfood exposure, or both. *Environ. Health Perspect.* 117, 784–789. <https://doi.org/10.1289/ehp.0800376>.
- Takeuchi, T., Tsutsumi, O., 2002. Serum bisphenol a concentrations showed gender differences, possibly linked to androgen levels. *Biochem. Biophys. Res. Commun.* 291, 76–78. <https://doi.org/10.1006/bbrc.2002.6407>.
- Tamayo-Uria, I., Maitre, L., Thomsen, C., Nieuwenhuijsen, M.J., Chatzi, L., Siroux, V., et al., 2019. The early-life exposome: description and patterns in six European countries. *Environ. Int.* 123, 189–200. <https://doi.org/10.1016/j.envint.2018.11.067>.
- Teeguarden, J.G., Waechter, J.M., Clewell, H.J., Covington, T.R., Barton, H.A., 2005. Evaluation of oral and intravenous route pharmacokinetics, plasma protein binding, and uterine tissue dose metrics of bisphenol A: a physiologically based pharmacokinetic approach. *Toxicol. Sci.* 85, 823–838. <https://doi.org/10.1093/toxsci/kfi135>.
- Thayer, K.A., Doerge, D.R., Hunt, D., Schurman, S.H., Twaddle, N.C., Churchwell, M.I., et al., 2015. Pharmacokinetics of bisphenol A in humans following a single oral administration. *Environ. Int.* 83, 107–115. <https://doi.org/10.1016/j.envint.2015.06.008>.
- Tsukioka, T., Terasawa, J., Sato, S., Hatayama, Y., Makino, T., Nakazawa, H., 2004. Development of analytical method for determining trace amounts of BPA in urine samples and estimation of exposure to BPA. *J. Environ. Chem.* 14, 57–63. <https://doi.org/10.5985/jec.14.57>.
- Völkel, W.V., Bittner, N., Dekant, W., 2005. Quantitation of bisphenol A and bisphenol A glucuronide in biological samples by high performance liquid chromatography-tandem mass spectrometry. *Drug Metab. Dispos.* 33, 1748–1757. <https://doi.org/10.1124/dmd.105.005454>.
- Vrijheid, M., 2013. The human early life exposome (HELIX): project rationale and design. In: ISEE Conf Abstr 2013:5849. <https://doi.org/10.1289/isee.2013.s-2-32-05>.
- Wagner, C., Zhao, P., Pan, Y., Hsu, V., Grillo, J., Huang, S.M., et al., 2015. Application of physiologically based pharmacokinetic (PBPK) modeling to support dose selection: report of an FDA public workshop on PBPK. *CPT Pharmacometrics Syst. Pharmacol.* 4, 226–230. <https://doi.org/10.1002/psp4.33>.