

A super cool scientific Title ...

Gautier VAILLANT

August 20, 2015

Acknowledgments

I would first like to gratefully thank Helmut Grubmüller and Bert de Groot for welcoming me in their lab. I also thank Carsten Kutzner who supervised my Internship. I also would like to Thank Bartosz Kohnke and Thomas Ullmann for providing me advice on my project.

Finally I would also like to thank Ivo Kabadschow and Andreas Beckmann who welcomed me in the Jülich Forschungszentrum to teach me the details of the FMM method.

Abstract

Simulating systems with long-range pairwise interactions is a frequently encountered challenge in scientific research. It plays an important role in Astrophysics to know the dynamics of galaxies, in plasma physics or in our case in biophysics. A direct evaluation of these pairwise interactions leads to $\mathcal{O}(N^2)$, which obviously scaled badly with a system of size N . Some other techniques, such as the PME (Particle Mesh Ewald) and the FMM (Fast Multipole Method) obtain a complexity of $\mathcal{O}(N \log(N))$ and $\mathcal{O}(N)$, respectively these techniques also allow a greater scalability of the system for parallel computations.

The aim of this internship is to study how the FMM compares to the PME, ie. which sets of parameters gives the same accuracy for both methods. We developed some tools to compare the data from different softwares.

La simulation de larges systemes de particules en interaction est tres importante pour le calcul scientifique. Elles jouent un role important en Astrophysique pour connaitre la dynamique des galaxies, en physique des plasmas ainsi que, dans notre cas, en Biophysique pour l'etude de machines moleculaires complexes.

Contents

1	Presentation of the Lab	6
2	Context of the Internship	8
3	Methods for computing electrostatic forces	11
3.1	Direct summation method	11
3.1.1	"Naive" $\mathcal{O}(N^2)$ Method	11
3.1.2	Possible improvements	13
3.2	Fourier Transform-Based methods	14
3.2.1	Ewald Summation	14
3.2.2	PME	16
3.3	Tree-based methods	18
3.3.1	Mathematical preliminaries	18
3.3.2	Workflow of the algorithm	20
	Splitting the Space	20
	Defining a Separation Criterion	21
	PASS 1 : Computing the Multipole moments	22
	PASS 2 : Transforming distant expansions into multi- poles	23
	PASS 3 : Shift Taylor-Like expansions down the tree .	24
	PASS 4 : Computing forces and Energies	24
4	Comparing FMM and PME accuracy	26
4.1	Presentation of GROMACS	26
4.1.1	Structure of a File	26
4.1.2	PME parameters	27
4.1.3	Command to launch a GROMACS simulation	27
4.2	Presentation of fmsolvr	28
4.2.1	FMM Parameters	28
4.3	Making GROMACS and fmsolvr comparable	29
4.3.1	File manipulation	29

Contents

	Making GROMACS only compute electrostatics forces	30
	Assure compatibility between *.qxyz and *.pdb files . .	30
	Output positions and forces to a Text file for GROMACS	30
4.4	First Comparisons and Dipole Correction	31
4.4.1	Silicamelt Simulation	31
4.4.2	Dipole correction	33
	Study of a two-particle system	33
	Back to the silicamelt simulation with dipole correction	37
4.5	Comparing with an analytical solution : the NaCl system . . .	38
4.5.1	Generating the system	38
4.5.2	Some explanations on the Owl cluster	39
4.5.3	Error plots	40
Appendices		45
A Glossary and Abbreviations		46
B Structure of Gromacs files		47
B.1	pdb file	47
B.2	top file	47
B.3	mdp file (longe-range electrostatics part)	49

Chapter 1

Presentation of the Lab

Max Planck Institute for Biophysical Chemistry

The Max Planck Institute for Biophysical Chemistry is part of the Max-Planck Institutes, which are an ensemble of research center throughout Germany (As the CNRS in France could be).

The Max-Planck Institute for Biophysical Chemistry, located in Göttingen, is one of the biggest research centers in the Biophysics field in Europe. Its research fields includes Molecular Dynamics Simulation, but also NMR or Biochemistry.

The departments are : (from <http://www.mpibpc.mpg.de/groups>)

- Patrick Cramer - Molecular Biology
- Gregor Eichele – Genes and Behavior
- Dirk Görlich – Cellular Logistics
- Christian Griesinger – NMR-based Structural Biology
- **Helmut Grubmüller – Theoretical and Computational Biophysics**
- Stefan W. Hell – NanoBiophotonics
- Herbert Jäckle – Molecular Developmental Biology
- Reinhard Jahn – Neurobiology
- Reinhard Lührmann – Cellular Biochemistry

- Marina V. Rodnina – Physical Biochemistry
- Melina Schuh – Meiosis
- Alec M. Wodtke – Dynamics at Surfaces

Department of Theoretical and Computational Biophysics

This department is led by both Helmut Grubmüller. It aims at understanding of the physics and function of proteins, protein complexes, and other biomolecular structures at the atomic level. For this purpose, dynamic complex atomistic computer simulations are carried out.

One area of research is on studying systems (Proteins, membranes) and molecular machines in order to better understand their dynamics (for example the molecular mechanisms of the ribosome), and another area is focusing on development. Concerning the part of the department dedicated to methods and theory, new statistical mechanics concepts, quantum hybrid methods, and efficient parallel simulation algorithms and codes are the methodological focus of the department.

Both lines of questions phrased above are closely interlinked. Progress in the understanding of the physics of proteins, on the one hand, enables improved and more realistic simulation techniques, which allow to study a growing number of biochemical processes in great detail. Through analysis of well-understood mechanisms, on the other hand, one can learn to separate relevant aspects in protein dynamics from irrelevant ones — which is prerequisite for the construction of effective protein models. In short, we find a close interplay between method development, algorithmic progress, and application.

Chapter 2

Context of the Internship

The context of this internship is driven by the "SPPEXA (Software for exascale computing) / GromEx" project funded by the DFG (Deutsche Forschungsgemeinschaft). The Idea of this project is to create a flexible and fast solver for computing forces and potentials, which is a preliminary for molecular simulations.

A poster¹ of the project can be found below :

¹from <http://www.mpibpc.mpg.de/grubmueller/sppexa>

German Priority Programme 1648 “Software for Exascale Computing”

SPPEXA – Findings & Goals

- Massive parallelism (on- and cross-chip) requires fundamentally new concepts
- Not “racks without brains”, but software is the key to this paradigm shift
- Fundamental research (→ DFG), in contrast to other (more application-oriented) initiatives (→ German Federal Ministry of E & R)
- Establish collaborative, interdisciplinary co-design of HPC applications and HPC methods
- Focus on six research directions:
 - Computational algorithms
 - Application software
 - Programming
 - System software
 - Data management and exploration
 - Software tools

SPPEXA – Implementation

- Two three-year funding phases
- Overall budget of 3,7 M € per year
- Funded via DFG’s strategy fund
- Interdisciplinary consortia of 3–5 groups
- Consortia address at least two of SPPEXA’s six research directions
- Two-stage application process with (1) sketches and (2) full proposals
- Global strategic coordination, following the established procedures of Collaborative Research Centres (SFB)
- Close collaboration with respective international programmes intended

SPPEXA – Chronology

- 2006: discussion in the German Research Foundation (DFG) on the necessity of a funding initiative for HPC software
- 2010: initiative out of German HPC community, referring to increasing activities on HPC software elsewhere (USA: NSF, DOE; Japan; China; G8)
- 2010: discussion with DFG’s Executive Committee, suggestion of a flexible, strategically initiated SPP
- 2011: submission of the proposal, international reviewing, and formal acceptance
- 2012: Review of project sketches and full proposals

SPPEXA – Current Status

- 68 sketches handed in, overall volume of 19 M € per year applied for
- 80 different universities, institutes and companies represented by 240 national and 15 international PIs
- 24 sketches invited for full proposals
- 13 full proposals accepted for funding
- Launch of programme and projects in January 2013

www.sppexa.de




Figure 2.1: Poster for the SPEXXA project

The current method for computing electrostatic forces is called the PME (Particle Mesh Ewald). One of its problems is a communication bottleneck that prohibits efficient parallelization across many (> 10000) CPU cores. The idea would be to replace this method with a new method called the Fast-Multipole Method which is based on a tree Structure and may allow an greater parallelization of the system as wanted.

So the Idea of the Internship is first to know how to tune those systems to know how they relate to each other. Both PME and FMM make approximations. For a fair comparison we want to compare the methods at equal

accuracy of the forces (and energies) they compute.

Chapter 3

Methods for computing electrostatic forces

The electrostatic potential of a group of N of charge q_i at position r_i is defined as :

$$V_C = \frac{1}{4\pi\epsilon_0} \sum_{i=1}^N \sum_{j>i} \frac{q_i q_j}{r_{ij}} \quad (3.1)$$

Computing this potential is very important in different areas of physics, such as plasma physics, or in our case molecular dynamics where the forces between charged particles need to be computed in an efficient and scalable way, the force being computed using

$$\vec{F}_i = -\vec{\nabla}_i V(\vec{r}) \quad (3.2)$$

3.1 Direct summation method

In this section, we will explain the most basic method to compute pairwise interactions and explain why the method leads to long computation times and sometimes .

3.1.1 "Naive" $\mathcal{O}(N^2)$ Method

The sum of all electrostatic forces F_i exerted on one particle can be written the following way:

3.1. Direct summation method

$$F_i = \sum_{i=1}^N \sum_{j>i} \frac{q_i q_j}{4\pi\epsilon_0 r_{ij}^2} \quad (3.3)$$

where q is the charge of one particle, and R_{ij} is the distance between particle i and particle j .

In the thesis we will simplify the equation (3.3) by just writing:

$$\vec{F}_i = f \cdot \sum_{i=1}^N \sum_{j>i} \frac{q_i q_j}{r_{ij}^2} \quad (3.4)$$

where $f = \frac{1}{4\pi\epsilon_0}$ is called the **electric conversion factor** and is equal to 138.935485(9) kJ.mol⁻¹.nm.e⁻²

The first, naive way to compute electrostatic forces is just to follow equation (3.4):

So if we consider a set of N charged particles, $N - 1$ interactions are needed to compute the force acting on one specific particle. So in order to know the forces of the set of particles, $N \cdot (N - 1)$ operations are needed, hence an algorithmic complexity of $\mathcal{O}(N^2)$. Indeed, in order to have to have a physically accurate simulation, it is need to surround the molecule we want to study into a solvent such as water : If the layer of water is finite, it can lead to some unexpected effects as shown [REF...]

This gives the following algorithm:

```

input : A set of  $N$  charged particles
output: A list of the forces for each particle

For each particle  $i$ ;
for  $i \leftarrow 1$  to  $N - 1$  do
    | add interaction between particle  $i$  and particle  $j$  ;
    for  $j \leftarrow i + 1$  to  $N$  do
        | force [ $i$ ]  $\leftarrow$  force [ $i$ ] + computeForce( $i, j$ ) ;
    end
end

```

Algorithm 1: Naive method

The complexity of such a computation limits its use to rather small systems and is not really usable for bigger systems such as proteins or astrophysical systems. Moreover it is also important to say that if we want to add periodic boundary conditions the $\mathcal{O}(N^2)$ method is not sufficient as the system is infinite.

3.1. Direct summation method

3.1.2 Possible improvements

A possible method to overcome this limitation for periodic systems is to limit the interaction to a certain radius : if the distance between two particles is greater than R_0 , then the force is set to 0.

So we have the following system :

$$\vec{F}_{A \rightarrow B} = \begin{cases} \frac{q_A q_B \hat{r}_{AB}}{|\vec{R}_{AB}|^2} & \text{if } R_{AB} < R_0 \\ 0 & \text{otherwise} \end{cases} \quad (3.5)$$

This technique is for example used for Lennard-Jones potentials ($V_{LJ} = 4\epsilon[(\frac{\sigma}{r})^{12} - (\frac{\sigma}{r})^6]$), where the intensity of the force is quickly decreasing. It allows to limit the number of interactions to only the close neighbors.

However, one of the problems of this cutoff technique, especially for long-range interactions such as coulombic interactions using a cut-off can lead to artifacts resulting from the sudden drop of the force to 0 at the cutoff. It was shown that this can lead to unphysical assemblies of particles at the cutoff distance as shown for instance in figure (3.1) .




Figure 3.1: Radial distribution function (RDF) $g(r)$ between the two central atoms in the head-group of lipids: Cutoff distances are indicated by arrows. from [ADD REFERENCE]

As we can see in figure 3.1, the radial distribution of the distance be-

3.2. Fourier Transform-Based methods

tween two atom shows a peak, corresponding to the cutoff of the system. This shows that by using a cutoff technique we might see some artifacts.

So we can see at least three reasons not to use the direct summation method : The first and main reason is that the method doesn't work with periodic boundary conditions. Then, is cutoff methods are used to solve the problem, artifacts might arise at cutoff distances. Finally the method is computationally inefficient.

3.2 Fourier Transform-Based methods

To compute the potentials and the forces of particles, Fourier-transform based techniques have been developed in order to overcome the problems stated above .

3.2.1 Ewald Summation

This subset of techniques comes from a theoretical physics technique called the Ewald summation.

in periodic boundary conditions, the Coulomb potential V_C is:

$$V = \sum_{n_x, n_y, n_z} \sum_i^N \sum_{j>i} \frac{q_i q_j}{r_{ij}} \quad (3.6)$$

where n_x, n_y, n_z are the box index vector.

The equation (3.6) is conditionally convergent and slow to converge. One technique discovered by Ewald is to split the potential in two absolutely convergent terms and one constant term:

$\frac{1}{r}$ can be written in the following way :

$$\frac{1}{r} = \frac{f(x)}{r} + \frac{1 - f(x)}{r} \quad (3.7)$$

The idea is to choose f , so that $\frac{f(x)}{r}$ is quickly decaying and can be computed in real space with a cutoff at high accuracy. At the same time, we want $\frac{1-f(x)}{r}$ is as smooth as possible and therefore be represented in reciprocal space with just a few \vec{k} vectors This gives a quick and accurate computation of the real and the reciprocal space where $V = V_{\text{direct}} + V_{\text{reciprocal}}$

3.2. Fourier Transform-Based methods

A good choice is often $f(x) = \text{erfc}(\alpha x)$, the complementary error function, where α is called the splitting parameter, and $\text{erfc}(x) = \frac{2}{\sqrt{\pi}} \int_x^{+\infty} e^{-t^2} dt$

so we obtain the following potentials :

For the direct space :

$$V_{\text{direct}} = \frac{1}{2} \sum_{\vec{n}} \sum_{i,j}^N q_i q_j \frac{\text{erfc}(\alpha r)}{r} \quad (3.8)$$

and for the reciprocal space :

$$V_{\text{reciprocal}} = \frac{1}{2\pi L} \sum_{i,j}^N q_i q_j \sum_{\vec{m}} \frac{\exp(-(\pi \vec{m}/\alpha)^2 + 2\pi i \vec{m} \cdot (r_i - r_j))}{\vec{m}^2} \quad (3.9)$$

where \vec{m} are the vectors in the reciprocal space, and L is the volume of the cell.

$$\begin{cases} \vec{m}_x = 2\pi \frac{\vec{n}_y \times \vec{n}_z}{\vec{n}_x \cdot (\vec{n}_y \times \vec{n}_z)} + n \cdot L_x \\ \vec{m}_y = 2\pi \frac{\vec{n}_z \times \vec{n}_x}{\vec{n}_y \cdot (\vec{n}_z \times \vec{n}_x)} + n \cdot L_y \\ \vec{m}_z = 2\pi \frac{\vec{n}_x \times \vec{n}_y}{\vec{n}_z \cdot (\vec{n}_x \times \vec{n}_y)} + n \cdot L_z \end{cases}, n \in \mathbb{Z}$$

Once the electrostatic potentials are obtained, it is possible to differentiate the potentials with respect to the position $p = (x, y, z)$ to obtain the force on each particle.

$$\vec{F}_i^{\text{direct}} = \vec{\nabla}_i V_{\text{direct}} \quad (3.10)$$

so we have:

$$\vec{F}_p^{\text{direct}} = q_i \sum_{i=1, i \neq j}^N \sum_{\vec{n}} q_j \frac{(r_{ij, \vec{n}})_p}{r_{ij, \vec{n}}^3} \{ \text{erfc}(\alpha(r_{ij, \vec{n}})) + \frac{2\alpha}{\sqrt{\pi}} r_{ij, \vec{n}} \exp(-(\alpha r_{ij, \vec{n}})^2) \} \quad (3.11)$$

$$\vec{F}_p^{\text{reciprocal}} = \frac{2q_i}{L} \sum_{i=1, i \neq j}^N \sum_{\vec{m} \neq 0} \frac{\vec{m}_p^*}{\vec{m}^{*2}} \exp(-(\frac{\pi \vec{m}}{\alpha L})^2) \sin \frac{2\pi}{L} \vec{m} \cdot r_{ij} \quad (3.12)$$

3.2. Fourier Transform-Based methods

Equations 3.11 and 3.12 can then be used to compute the force on each particle by adding the direct space and the reciprocal space contribution.

3.2.2 PME

One way to improve this method is to compute the Fourier sum using the FFT (Fast Fourier Transform). This allows the algorithm to get a complexity of $\mathcal{O}(N \log N)$. There is different methods that are based on the Ewald summation, namely the **P3M** (Particle-Particle Particle-Mesh Method) or the **FFP** (Fast Fourier Poisson method).

In our case, we will focus on the **PME** as it is the method currently used in GROMACS.

We remind that the potential for the reciprocal space $V_{\text{reciprocal}}$ is written in equation (3.9):

$$V_{\text{reciprocal}} = \frac{1}{2\pi V} \sum_{i,j}^N q_i q_j \sum_{\mathbf{n}^*} \frac{\exp(-(\pi \vec{m}/\alpha)^2 + 2\pi i \vec{m} \cdot (r_i - r_j))}{m^2}$$

This equation can be rewritten as :

$$V_{\text{reciprocal}} = \frac{1}{2\pi V} \sum_{\mathbf{n}^* \neq 0}^N \frac{\exp(-(\pi \vec{m}/\alpha)^2)}{\vec{m}^2} S(-\vec{m}) S(\vec{m})$$

where $S(\mathbf{n}^*)$ is defined as the Structure factor:

$$S(\mathbf{n}^*) = \sum_{k=1}^N q_k \exp(2\pi i \vec{m} \cdot \mathbf{r}) \quad (3.13)$$

The idea of the PME is to approximate the structure factor $S(\vec{m})$ by the 3D Fourier Transform of the charge matrix, which is obtained by interpolating the charges to a discrete grid of size $p_x \times p_y \times p_z$. Let also define $\mathcal{F}(Q)$ the 3D FFT of Q . The charges q_i are mapped to the grid using interpolation. Originally, Lagrange interpolation was used, but now a b-spline interpolation is used. The order of interpolation is called the *PME Order*

then the Structure factor can be approximated as its FFT :

$$S(\vec{m}) \approx \tilde{S}(\vec{m}) = \mathcal{F}(Q)(\vec{m}) \quad (3.14)$$

3.2. Fourier Transform-Based methods

so the reciprocal energy can also be approximated by:

$$V_{\text{reciprocal}} \approx \tilde{V}_{\text{reciprocal}} = \frac{1}{2\pi V} \sum_{\mathbf{n}^* \neq 0}^N \frac{\exp(-(\pi \vec{m} / \alpha)^2)}{m^2} \mathcal{F}(Q)(\mathbf{m}) \mathcal{F}(Q)(-\mathbf{m}) \quad (3.15)$$

It can be shown that the complexity of such a system is $\mathcal{O}(N \log(N))$ which is a much bigger improvement compared to the $\mathcal{O}(N^2)$ complexity of the direct algorithm. Then it also allows to handle infinite systems with periodic boundary conditions.

However, the algorithm doesn't scale well for large scale parallelism (above 10000 CPU cores)[REFERENCE ?]. The bottleneck is located in the computation of the FFT, where an all-to-all communication is required, hence the need for a more scalable algorithm for the computations of electrostatic forces.




Figure 3.2: Parallel efficiency of three "real-world" simulations using Intel's MPI. Horizontal grey line indicates perfect scaling.

3.3. Tree-based methods

3.3 Tree-based methods

Now we will look at an algorithm that scales linearly with the number of charges N and which does not have inherent parallel communication bottleneck as [ref]. Furthermore, it can be applied to systems with periodic as well to systems with open boundaries. Since it approximates the actual charge distribution by a set of multipoles, it is called the "Fast Multipole method" (FMM)

3.3.1 Mathematical preliminaries

Let's move back to the potential created by a charged particle as in (??),

$$V = \frac{1}{d}$$

Let two particles $A(a, \alpha, \beta)$ and $R(r, \theta, \phi)$, separated by a distance d : The distance between the particles is $d = |a - r|$. We would like to achieve is to "factorize the addition", having a product of one function depending only on a and one depending only on r .




Figure 3.3: Two particles $A(a, \alpha, \beta)$ and $R(r, \theta, \phi)$, separated by a distance d

The inverse distance can be written as :

$$\frac{1}{d} = \frac{1}{|a - r|} = \frac{1}{\sqrt{a^2 + r^2 - 2ar \cos(\gamma)}}$$

This distance can then be written as a series of Legendre polynomials $P_l(u)$ of degree l if $r \gg a$:

$$\frac{1}{d} = \frac{1}{\sqrt{a^2 + r^2 - 2ar \cos(\gamma)}} = \sum_{l=0}^{+\infty} P_l(u) \mu^l \quad (3.16)$$

where $\mu = \frac{a}{r}$ and $u = \cos \gamma$.

3.3. Tree-based methods

with some manipulation as explained [ADD REFERENCE], we obtain :

$$\frac{1}{|r-a|} = \sum_{l=0}^{+\infty} \sum_{m=-l}^{+l} \frac{(l-m)!}{(l+m)!} \frac{a^l}{r^{l+1}} P_{lm}(\cos(\alpha)) P_{lm}(\cos(\theta)) e^{-im(\beta-\phi)} \quad (3.17)$$

in order to approximate the scheme, we can truncate the infinite series to a certain order p , we call this order the **multipole order**

$$\frac{1}{|r-a|} \simeq \sum_{l=0}^{\overset{p}{\color{red}}} \sum_{m=-l}^{+l} \frac{(l-m)!}{(l+m)!} \frac{a^l}{r^{l+1}} P_{lm}(\cos(\alpha)) P_{lm}(\cos(\theta)) e^{-im(\beta-\phi)} \quad (3.18)$$

We can now rewrite the summation the following way :

$$\frac{1}{|r-a|} \simeq \sum_{l=0}^p \sum_{m=-l}^{+l} \underbrace{\frac{a^l}{(l+m)!} P_{lm}(\cos(\alpha)) e^{-im\beta}}_{O_{lm}(\mathbf{a})} \underbrace{\frac{(l-m)!}{r^{l+1}} P_{lm}(\cos(\theta)) e^{+im\phi}}_{M_{lm}(\mathbf{r})} \quad (3.19)$$

hence,

$$\frac{1}{|r-a|} \simeq \sum_{l=0}^p \sum_{m=-l}^{+l} O_{lm}(\mathbf{a}) M_{lm}(\mathbf{r}) \quad (3.20)$$

Thus it is possible to "factorize" the inverse of the the distance $\frac{1}{d}$ so we can obtain two independent terms $O_{lm}(\mathbf{a})$ and $M_{lm}(\mathbf{r})$ for two particles

We can then define the multipole moment $\omega_{lm}(q, \mathbf{a}) = q O_{lm}(\mathbf{a})$ and the Taylor-like moment $\mu_{lm}(q, \mathbf{r}) = q M_{lm}(\mathbf{r})$

One obtains the following "bipolar expansion" for two particles associated with two different origins :

$$\frac{1}{|\mathbf{a}_1 - \mathbf{a}_2 + \mathbf{R}|} = \sum_{l=0}^{+\infty} \sum_{j=0}^{+\infty} \sum_{m=-l}^{+l} \sum_{k=-j}^{+j} (-1)^j \cdot O_{lm}(\mathbf{a}_1) \cdot M_{l+j, m+k}(\mathbf{R}) \cdot O_{jk}(\mathbf{a}_2) \quad (3.21)$$

3.3. Tree-based methods




Figure 3.4: Scheme of two particles expanded according to two different origins [from ADD REFERENCE]

3.3.2 Workflow of the algorithm

Once some mathematical preliminaries are set, it is possible to explain the workflow of the algorithm.

Splitting the Space

One of the ideas of the FMM is to approximate far-away charges by multipoles expansions, whereas interactions between near charges are calculated directly without any approximation. So that's why we need to split the simulation box in sub-boxes, so that near field and far field contributions can be computed everywhere.

So first step of the scheme is to split the space in order to generate different groups. The idea is to recursively split the space in eight octants, created the so-called structure of an *octree*. So the number of boxes is 8^{D-1} , where D is the depth of the oct-tree.

3.3. Tree-based methods




Figure 3.5: Example of a division of the space in boxes. The example is in 2D so the structure of the space is called a *quadtree*.

The number of subdivisions is called the *depth* D . For example, in figure 3.5, we can observe from left to right the depths 0, 1, 2 and 3.

Then each particle is assigned to the box of lowest depth it is found in.

Defining a Separation Criterion

For the further computations it is needed to define a separation criterion ws : It is the ws^{th} next neighbors of a given depth. Two boxes A and B are called *next neighbors* if they are at the same level and box B is enclosed by a box of size $(2ws + 1)^3$ around the center of A




Figure 3.6: Example of the influence of the separation criterion ws for a given box A, from left to right, $ws = 1, 2, 3$.

This separation criterion divides the space into two parts. If two particles are close so they are in the same grey space as defined figure (3.6), then, their interaction is computed directly, otherwise their interaction is approximated via multipoles.

It is most of the time not possible to set the separation criterion ws to 0, as the multipole expansion might overlap. Using a bigger ws means that the order of the multipole expansions need to be less important for the same

3.3. Tree-based methods

precision, however this will increase the number of direct $\mathcal{O}(N^2)$ operation, which are more computationally expensive.

The algorithm by itself will be done in separate phases we call "passes" : there will be 4 different passes:

PASS 1 : Computing the Multipole moments

PASS 2 : Transforming distant expansions into multipoles

PASS 3 : Shift Taylor-Like expansions down the tree

PASS 4 : Computing forces and Energies

PASS 1 : Computing the Multipole moments

Once every particle is assigned to its box at the lowest level, we will compute the multipole moment ω_{lm}^j of each box.

$$\omega_{lm}^j(q, \mathbf{a}) = q^j a_j^l \tilde{P}_{lm}(\cos(\alpha_j)) e^{-im\beta_j} \quad (3.22)$$

To construct the multipoles at the higher levels from the existing multipoles at the lowest level, each multipole order is moved to the Center of the parent box using the so-called *M2M* (Multipole to Multipole) operator:

$$\omega(\mathbf{a} + \mathbf{b}) = \sum_{j=0}^k \sum_{k=-j}^j \omega_{jk}(\mathbf{a}) O_{l-j, m-k}(\mathbf{b}) \quad (3.23)$$

$O_{l-j, m-k}(\mathbf{b})$ is called the *M2M* operator.




Figure 3.7: Example in 1-D showing the shifting of the multipole moments using the M2M operator.

3.3. Tree-based methods

PASS 2 : Transforming distant expansions into multipoles

PASS2 transforms distant expansions (beyond the separation criterion) ω_{lm} into a Taylor-like moment μ_{lm} . This transformation is done over at most 189 boxes. This transformation is applied for each depth of the tree.






Figure 3.8: Example in 1-D showing the transformation of distant expansions into multipoles

In order to compute this transformation, two different rules have to be applied:

1. Take the non-separated parent boxes, as shown sub-figure (a) below.
2. Then the children of the parent box are selected, taking into account the separation criterion for the child boxes.



(a) First, the parents of the box are chosen



(b) Then, then children of the parents are chosen. Then the separation criterion can apply

Figure 3.9: Iteration Rules for the transformation of the multipole expansions

3.3. Tree-based methods

Mathematically, the transformation from multipole expansions to multipole moments is given using the M2L (Multipole to Local) Operator

$$\mu_{lm}(\mathbf{b} - \mathbf{a}) = \sum_{j=0}^{+\infty} \sum_{k=-j}^j M_{j+l,k+m}(\mathbf{b}) \cdot \omega_{jk}(\mathbf{a}) \quad (3.24)$$

where $M_{j+l,k+m}(\mathbf{b})$ is the M2L operator : It allows to exchange, as figure (??) shows, information between boxes of the same level.

PASS 3 : Shift Taylor-Like expansions down the tree

Once the local μ_{lm} expansions are computed, the expansion are shifted to the lowest level of the tree using the L2L (Local to Local) operator.




Figure 3.10: Example in 1-D showing the shifting of the local moments using the L2L operator.

This shift operation is done by the following formula :

$$\mu_{lm}(\mathbf{r} - \mathbf{b}) = \sum_{j=l}^p \sum_{k=-j}^j O_{j-l,k-m}(\mathbf{b}) \cdot \mu_{jk}(\mathbf{a}) \quad (3.25)$$

where $O_{j-l,k-m}(\mathbf{b})$ is the L2L operator : It allows to exchange, as figure (??) shows, information between levels downwards.

PASS 4 : Computing forces and Energies

After PASS 3, we have in every box at the lowest level both the multipole expansion as well as the local expansion, thus everything to compute the far-field contribution to the forces and the potential of the system.

3.3. Tree-based methods




Figure 3.11: Situation after PASS 3

So we have for the electric field :

$$E_{FF} = \sum_{l=0}^p \sum_{m=-l}^{+l} \mu_{lm}(\mathbf{r}) \omega_{lm}(\mathbf{a}) \quad (3.26)$$

For the potential :

$$\Phi_{FF} = - \sum_{l=0}^p \sum_{m=-l}^{+l} \mu_{lm}(\mathbf{r}) \nabla_{\mathbf{a}_i} [a_i^l \tilde{P}_{lm}(\cos \alpha_i) e^{-im\beta_i}] \quad (3.27)$$

and for the force :

$$\mathbf{F}_{FF} = - \sum_{l=0}^p \sum_{m=-l}^{+l} \mu_{lm}(\mathbf{r}) \nabla_{\mathbf{a}_i} [a_i^l \tilde{P}_{lm}(\cos \alpha_i) e^{-im\beta_i}] \quad (3.28)$$

Then the near-field contribution can be easily computed :

Chapter 4

Comparing FMM and PME accuracy

In this chapter I will more precisely explain my work done in the Lab, which consisted in comparing the PME methods, which is for example used by gromacs, and the FMM method, used by a solver made between the Julich Forschungszentrum and the Max Planck Institute for Biophysics in Göttingen.

4.1 Presentation of GROMACS

GROMACS (GRoningen MACHine for Chemical Simulations) is a simulation software used for Molecular dynamics. It was originally developed in the Biophysical Chemistry department of University of Groningen but it is now developed around the world. GROMACS is made to be as fast as possible using all possible techniques to improve its performance (MPI, GPU computing)

4.1.1 Structure of a File

The program is used by modifying text files that giving information on the structure of the system that has to be simulated and the parameters of the simulation. The files are the following

- The *.pdb file are the *Protein DataBank* file : In this file is denoted the position and the type of the particles of the system. It also gives the size of the simulation box

4.1. Presentation of GROMACS

- The *.top files (namelt *topology* files) are the file where the properties of the atoms are defined. These properties are for example the charge, the Van-der-Waals parameters or the binding forces of the system. This configuration is often done with force field files such as AMBER or CHARMM
- The *.mdp (MD parameter file) files are defining the physical and computational parameters of the simulation. It is defining the methods used for the simulation. In our case we are mostly interested in the electrostatics part of the computation ; it is then possible to select the method (PME or Cut-Off methods) and the parameters for the PME

4.1.2 PME parameters

In this paragraph I will explain the parameters it is possible to play with:

CutOff The first parameter is the CutOff : It allows to set the Cut-Off radius for the Cut-Off method, but it also sets the difference between the direct part and the reciprocal part in the PME method

Fourier Spacing The other important parameter is the fourier spacing. I remind that in the PME method a 3D FFT is done in order to compute the energies and the forces of the system. The FFT is so computed on a grid : The dimension of the grid is given by the *Fourier Spacing* parameter.

PME Order The PME order gives the order of the charge interpolation on the grid. For instance, PMEorder= 4 corresponds to a cubic interpolation.

4.1.3 Command to launch a GROMACS simulation

The workflow to launch a GROMACS Simulation is the following :

First generate a .tpr binary file containing all the information about the simulation. The file is processed using the *grompp* (Gromacs Preprocessor). The command is :

```
grompp -f mdpfile
-p topFile.top
-c pdbFile.pdb
-o tprFile.tpr
```

where the input files are the *.mdp , *.top and *.pdb (Respectively properties of the Simulation, of the atom, and the position of the Atoms). The *.tpr file is the output file.

4.2. Presentation of fmsolvr

It is then possible to run the simulation using the program `mdrun`, which as its name stands, runs the md simulation. A possible example is :

```
mdrun -s tprfile.tpr
      -ntomp = 4
      -ntmpi = 4
      -nsteps= 0
```

Where the `.tpr` File is now the input file, containing everything about the simulation, `-nsteps` is the number of time steps needed (In the case of this Intership, no time intergration is needed as just the forces are required). It is also possible to choose the number of CPU cores needed for the simulation (Using MPI and OpenMP).

4.2 Presentation of fmsolvr

The software used to compute the electrostatic forces with the FMM is not `gromacs` (at least not for the moment). It is developped by both the Max-Planck Institut for Biophysical Chemistry and the Jülich Forschungszentrum . In the following chapters, we will call this code *fmsolvr*.

The codebase is mostly written in C++. There exists several git branches for the program, allowing different versions of the system. The First version is a sequential version, which is the "basic" version of the FMM code . There also exists a version which allows periodic boundary conditions : We will use this version a lot as we need to compare it to the PME, which, by construction, uses periodic boundary conditions.

The input file, is a `*.hpp` file containing 4 arrays of size N , where N is the number of atoms in the simulation : one charge array q , and three array x, y, z for the positions of the atoms.

4.2.1 FMM Parameters

The simulation can be launched using the following command :

```
DEPTH=${DEPTH}  MULTIPOLEORDER=${MULTIPOLEORDER}  WS=${WS}
OPENBOUNDARY=0  UNITBOX=${UNITBOX}  CENTER=c
./fmmtest $inputFile.qxyz $outputFile.dat \\\
```

```
DEPTH=$DEPTH  MULTIPOLEORDER=$MULTIPOLEORDER  WS=$WS
OPENBOUNDARY=0  UNITBOX=$UNITBOX  CENTER=c
./fmmtest $inputFile.qxyz $outputFile.dat
```

4.3. Making GROMACS and fmsolvr comparable

The environment variables are the following :

DEPTH The maximal subdivision of the space : if DEPTH= n the space will be divided in 8^n boxes.

MULTIPOLEORDER The Order of truncation in a the series, for instance :

$$\frac{1}{|r-a|} \simeq \sum_{l=0}^{\textcolor{red}{p}} \sum_{m=-l}^{+l} \frac{(l-m)!}{(l+m)!} \frac{a^l}{r^{l+1}} P_{lm} \cos(\alpha) P_{lm} \cos(\theta) e^{-im(\beta-\alpha)}$$

15:20 - 16:00 - Gautier VAILLANT

where the multipole order is the red $\textcolor{red}{p}$ in the first summation.

WS This gives the separation criterion $ws \geq 1$, which separates the far-field computation from the near space.

OPENBOUNDARY if OPENBOUNDARY = 0, periodic boundary conditions will be used, if OPENBOUNDARY=1, then openboundaries will be used.

UNITBOX Gives the size of the simulation box (in nm).

4.3 Making GROMACS and fmsolvr comparable

The first part, if we want to make the PME and the FMM comparable, is to use the same simulation for both programs, hence requiring coding some tools transforming a GROMACS-compatible file to fmsolvr-compatible one. Then, we saw that the results we obtained weren't good enough as one system was using some dipole correction, so we needed to implements the dipole correction to the FMM system.

4.3.1 File manipulation

In this section will be explained the file modifications needed in order to have the simulations comparable on both systems.

4.3. Making GROMACS and fmsolvr comparable

Making GROMACS only compute electrostatics forces

In this Internship we just want to compute the electrostatic forces on our system. However, what we obtain in the GROMACS output file is the sum of all forces on each atom. These forces may include forces such as for instance Lennard-Jones potentials. ($V_{LJ} = 4\epsilon[(\frac{\sigma}{r})^{12} - (\frac{\sigma}{r})^6]$).

So the first thing to do is to modify the topology file (*.top), so there is no more Lennard Jones potential at all ; this snippet is added to the *.top file:

```
[ atomtypes ]
type atnum      mass      charge ptype      sigma  epsilon\\
CLA      17      35.450000    0.000  A    0.00000000000000  0.000000\\
SOD      11      22.989770    0.000  A    0.00000000000000  0.000000\\
```

In the expression of the potential, we set $\epsilon = 0$ and $\sigma = 0$, so it gives $V_{LJ} = 0$. So we have modified the files such that there is only the electrostatic interaction at play in the simulation.

Assure compatibility between *.qxyz and *.pdb files

The next thing to do is to have the same simulation system for both methods : So I needed to make a few scripts to transform a *.pdb file in a *.qxyz or a *.hpp file.

The scripts are written in python, the workflow is the following:

$*.hpp \xleftarrow{\text{qxyz2hpp.py}} *.qxyz \xrightarrow{\text{qxyz2gromacs.py}} *.pdb$

So it is now possible to move the positions of the atoms one software to another software. (PME to FMM or FMM to PME)

Output positions and forces to a Text file for GROMACS

The last preliminary thing to do is to make the GROMACS Simulation print the positions and the forces in an easy and detailed way. We decided to modify the Gromacs code to add after the so-called `do_force(...)` function, which computes the forces for the system, a routine that prints the force as well as the positions in order to be easily used afterwards.

4.4. First Comparisions and Dipole Correction

4.4 First Comparisions and Dipole Correction

4.4.1 Silicamelt Simulation

The first thing we did afterwards it to take a good precision for both methods and compare the forces.

The model we are using is called the "Silicamelt" system which is a system composed of SiO_2 atoms : It is a system compased of about 100000 atoms, the silicon atoms (Si) have a $+2.4e$ charge and the Oxygen has a charge of $-1.2e$, where e is the elementary charge $e \approx 1.6 \cdot 10^{-19}C$. The size of the simulation box is 124.120 nm.

The parameters for the PME simulation are cutoff=1.2nm, PME order=12 and fourier spacing=0.015 nm.

The paramerters for the FMM are : Depth=4, Multipole Order=40, and WS(Separation Criterion) =4. The simulations is done with periodic boundaries

The relative error is computed the following way:

$$error_{relative} = \frac{F_{FMM} * F_{PME}}{F_{PME}} \quad (4.1)$$

Then the following histogram can be plotted:

4.4. First Comparisons and Dipole Correction




Figure 4.1: Histogram representing the relative error for a silicamelt simulation : The histogram is computed before any dipole correction.

4.4. First Comparisions and Dipole Correction




Figure 4.2: Figure representing the Magnitude of the force in the x-axis and the error in relative error in the y-axis. It is possible to see that the biggest errors are when the magnitude of the force is smaller.

Some statistical information about the distribution gives us a maximum error of about 56.35%, this maximum error is much bigger than expected and is therefore not acceptable.

As shown figure (4.2) we study the distribution of errors according to the magnitude of system we can see that the biggest errors are some smaller magnitudes.

4.4.2 Dipole correction

Study of a two-particle system

In order to understand these big errors, we first decided to study how 2 particles behave. Then if the errors are small enough, it is possible to move

4.4. First Comparisions and Dipole Correction

to bigger systems.

We also observed that if we set 4 particles in a square in a quadrupole, so there is no dipole here, the value of the FMM system and the PME system are matching

The hypothesis is that the dipole moment may be corrected in one system and not in another system. So the idea is to take two particles, one positively charged and the other negatively charged ; the two particles will form a dipole, namely :

$$\vec{p} = q\vec{d} \quad (4.2)$$

where \vec{d} is the displacement vector pointing from the negative charge to the positive charge.

Then we can vary the distance between the two particles, compute the force using both the FMM method and the PME method and compare the force.




Figure 4.3: Figure representing the distance between two particles the x-axis (here 55,60,65,70 nm) and the Magnitude of the force for PME (Blue Square), FMM without correction (Black point) and with correction (red star). It is possible to see that the dipole correction makes the FMM magnitude much closer to the PME.

4.4. First Comparisions and Dipole Correction

The dipole correction is done the following way :

It is possible to modify the potential in the FMM code using the following equation:

$$\nabla_i E_{\text{after correction}} = \nabla_i E_{\text{before correction}} - \frac{4\pi}{3 \cdot V} q_i \mathbf{d} \quad (4.3)$$

where V is the volume of the simulation box, q_i the charge of the particle and \mathbf{d} is the dipole moment.

The dipole moment can be computed from the multipole moments, which contains the dipole moment : The multipole moments are contained in a triangular array (because of symmetry and memory space reasons) the following way :




Figure 4.4: Structure of the array containing the multipoles $\omega_{lm}\mathbf{a}$) : the 'm' values are on the y-axis and the 'l'-values are on the x-axis. The dipole moment is contained in the boxes (1,1), (1,0) and (1,-1)

As showed in figure (4.4), it is possible to obtain the dipole moment from the multipole ones, using the following formula :

$$\begin{cases} \vec{d}_x = -2 * \Re(\omega_{1,1}) \\ \vec{d}_y = 2 * \Im(\omega_{1,1}) \\ \vec{d}_z = -\Re(\omega_{1,0}) \end{cases} \quad (4.4)$$

Then the correction can be applied using equation (4.3)

4.4. First Comparisions and Dipole Correction

The code used for this correction, in C++11, is the following :

```
for (size_t i = 0; i < n; ++i) {

    auto cell_volume = (abc.a).x * (abc.b).y * (abc.c).z; // Only
        works with square Cell Unit
    std::cout << "Periodic Vector = " << (abc.a).x << std::endl;
    std::cout << "Volume " << cell_volume << std::endl;

    std::cout << Real3(ordered_particles[i]) << "\t"
        <<reference_center << std::endl;

    auto dr = Real3(ordered_particles[i]) - reference_center;
    std::cout << "dr = " << dr << std::endl;

    const auto& omega = tree_omega[0][0];
    dipole_moment.x = -omega.get(1,1).real()*2;
    dipole_moment.y = omega.get(1,1).imag()*2;
    dipole_moment.z = -omega.get(1,0).real();

    // Potential Correction
    auto correction = (4.0*PI/(3.0 * cell_volume )) * (dr.x *
        dipole_moment.x) + (dr.y * dipole_moment.y) + (dr.z *
        dipole_moment.z);
    std::cout << "Potential Before = " << potential[i] << std::endl;
    std::cout << "Correction = " << correction << std::endl;
    potential[i] -= correction;
    std::cout << "Potential After = " << potential[i] << std::endl;
    /*
    efield[i].x -= 4*PI/3 * dipole_moment.x;
    efield[i].y -= 4*PI/3 * dipole_moment.y;
    efield[i].z -= 4*PI/3 * dipole_moment.z;
    */

    Real q = -(-ordered_particles[i].s); // negative gradient
    force[i] = efield[i] * q - dipole_moment *
        (4.*PI/(3.0*cell_volume)) * q ;
    Ec2 += q * potential[i];
}
```

4.4. First Comparisions and Dipole Correction

```
Ec += Ec2 * 0.5;
```

Back to the silicamelt simulation with dipole correction

Now we can relaunch the simulation with the same parameters as before, including the dipole correction. We obtain now a maximum error of 3.52%, which is a much bigger improvement compared to the 56.35% of the former simulation.




Figure 4.5: Histogram representing the relative error for a silicamelt simulation : The histogram is computed with the dipole correction. The scale in x-direction is the same as in figure (4.1)

4.5. Comparing with an analytical solution : the NaCl system




Figure 4.6: Figure representing the Magnitude of the force in the x-axis and the error in relative error in the y-axis. The blue points represent the force before the dipole correction, then red points after the dipole correction

4.5 Comparing with an analytical solution : the NaCl system

One other thing we want to try is to see the convergence of the scheme. The idea is to know which set of parameters for the FMM has the same accuracy than a given set of parameters for the PME method.

In order to study that, we want to have a system which is analytically solvable with periodic boundary conditions : That is why we decided to use a cristal as a system.

4.5.1 Generating the system

The system used is based on a NaCl cristal structure. As a cristal is a stable structure, the force acting on each atom has to be equal to 0. The potential felt by one atom can also be computed with a series ; this series converges

4.5. Comparing with an analytical solution : the NaCl system

to a value called the **Madelung Constant** which is equal for this cristal structure to approximately 3.495 V.

The generation is done with a python script writing the charges and the positions of the atoms in a *.qxyz file, then translated into a pdb file to obtain cristal structure necessary for GROMACS.




Figure 4.7: NaCl system generated with the script described above : The Blue atoms are Na atoms and the Green ones are Cl atoms. The Following image has been rendered using the visualization software VMD (Visual Molecular Dynamics)

4.5.2 Some explanations on the Owl cluster

Now I will quickly explain how to submit a certain program to the cluster of the lab, which is called *owl*.

Owl is the cluster of the lab used for the MD simulations : It contains about 12500 processor cores and 70 GPUs. The subnit system is done using

4.5. Comparing with an analytical solution : the NaCl system

sgc.

The cluster is accessible via an ssh connection : then it is possible to submit to the cluster using the following script:

```
#!/bin/bash
#$ -S /bin/bash
#$ -l mem_free=10G,h_data=5G,h_rt=96:00:00,h_cpu=96:00:00
#$ -ac SLOTSCPU=1
#$ -j yes
#$ -v TMPDIR=$TMPDIR/add_ion.$JOB_ID
#$ -pe *_fast 1

submit program
```

For the PME, there already exists a script called `g_submit`, which allows to submit the simulation defined *.tpr file to the cluster. An example of a script is given below :

```
g_submit -N ${JOB_NAME}\
        -days ${NUMBER_DAYS}\
        -double\
        -source $GMXPATH\
        -s ${tprfile}\
        -nomail\
```

this allows the script to choose the version of gromacs compiled with double precision (`-double`) using the `tprfile` for a certain number of days `NUMBER_DAYS`

4.5.3 Error plots

One way to measure the accuracy is to compare the force obtained experimentally with the analytic solution. In order to test the accuracy of one set of parameters, we will compute the sum of the euclidian norm for each atom :

$$\text{error} = \sum_{i=0}^N \|\vec{F}_i - \vec{0}\| = \sum_{i=0}^N \|\vec{F}_i\| \quad (4.5)$$

4.5. Comparing with an analytical solution : the NaCl system




Figure 4.8: Figure representing the Magnitude of the force in the x-axis and the error in relative error in the y-axis. The blue points represent the force before the dipole correction, then red points after the dipole correction

Conclusion

Bibliography

- [1] A. Arnold and C. Holm. Efficient methods to compute long-range interactions for soft matter systems. In *Advanced computer simulation approaches for soft matter sciences II*, pages 59–109. Springer, 2005.
- [2] S. Boresch and O. Steinhauser. Presumed versus real artifacts of the ewald summation technique: The importance of dielectric boundary conditions. *Berichte der Bunsengesellschaft für physikalische Chemie*, 101(7):1019–1029, 1997.
- [3] H. Dachsel. Corrected article:“an error-controlled fast multipole method”[j. chem. phys. 131, 244102 (2009)]. *The Journal of chemical physics*, 132(11):119901, 2010.
- [4] T. Darden, D. York, and L. Pedersen. Particle mesh ewald: An $n \log(n)$ method for ewald sums in large systems. *The Journal of chemical physics*, 98(12):10089–10092, 1993.
- [5] L. Greengard. Fast algorithms for classical physics. *Science*, 265(5174):909–914, 1994.
- [6] B. Hess, D. van Der Spoel, and E. Lindahl. Gromacs user manual version 5.0.4. *University of Groningen, Netherland*.
- [7] A. T. Ihler. An overview of fast multipole methods. *Area Exam*, April, 2004.
- [8] I. Kabadshow. *Periodic Boundary Conditions and the Error-Constrolled Fast Multipole Method*. PhD thesis, Juelich Forschungszentrum, 2012.
- [9] C. Kutzner, R. Apostolov, B. Hess, and H. Grubmüller. Scaling of the gromacs 4.6 molecular dynamics code on supermuc. *Advances in Parallel Computing*, 25, 2013.

Bibliography

- [10] S. Pronk, S. Páll, R. Schulz, P. Larsson, P. Bjelkmar, R. Apostolov, M. R. Shirts, J. C. Smith, P. M. Kasson, D. van der Spoel, et al. Gromacs 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. *Bioinformatics*, page btt055, 2013.
- [11] A. Y. Toukmaji and J. A. Board. Ewald summation techniques in perspective: a survey. *Computer physics communications*, 95(2):73–92, 1996.

Appendices

Appendix A

Glossary and Abbreviations

MD Molecular Dynamics

GROMACS GRoningen MACHine for Chemical Simulations

PME Particle Mesh Ewald

FMM Fast Multipole Method

Owl MPI-BPC Cluster Used to make the Simulations

MPI Message Passing Interface

Omp OpenMP

Appendix B

Structure of Gromacs files

B.1 pdb file

```
TITLE Generated by qxyz2gromacs.py
REMARK THIS IS A SIMULATION BOX
CRYST1 100.000 100.000 100.000 90.00 90.00 90.00 P 1 1
MODEL 1
ATOM 1 NA NA 2 0.000 0.000 0.000 1.00
ATOM 2 NA NA 4 5.000 5.000 0.000 1.00
ATOM 3 NA NA 6 5.000 0.000 5.000 1.00
ATOM 4 NA NA 8 0.000 5.000 5.000 1.00
ATOM 5 NA NA 10 0.000 0.000 10.000 1.00
ATOM 6 NA NA 12 5.000 5.000 10.000 1.00
ATOM 7 NA NA 14 5.000 0.000 15.000 1.00
ATOM 8 NA NA 16 0.000 5.000 15.000 1.00
ATOM 9 NA NA 18 0.000 0.000 20.000 1.00
...
...
TER
ENDMDL
```

B.2 top file

```
; topology file for a NaCl melt
;

; Include forcefield parameters
#define _FF_CHARMM
```

B.2. top file

```
[ defaults ]
; nbfunc          comb-rule      gen-pairs      fudgeLJ fudgeQQ
1                yes            1.0          1.0

[ atomtypes ]
;type atnum      mass      charge ptype      sigma  epsilon
CLA    17      35.450000    0.000  A  0.000000000000  0.00000
SOD    11      22.989770    0.000  A  0.000000000000  0.00000

; Include topology for ions
[ moleculetype ]
; molname      nrexcl
NA              1

[ atoms ]
; id      at type      res nr  residu name at name  cg nr  charge
1    SOD      1    NA      NA      1    1

[ moleculetype ]
; molname      nrexcl
CL              1

[ atoms ]
; id      at type      res nr  residu name at name  cg nr  charge
1    CLA      1    CL      CL      1   -1

[ system ]
; Name
NaCl melt

[ molecules ]
; Compound
NA      500
CL 500
```


B.3. mdp file (longe-range electrostatics part)

B.3 mdp file (longe-range electrostatics part)

```
; OPTIONS FOR ELECTROSTATICS AND VDW

; Method for doing electrostatics
coulombtype           = PME
coulomb-modifier       = Potential-shift-Verlet
rcoulomb_switch       = 0
rcoulomb              = 0.1

; Relative dielectric constant for the medium and the reaction field
epsilon_r             = 1
epsilon-rf            = 0

; Method for doing Van der Waals
vdw-type              = Cut-Off
vdw-modifier          = Potential-shift-Verlet

; cut-off lengths
rvdw_switch           = 0
rvdw                  = 0.1

; Apply long range dispersion corrections for Energy and Pressure
DispCorr              = No

; Extension of the potential lookup tables beyond the cut-off
table-extension       = 1

; Separate tables between energy group pairs
energygrp-table       =

; Spacing for the PME/PPPM FFT grid
fourierspacing        = 0.01

; FFT grid size, when a value is 0 fourierspacing will be used
fourier-nx            = 0
fourier-ny            = 0
fourier-nz            = 0

; EWALD/PME/PPPM parameters
pme-order             = 4
```

B.3. mdp file (longe-range electrostatics part)

```
ewald-rtol          = 1E-5
ewald-rtol-lj        = 0.001
lj-pme-comb-rule     = Geometric
ewald-geometry       = 3d
epsilon-surface      = 0
```