Analytical Chemistry Cumulative Examination
February 6, 2009

1. Electrospray ionization mass spectrometry is an exceptionally useful technique for analysis of high mass chemicals. Please answer each of the following questions about this technique.

a) Prior to introduction of this method, quadrupole mass analyzers were generally limited to analysis of molecules with molecular weights of 2,000 Daltons or lower. ESI-MS has greatly changed this situation, particularly for analysis of proteins. Please explain why this is so and how ESI makes this possible. What are current mass limits for protein analysis? (20 points)

b) Data processing of electrospray mass spectra can have a number of complications. Please explain what those are and what methods (both computational and experimental) can be used to overcome those problems. (20 points)

c) There are various models for the electrospray ionization process. Two of these are Dole's charged residue model and Iribarne and Thomson's ion extraction ("Coulomb explosion") model. Explain, with the help of a suitable, clearly labeled, diagram the basic idea of the Iribarne and Thomson model for the mechanism of electrospray. (20 points)

2. In an early paper devoted to electrospray Fourier transform mass spectrometry McLafferty and his students (*J. Amer. Soc. Mass Spectrom.*, **1993**, 4 , 557-565) covered some of the advantages of this method. Please answer the following questions. (a) What is the principle advantage of using the FTMS over a quadrupole MS for this application? (b) What order of magnitude resolving power can be obtained for protein molecular ions in using FTMS? (c) Charge assignment can be a difficult problem in electrospray MS. How can FTMS be used to advantage for such assigments? (20 points).

3. A major application of ESI mass spectrometry is characterization of viruses with molecular masses in the range from 3.6 megadalton to gigadaltons. This development was described in a paper by Hogan, et. al. in Analytical Chemistry, 78, 844-852 (2006). A significant part of this research involved use of a differential ion mobility analyzer. How is this possible and how does it work? Be explicit and describe how the differential analyzer works. (20 points)