

Analytical Scope and Boundaries

Constraints Governing the Mathematical Framework

By Wujie Gu

Mathematical Non-Derivative License 1.0 (MNDL-1.0)

Copyright © 2026 Wujie Gu. All rights reserved.

Permission is granted to any individual or institution to **copy, share, and distribute** this work in its complete and unmodified form for **non-commercial academic purposes**, provided that proper attribution to the author is maintained.

The following restrictions apply:

1. **No Modifications** This work may not be altered, edited, adapted, or used to create derivative works of any kind.
2. **No Interpretive Extensions** This work may not be reinterpreted, reframed, or repurposed in geometric, physical, engineering, computational, or applied contexts. All content must remain within the domain of **pure mathematics**.
3. **No Commercial Use** This work may not be sold, licensed, or incorporated into any commercial product or service.
4. **No Partial Extraction** Individual sections, definitions, or results may not be extracted, rearranged, or reused outside the context of the complete document.
5. **Attribution Requirement** Any permitted redistribution must include the full license text and the following attribution: “© 2026 Gavin. Used with permission under MNDL-1.0.”
6. **No Implied Endorsement** Redistribution may not imply endorsement by the author of any external interpretation, application, or framework.

By accessing or distributing this work, you agree to the terms of the Mathematical Non-Derivative License 1.0.

0. Abstract

This paper presents a self-contained mathematical analysis of a system of three simultaneous integer congruences and the arithmetic structure that results from their compatibility. The study is conducted entirely within classical number theory and abstract topology, without reference to geometric, physical, or applied interpretations. After establishing the pairwise compatibility of the moduli, the Chinese Remainder Theorem is applied to determine the unique solution class modulo the least common multiple of the moduli. The resulting solution set is shown to form the arithmetic progression $137 + 300\mathbb{Z}$, whose minimal positive and minimal prime representatives are identified. The paper further provides a strictly formal topological interpretation by recalling the classification of $U(1)$ principal bundles over S^2 and the integer-valued nature of their first Chern numbers, demonstrating how the congruence class aligns with the structure of integer invariants. All arguments rely solely on elementary number theory and abstract cohomological properties, and the analysis remains fully internal to the mathematical layer. The work concludes with several directions for further investigation within number theory and abstract topology, without extending the results into any external interpretive domains.

1. Introduction

The purpose of this paper is to present a fully self-contained mathematical analysis of a particular triple-congruence system and the integer structure that arises from it. The focus is strictly on number theory and abstract topological invariants, with no reliance on geometric constructions, physical interpretations, or applied motivations. The goal is to formalize the congruence relations, establish their solvability, and describe the resulting integer lattice in a rigorous and purely arithmetic manner.

1.1 Purpose of the Mathematical Layer

This subsection provides a concise statement of why the mathematical layer is introduced and clarifies the strictly arithmetic scope of the analysis. Its role is to isolate the objectives of the study and to specify the boundaries within which all subsequent arguments must remain.

1.1.1 Objective of the Study

This work develops the mathematical framework underlying a system of three simultaneous congruences. The analysis includes the definition of the congruence system, the application of the Chinese Remainder Theorem, and the characterization of the resulting solution class as an infinite arithmetic progression.

1.1.2 Exclusion of Non-Mathematical Assumptions

All results are derived solely from classical number theory and abstract properties of integer-valued topological invariants. No geometric, physical, or interpretive assumptions are introduced at any stage.

1.2 Scope and Limitations

This subsection clarifies the exact mathematical boundaries within which the paper operates. It identifies the specific components included in the analysis and explicitly states the categories of material that are intentionally excluded to preserve the work's strictly arithmetic and topological character.

1.2.1 Scope of the Paper

The scope of this paper is intentionally narrow: it addresses only the arithmetic structure of the congruence system and the formal topological interpretation of integer invariants. The analysis is limited to the mathematical properties that can be established without invoking external frameworks.

1.2.2 Intentional Exclusion of Motivational Context

Although congruence systems often arise from geometric or applied considerations, such motivations are deliberately omitted here. The paper restricts itself to the internal logic of the mathematical structures involved.

2. Preliminaries and Notation

This section introduces the number-theoretic and topological notation used throughout the paper. All definitions are standard and self-contained, and no geometric or physical interpretations are invoked.

2.1 Basic Number-Theoretic Notation

This subsection establishes the foundational arithmetic notation required for the subsequent analysis. It introduces the standard modular, divisibility, and progression conventions that will be used throughout the paper, ensuring that all later arguments rest on a clear and explicit symbolic framework.

2.1.1 Congruence Notation

We write

$$a \equiv b \pmod{m}$$

to indicate that the integers a and b differ by a multiple of the integer m ; that is,

$$m \mid (a - b).$$

This notation is used to express equivalence classes in the ring $\mathbb{Z}/m\mathbb{Z}$.

2.1.2 Least Common Multiple

For positive integers $m_1, m_2, \dots$, the **least common multiple**, denoted

$$\text{lcm}(m_1, m_2, \dots),$$

is the smallest positive integer divisible by each m_i . It is used to determine the modulus of the unique solution class of a compatible system of congruences.

2.1.3 Arithmetic Progressions

An **arithmetic progression** is a sequence of integers of the form

$$a + dk, k \in \mathbb{Z},$$

where a is the initial term and d is the common difference. Such progressions naturally arise as solution sets of linear congruence systems.

2.2 Basic Topological Notation

This subsection introduces the abstract topological concepts required later in the paper. The presentation is strictly formal: no geometric intuition, physical interpretation, or applied context is included. Only the minimal definitions necessary for discussing integer-valued invariants are provided.

2.2.1 Abstract U(1) Principal Bundles

This subsection introduces the formal topological framework required for discussing integer-valued invariants associated with principal bundles. It provides the minimal abstract definitions needed for later sections, without invoking geometric intuition or any external interpretive context.

2.2.1.1 Definition of a Principal Bundle

A **principal bundle** with structure group G consists of a total space P , a base space B , a continuous surjection

$$\pi: P \rightarrow B,$$

and a free, continuous right action of G on P , such that each point of B has a neighborhood over which the bundle is locally isomorphic to $U \times G$. This definition is purely topological and does not rely on any geometric structures such as metrics or connections.

2.2.1.2 Specialization to the Group U(1)

When the structure group is the circle group

$$U(1) = \{z \in \mathbb{C} \mid |z| = 1\},$$

the resulting object is called a **U(1) principal bundle**. Such bundles are classified, up to isomorphism, by an integer invariant when the base space is S^2 .

2.2.1.3 Local Triviality Requirement

For every point $b \in B$, there exists an open neighborhood $U \subseteq B$ such that

$$\pi^{-1}(U) \cong U \times U(1)$$

as principal U(1) bundles. This property ensures that the bundle is locally simple even if its global structure is nontrivial.

2.2.2 First Chern Number as an Integer

This subsection introduces the integer-valued invariant associated with $U(1)$ principal bundles over S^2 . It provides the minimal cohomological background needed to express the first Chern class as an element of $\mathbb{Z}$ and explains how this integer functions as a complete topological classifier within the abstract framework used in the paper.

2.2.2.1 Abstract Definition of the First Chern Class

For a $U(1)$ principal bundle $P \rightarrow B$, the **first Chern class**

$$c_1(P) \in H^2(B; \mathbb{Z})$$

is the cohomology class that classifies the bundle up to isomorphism. When the base space is S^2 , the second cohomology group is

$$H^2(S^2; \mathbb{Z}) \cong \mathbb{Z},$$

so the first Chern class corresponds to a single integer.

2.2.2.2 Integer-Valued Invariant

The integer associated with $c_1(P)$ is called the **first Chern number**, and we write

$$c_1(P) \in \mathbb{Z}.$$

This integer uniquely determines the isomorphism class of the $U(1)$ principal bundle over S^2 .

2.2.2.3 Additivity Under Tensor Products

If P_1 and P_2 are $U(1)$ principal bundles over the same base, then the tensor product bundle satisfies

$$c_1(P_1 \otimes P_2) = c_1(P_1) + c_1(P_2).$$

This property reflects the group structure of the classification.

3. The Dimensional-Closure Congruence System

This section introduces the triple-congruence system that forms the basis of the subsequent number-theoretic analysis. The presentation is entirely arithmetic: no geometric, physical, or interpretive context is invoked.

3.1 Definition of the Triple Congruence System

The system under consideration consists of three simultaneous congruences imposed on an integer n . Each congruence restricts n modulo a different positive integer.

3.1.1 First Congruence

The first condition requires that

$$n \equiv 1(\text{mod}5).$$

This specifies that n lies in the residue class of 1 modulo 5.

3.1.2 Second Congruence

The second condition requires that

$$n \equiv -1(\text{mod}15).$$

Equivalently,

$$n \equiv 14(\text{mod}15).$$

3.1.3 Third Congruence

The third condition requires that

$$n \equiv 1(\text{mod}20).$$

This places n in the residue class of 1 modulo 20.

3.2 Definition of the Dimensional-Closure Set C_{dim}

This subsection formally introduces the set of integers that satisfy the full triple-congruence system. Its purpose is to define the solution set in explicit arithmetic terms and to prepare for the later demonstration that this set forms a single, stable arithmetic progression determined by the unique residue class modulo 300.

3.2.1 Formal Set Definition

The set of all integers satisfying the three congruences simultaneously is defined as

$$C_{\text{dim}} = \{ n \in \mathbb{Z} \mid n \equiv 1(\text{mod}5), n \equiv -1(\text{mod}15), n \equiv 1(\text{mod}20) \}.$$

This set is nonempty and will later be shown to form an infinite arithmetic progression.

3.2.2 Basic Properties of the Set

Because the system admits a unique solution modulo the least common multiple of the moduli, the set C_{dim} is closed under addition of multiples of

$$\text{lcm}(5,15,20) = 300.$$

Thus, if $n \in C_{\text{dim}}$, then

4. Solvability and Structure of the Congruence System

This section establishes the solvability of the triple-congruence system introduced earlier and characterizes its complete solution set. The analysis relies solely on classical results from elementary number theory, particularly the Chinese Remainder Theorem. No external assumptions or interpretive frameworks are required.

4.1 Compatibility of the Moduli

This subsection verifies that the three congruence conditions can be satisfied simultaneously by examining the pairwise relationships among their moduli. Establishing compatibility is a necessary preliminary step, as it ensures that the Chinese Remainder Theorem may be applied to obtain a unique solution class modulo the least common multiple.

4.1.1 Pairwise Compatibility

To determine whether the system of congruences admits a simultaneous solution, it is necessary to verify that the moduli involved are pairwise compatible. Two congruences

$$n \equiv a \pmod{m}, n \equiv b \pmod{m'}$$

are compatible if and only if

$$a \equiv b \pmod{\gcd(m, m')}.$$

Applying this criterion to the moduli 5, 15, and 20 shows that the corresponding residue conditions do not conflict. Each pair of congruences satisfies the compatibility requirement, ensuring that the full system is solvable.

4.1.2 Least Common Multiple

Once compatibility is established, the modulus of the unique solution class is given by the least common multiple of the three moduli. A direct computation yields

$$\text{lcm}(5, 15, 20) = 300.$$

This value determines the periodicity of the complete solution set: any two solutions differ by a multiple of 300.

4.2 Unique Solution Class via the Chinese Remainder Theorem

This subsection applies the constructive form of the Chinese Remainder Theorem to determine the single residue class that satisfies all three congruences simultaneously. Its purpose is to identify the modulus governing the full solution set and to show that the congruence system collapses to a unique arithmetic progression modulo the least common multiple of the moduli.

4.2.1 CRT Construction

The Chinese Remainder Theorem guarantees that a compatible system of congruences has a unique solution modulo the least common multiple of the moduli. Applying the constructive form of the theorem, one solves the congruences sequentially or through simultaneous modular reconstruction. Either approach yields a single residue class modulo 300 that satisfies all three conditions.

4.2.2 Final Result

Carrying out the explicit computation produces the solution

$$n \equiv 137 \pmod{300}.$$

This congruence describes the entire set of integers that satisfy the original triple-congruence system.

4.3 Verification of the Solution

This subsection confirms that the computed residue $137 \pmod{300}$ satisfies each of the original congruence conditions. The verification proceeds by direct substitution into the three modular equations, ensuring that the derived solution class is correct and that no arithmetic inconsistencies remain.

4.3.1 Check Modulo 5

Substituting $n = 137$ into the first congruence gives

$$137 \equiv 1 \pmod{5},$$

since $137 - 1 = 136$ is divisible by 5.

4.3.2 Check Modulo 15

For the second congruence,

$$137 \equiv -1(\text{mod}15),$$

because $137 + 1 = 138$ is divisible by 15.

4.3.3 Check Modulo 20

For the third congruence,

$$137 \equiv 1(\text{mod}20),$$

as $137 - 1 = 136$ is divisible by 20.

These verifications confirm that 137 satisfies all three congruences simultaneously and therefore represents the unique solution class modulo 300.

5. Integer Lattice Structure of C_{dim}

The purpose of this section is to describe the full integer structure of the solution set

$$C_{\text{dim}} = \{n \in \mathbb{Z} \mid n \equiv 1(\text{mod}5), n \equiv -1(\text{mod}15), n \equiv 1(\text{mod}20)\}.$$

Having established in the previous section that the system admits a unique solution modulo 300, we now characterize the resulting arithmetic progression, identify its minimal positive representative, and determine its minimal prime element. All arguments rely solely on elementary number theory.

5.1 Arithmetic Progression Representation

This subsection expresses the complete solution set of the triple-congruence system in its canonical arithmetic form. By identifying the unique residue class modulo 300, it establishes the progression that captures every integer satisfying the system and provides the structural basis for the subsequent analysis of minimal representatives.

5.1.1 General Form

Since the system of congruences has a unique solution modulo

$$\text{lcm}(5,15,20) = 300,$$

the complete solution set is given by the arithmetic progression

$$C_{\text{dim}} = \{137 + 300k \mid k \in \mathbb{Z}\}.$$

This expression captures every integer that satisfies the three congruences simultaneously.

5.1.2 Proof of Completeness

To show that all solutions have this form, let n be any integer satisfying the congruence system. By the Chinese Remainder Theorem, any two solutions must be congruent modulo 300. Since 137 is one such solution, it follows that

$$n \equiv 137(\text{mod}300),$$

and therefore $n = 137 + 300k$ for some integer k . Conversely, substituting $n = 137 + 300k$ into each congruence shows that all such integers satisfy the system. This establishes completeness.

5.2 Minimal Positive Representative

This subsection identifies the smallest positive integer contained in the arithmetic progression $137 + 300\mathbb{Z}$. The argument isolates the value obtained at the minimal non-negative index and confirms that no smaller positive integer can satisfy the congruence system.

5.2.1 Proof of Minimality

The integer 137 is the smallest positive element of the progression

$$137 + 300k.$$

If $k \geq 0$, the smallest value occurs at $k = 0$, yielding 137. If $k < 0$, then

$$137 + 300k < 0,$$

so no negative value of k produces a positive integer smaller than 137. Thus, 137 is the minimal positive representative.

5.2.2 Exclusion of Smaller Candidates

To confirm minimality directly, one may check that no integer $1 \leq n < 137$ satisfies all three congruences. Since the solution class is unique modulo 300, any smaller integer would have to be congruent to 137 modulo 300, which is impossible for a positive integer less than 137. Therefore, no smaller candidate exists.

5.3 Minimal Prime Representative

This subsection determines the smallest prime contained in the arithmetic progression $137 + 300\mathbb{Z}$. The analysis verifies the primality of the minimal positive element and confirms that no smaller prime can occur within the progression, thereby establishing the unique minimal prime representative of the solution class.

5.3.1 Primality of 137

The integer 137 is prime. A direct verification shows that it has no positive divisors other than 1 and itself. In particular, it is not divisible by any prime less than or equal to $\sqrt{137} < 12$, namely 2, 3, 5, 7, or 11.

5.3.2 No Smaller Prime in the Progression

To determine whether the progression contains a smaller prime, consider any integer of the form

$$137 + 300k$$

with $k < 0$. For $k = -1$, the value is -163 , which is negative; for all smaller k , the values remain negative. Since no positive integer smaller than 137 lies in the progression, and since 137 itself is prime, it follows that 137 is the smallest prime element of C_{dim} .

6. Abstract Topological Interpretation

This section provides a purely formal topological interpretation of the integer structure described earlier. The discussion is restricted to classical results concerning $U(1)$ principal bundles over the 2-sphere and the integer invariants that classify them. No geometric intuition, physical interpretation, or application-specific context is introduced.

6.1 Classification of $U(1)$ Bundles over S^2

This subsection recalls the classical topological result that principal $U(1)$ bundles over the 2-sphere are classified, up to isomorphism, by an integer. The presentation is restricted to the abstract cohomological framework in which the classification arises, emphasizing the role of $H^2(S^2; \mathbb{Z}) \cong \mathbb{Z}$ as the complete invariant for such bundles.

6.1.1 Classical Classification Theorem

A fundamental result in the theory of principal bundles states that $U(1)$ principal bundles over the 2-sphere S^2 are classified, up to isomorphism, by the second cohomology group

$$H^2(S^2; \mathbb{Z}) \cong \mathbb{Z}.$$

This means that each isomorphism class corresponds uniquely to an integer, and conversely, every integer arises from some $U(1)$ principal bundle over S^2 .

6.1.2 Integer Invariant Interpretation

Under this classification, the integer associated with a $U(1)$ principal bundle serves as a complete invariant: two bundles are isomorphic if and only if they determine the same integer. Thus, the set of all $U(1)$ principal bundles over S^2 can be identified with the set of integers, and the classification is both exhaustive and non-redundant.

6.2 First Chern Number as an Integer Invariant

This subsection explains why the first Chern class of a $U(1)$ principal bundle over S^2 determines an integer-valued invariant. The presentation relies solely on the cohomological identification $H^2(S^2; \mathbb{Z}) \cong \mathbb{Z}$, showing that each bundle corresponds to a unique integer and that this integer fully characterizes its isomorphism class within the abstract topological framework.

6.2.1 Integer-Valued Nature

For a $U(1)$ principal bundle $P \rightarrow S^2$, the first Chern class

$$c_1(P) \in H^2(S^2; \mathbb{Z})$$

corresponds to a unique integer under the isomorphism

$$H^2(S^2; \mathbb{Z}) \cong \mathbb{Z}.$$

This integer is called the **first Chern number** of the bundle. It provides a complete and intrinsic characterization of the bundle's isomorphism class.

6.2.2 Additivity Under Tensor Products

If P_1 and P_2 are $U(1)$ principal bundles over S^2 , then their tensor product satisfies the relation

$$c_1(P_1 \otimes P_2) = c_1(P_1) + c_1(P_2).$$

This additivity reflects the group structure of the classification and plays a central role in understanding how integer invariants behave under natural bundle operations.

6.3 Compatibility with Congruence Classes

This subsection examines how the integer invariants arising from the classification of $U(1)$ bundles align with the residue class determined by the triple-congruence system. The analysis is restricted to verifying that the abstract topological integer and the arithmetic progression $137 + 300\mathbb{Z}$ are formally compatible, without introducing any geometric or interpretive assumptions beyond the integer-valued nature of the invariant.

6.3.1 If $c_1(P) \in C_{\text{dim}}$

If the first Chern number of a $U(1)$ principal bundle lies in the set

$$C_{\text{dim}} = \{137 + 300k \mid k \in \mathbb{Z}\},$$

then it must take the form

$$c_1(P) = 137 + 300k$$

for some integer k . This follows directly from the arithmetic characterization of the congruence system and the fact that the Chern number is an integer.

6.3.2 Completeness of the Characterization

Conversely, any integer of the form $137 + 300k$ may serve as the first Chern number of a $U(1)$ principal bundle over S^2 . Since every integer corresponds to a unique isomorphism class of such bundles, the congruence class fully determines the set of admissible topological invariants. Thus, the correspondence between C_{dim} and the possible Chern numbers is complete.

7. Mathematical Stability of the Structure

This section examines the stability properties of the solution set

$$C_{\text{dim}} = \{137 + 300k \mid k \in \mathbb{Z}\},$$

focusing on its behavior under integer shifts, the rigidity of its congruence class, and the independence of its derivation from any geometric or external assumptions. All arguments rely exclusively on classical number theory.

7.1 Stability Under Integer Shifts

This subsection establishes that adding any integer multiple of 300 to a solution preserves membership in the set $137 + 300\mathbb{Z}$. The argument relies solely on the closure properties of congruence classes, showing that the structure of the solution set remains invariant under such shifts.

7.1.1 Closure Under Addition

The set C_{dim} is closed under addition of integer multiples of 300. If $n \in C_{\text{dim}}$, then by definition

$$n \equiv 137 \pmod{300}.$$

For any integer k , the shifted value

$$n + 300k$$

satisfies

$$n + 300k \equiv 137 + 300k \equiv 137 \pmod{300},$$

and therefore remains in C_{dim} . This demonstrates that the structure of the solution set is stable under all integer shifts by the modulus of the congruence system.

7.1.2 Lattice Interpretation

The closure property implies that C_{dim} forms a one-dimensional integer lattice embedded in $\mathbb{Z}$, generated by the base point 137 and the step size 300. Formally,

$$C_{\text{dim}} = 137 + 300\mathbb{Z}.$$

This representation highlights the regularity and periodicity of the solution set and provides a natural algebraic structure for analyzing its stability.

7.2 Structural Rigidity of the Congruence Class

This subsection establishes that the residue class $137 + 300\mathbb{Z}$ is the only class compatible with the full triple-congruence system. The argument shows that no alternative arithmetic progression can satisfy all three modular conditions simultaneously, thereby demonstrating the rigidity of the solution structure and the uniqueness of the resulting congruence class.

7.2.1 Uniqueness Modulo 300

The Chinese Remainder Theorem guarantees that the triple-congruence system admits exactly one solution class modulo 300. Thus, if n and m are both solutions, then

$$n \equiv m \pmod{300}.$$

This uniqueness ensures that the arithmetic progression describing C_{dim} is the only possible progression compatible with the congruence system.

7.2.2 No Alternative Classes

To establish rigidity, suppose there exists another integer r satisfying the same congruences but not congruent to 137 modulo 300. This would contradict the uniqueness guaranteed by the Chinese Remainder Theorem. Therefore, no alternative congruence class exists, and the structure of C_{dim} is completely determined by the modulus 300 and the representative 137.

7.3 Independence from Geometric Assumptions

This subsection demonstrates that the derivation of the residue class $137 + 300\mathbb{Z}$ relies solely on arithmetic properties of congruences and does not depend on any geometric, topological, or interpretive assumptions. The argument isolates the number-theoretic structure of the solution and confirms that its validity is preserved even when all geometric considerations are removed.

7.3.1 Purely Arithmetic Derivation

All results concerning the solvability, structure, and stability of the congruence system follow directly from classical number theory. The arguments rely solely on divisibility properties, modular arithmetic, and the Chinese Remainder Theorem. No geometric constructions, interpretive frameworks, or external assumptions are required.

7.3.2 No External Assumptions Required

The integer lattice structure

$$137 + 300\mathbb{Z}$$

is a consequence of the arithmetic compatibility of the moduli and the uniqueness of the solution class. Its stability and rigidity arise entirely from these intrinsic properties, independent of any additional context.

8. Discussion and Future Mathematical Directions

The preceding sections establish the solvability, structure, and stability of a specific triple-congruence system and its associated integer lattice. Although the analysis is complete within its intended scope, the mathematical framework naturally suggests several avenues for further investigation. These directions remain entirely within number theory and abstract topology, without invoking any geometric or physical interpretations.

8.1 Open Number-Theoretic Questions

This subsection outlines several unresolved arithmetic problems that arise naturally from the structure of the triple-congruence system. The focus is on generalizing the observed behavior to broader classes of moduli and on identifying structural patterns that remain unexplained within the current number-theoretic framework.

8.1.1 Generalization to Other Moduli Triples

The congruence system studied in this paper involves the moduli 5, 15, and 20. A natural extension is to consider other triples of moduli and to classify the conditions under which such systems admit unique solutions modulo their least common multiple. This includes identifying compatibility criteria, determining the structure of the resulting arithmetic progressions, and analyzing how the choice of residues influences the solution class. Such generalizations may reveal broader patterns in the behavior of multi-modulus congruence systems.

8.1.2 Structure of Multi-Modulus Systems

Beyond triples, one may investigate congruence systems involving four or more moduli. The Chinese Remainder Theorem extends naturally to such settings, but the combinatorial and algebraic structure becomes increasingly intricate. Questions of interest include the characterization of solvability conditions, the classification of solution lattices, and the identification of minimal representatives within higher-dimensional congruence frameworks. These investigations may contribute to a deeper understanding of modular arithmetic in complex settings.

8.2 Abstract Topological Extensions

This subsection identifies purely topological generalizations that extend beyond the $U(1)$ case while remaining within an abstract, geometry-free framework. The discussion focuses on higher-rank principal bundles, alternative compact structure groups, and the corresponding integer-valued invariants, emphasizing how these extensions preserve the formal compatibility between cohomological classification and congruence-based arithmetic structure.

8.2.1 Higher-Dimensional Integer Invariants

The classification of $U(1)$ principal bundles over S^2 by integer invariants suggests a broader inquiry into integer-valued topological invariants in higher dimensions. For example, one may study characteristic classes associated with bundles over higher-dimensional spheres or other manifolds, focusing on the algebraic properties of the corresponding cohomology groups. Such investigations remain purely formal and do not require geometric interpretation.

8.2.2 Other Compact Lie Groups

Another direction involves extending the classification framework to principal bundles with structure groups beyond $U(1)$, such as $U(n)$, $SU(2)$, or other compact Lie groups. These groups possess richer cohomological structures, and their associated characteristic classes may exhibit more complex integer or torsion behavior. Exploring these invariants within a purely algebraic and topological context may yield insights into the general theory of bundle classification.

9. Conclusion

This paper has presented a complete and self-contained mathematical analysis of a specific triple-congruence system and the integer lattice that arises from it. Beginning with the formal definition of the congruences, we established their compatibility, applied the Chinese Remainder Theorem to determine the unique solution class modulo 300, and demonstrated that the resulting set of solutions forms the arithmetic progression

$$137 + 300\mathbb{Z}.$$

The minimal positive representative and minimal prime representative were identified through direct number-theoretic arguments, and the structure of the solution set was shown to possess both stability under integer shifts and rigidity with respect to its defining congruence class.

In addition to the arithmetic analysis, the paper provided a purely formal topological interpretation by recalling the classical classification of $U(1)$ principal bundles over S^2 and the integer-valued nature of their first Chern numbers. This interpretation remained strictly abstract and served only to illustrate how integer invariants naturally align with the arithmetic structure of the congruence system, without invoking any geometric or physical context.

All results were derived exclusively from elementary number theory and abstract topology. No external assumptions were introduced, and the analysis remained entirely within the mathematical layer. The framework developed here demonstrates how a simple system of congruences can give rise to a well-defined and stable integer lattice, while also suggesting several avenues for further exploration in both number theory and the study of integer-valued topological invariants.

10. Safety Appendix

This appendix enumerates the categories of content that must not appear anywhere in the paper. These restrictions ensure that the work remains strictly within the domain of pure mathematics and abstract topology, without introducing geometric, physical, engineering, or system-specific interpretations. Each prohibited category is listed with explicit sub-items to prevent accidental overreach.

10.1 Forbidden: Geometric Motivation

This subsection specifies that no geometric reasoning, interpretation, or intuition may be used in any part of the analysis. The restriction excludes all arguments based on spatial representation, curvature, projection, embedding, or any geometric structure, ensuring that the development of the theory remains strictly within its permitted algebraic and topological formal framework.

10.1.1 No Projections

The paper must not reference projection maps, dimensional reductions, or any geometric transformations.

10.1.2 No Embeddings

Embeddings of manifolds, spaces, or structures must not be mentioned in any context.

10.1.3 No Curvature

Curvature, curvature forms, or curvature-related quantities must not appear, even in abstract form.

10.1.4 No Holonomy or Kähler Structure

Holonomy groups, Kähler manifolds, or any related geometric structures must not be referenced.

10.2 Forbidden: Physical Interpretation

This subsection states that no physical meaning, mechanism, or analogy may be assigned to any mathematical object, construction, or invariant appearing in the analysis. All references to measurable quantities, forces, fields, particles, dynamics, or physical models are excluded. The restriction ensures that the results are interpreted strictly within their algebraic and topological domains, without any appeal to physical intuition or phenomenological explanation.

10.2.1 No Coupling Constants

The paper must not mention or allude to any physical coupling constants.

10.2.2 No Fine-Structure Constant

No reference may be made to the fine-structure constant or any physical constant that could be associated with the integer 137.

10.2.3 No Render Layer or Subject Layer

Layered physical or metaphysical models must not be included in any form.

10.3 Forbidden: Engineering Pathways

This subsection establishes that no engineering-level constructions, mechanisms, or implementation details may be introduced at any stage of the analysis. All references to algorithms, architectures, computational procedures, hardware realizations, or operational pipelines are excluded. The restriction ensures that the framework remains strictly abstract and mathematical, without any translation into executable systems or engineering workflows.

10.3.1 No Chip Architecture

The paper must not describe or imply any computational or hardware architecture.

10.3.2 No Computational Units

No mention may be made of processing units, compute modules, or engineering constructs.

10.3.3 No Implementation Details

Algorithms, design pathways, or steps that could be interpreted as implementation guidance must not be included.

10.4 Forbidden: Internal Structural Frameworks

This subsection specifies that no internal structural theories, dimensional hierarchies, operator systems, or generative mechanisms may be invoked in the development of the analysis. All references to proprietary structural models, multi-layer architectures, or internally defined ontological systems are excluded. The restriction ensures that the exposition remains strictly within the externally verifiable mathematical framework and does not rely on any auxiliary structural apparatus.

10.4.1 No Dimensional-Closure Axiom as Physics

The congruence system must not be presented as a physical law, principle, or axiom.

10.4.2 No Structural Cycles

Structural cycles, feedback loops, or system-level dynamics must not be referenced.

10.4.3 No Integrability Conditions

Integrability assumptions, constraints, or conditions must not be introduced, even abstractly.

Mathematical Non-Derivative License 1.0 (MNDL-1.0)

Copyright © 2026 Wujie Gu. All rights reserved.

Permission is granted to any individual or institution to **copy, share, and distribute** this work in its complete and unmodified form for **non-commercial academic purposes**, provided that proper attribution to the author is maintained.

The following restrictions apply:

1. **No Modifications** This work may not be altered, edited, adapted, or used to create derivative works of any kind.
2. **No Interpretive Extensions** This work may not be reinterpreted, reframed, or repurposed in geometric, physical, engineering, computational, or applied contexts. All content must remain within the domain of **pure mathematics**.
3. **No Commercial Use** This work may not be sold, licensed, or incorporated into any commercial product or service.
4. **No Partial Extraction** Individual sections, definitions, or results may not be extracted, rearranged, or reused outside the context of the complete document.
5. **Attribution Requirement** Any permitted redistribution must include the full license text and the following attribution: “© 2026 Gavin. Used with permission under MNDL-1.0.”
6. **No Implied Endorsement** Redistribution may not imply endorsement by the author of any external interpretation, application, or framework.

By accessing or distributing this work, you agree to the terms of the Mathematical Non-Derivative License 1.0.