

PRINT IN CAPITAL LETTERS - STAY WITHIN THE BOX All Fields Required-unless otherwise specified

Last Name			First Name			Middle Initial			Gender			
									☐ Male ☐ Female			
Date of Birth			Weight			Height			Neck Size			
Month	Day	Year	Pounds	Feet	Inches	Inches						
I.D. Number (optional)												

Tally ARES
Risk Points

Neck Size
+2 Male ≥ 16.5
+2 Female ≥ 15

Score

Co-morbidities
+1 for each Yes
response

Score

Do not assign
any points for
these eight
responses

COMPLETELY FILL IN ONE SQUARE FOR EACH QUESTION - ANSWER ALL QUESTIONS

Have you been diagnosed or treated for any of the following conditions?

High blood pressure	Yes ☐	No ☐	Stroke	Yes ☐	No ☐
Heart disease	Yes ☐	No ☐	Depression	Yes ☐	No ☐
Diabetes	Yes ☐	No ☐	Sleep Apnea	Yes ☐	No ☐
Lung disease	Yes ☐	No ☐	Nasal oxygen use	Yes ☐	No ☐
Insomnia	Yes ☐	No ☐	Restless legs syndrome	Yes ☐	No ☐
Narcolepsy	Yes ☐	No ☐	Morning Headaches	Yes ☐	No ☐
Sleep Medication	Yes ☐	No ☐	Pain Medication e.g. vicodin, oxycontin	Yes ☐	No ☐

Epworth Sleepiness Scale: How likely are you to doze off or fall asleep in the following situations, in contrast to just feeling tired? This refers to your usual way of life in recent times. Even if you have not done some of these things recently, try to work out how they would have affected you. Use the following scale to mark the most appropriate box for each situation. (M.W. Johns, Sleep 1991)

0 = would never doze 1 = slight chance of dozing
2 = moderate chance of dozing 3 = high chance of dozing

	0	1	2	3
Sitting and reading	☐	☐	☐	☐
Watching TV	☐	☐	☐	☐
Sitting, inactive, in a public place (theater, meeting, etc)	☐	☐	☐	☐
As a passenger in a car for an hour without a break	☐	☐	☐	☐
Lying down to rest in the afternoon when circumstances permit	☐	☐	☐	☐
Sitting and talking to someone	☐	☐	☐	☐
Sitting quietly after lunch without alcohol	☐	☐	☐	☐
In a car, while stopped for a few minutes in traffic	☐	☐	☐	☐

Epworth Score
Total the
values from all
8 questions.
If 11 or less
Score = 0
If 12 or more
Score = 2

Score

Frequency (Check one for each question): Never +0, Rarely +1 times/wk, Sometimes +2 times/wk, Frequently +3 times/wk, Almost Always +4 times/wk.

On average in the past month, how often have you snored or been told that you snored?

Never +0 ☐ Rarely +1 ☐ Sometimes +2 ☐ Frequently +3 ☐ Almost always +4 ☐

Do you wake up choking or gasping?

Never +0 ☐ Rarely +1 ☐ Sometimes +2 ☐ Frequently +3 ☐ Almost always +4 ☐

Have you been told that you stop breathing in your sleep or wake up choking or gasping?

Never +0 ☐ Rarely +1 ☐ Sometimes +2 ☐ Frequently +3 ☐ Almost always +4 ☐

Do you have problems keeping your legs still at night or need to move them to feel comfortable?

Never ☐ Rarely ☐ Sometimes ☐ Frequently ☐ Almost always ☐

Assign points for
each of the first
three responses

I have personally completed this questionnaire.
Signature

Date

Phone Number

Epworth Sleepiness Scale:

If points total = 3 or lower (no risk)
4 or 5 (low risk), 6 to 10
(high) and 11 or more (very high risk)

Point Total