

Multivariate Outliers w Mahalanobis Distance

2023-12-31

```
###IDENTIFYING MULTIVARIATE OUTLIERS USING MAHALANOBIS DISTANCE###
```

```
#Identifying multivariate outliers in your data can be helpful #if you want to determine whether the correlations between multiple #values or variables are unusually strong for individual cases/customers/participants#####
```

```
###STEP 1###
```

Identify what variables are in linear combination.

This could be, for example, a group of independent

variables used in a multiple linear regression or

a group of dependent variables used in a MANOVA.

Usually, this will include your predictor variable,

any outcome variables, and any mediators or moderators.

```
data <- read_csv("Cues to Infidelity - MEN ONLY 7.12.23.csv")

## Rows: 239 Columns: 554
## -- Column specification -----
## Delimiter: ","
## chr (6): StartDate, EndDate, RecordedDate, M_SC_MRSI_1, M_SC_MRSI_4_2, M_S...
## dbl (462): USE, CUES, SEX_CUES, EMO_CUES, MISC_CUES, REACT, SUS, SOI, VAI_A...
## lgl (86): Q566_1, Q566_2, W_PDIS_A_1, W_PDIS_A_2, W_PDIS_A_3, W_PDIS_A_4, W...
##
## i Use `spec()` to retrieve the full column specification for this data.
## i Specify the column types or set `show_col_types = FALSE` to quiet this message.

###STEP 2###
```

Create a dataframe containing all of the variables you identified in step 1.

```
df <- data.frame(data$REACT, data$SUS, data$SOI, data$SMIRB)

head(df, 10)
```

```
## data.REACT data.SUS data.SOI data.SMIRB
## 1          5.375 1.066667 0.8888889 1.000000
```

```
## 2      4.625 1.133333 2.444444 1.857143
## 3      4.625 1.066667 1.888889 2.428571
## 4      5.125 1.533333 1.777778 1.000000
## 5      5.500 1.666667 1.555556 4.142857
## 6      3.750 1.400000 1.111111 1.000000
## 7      1.750 1.400000 1.000000 1.285714
## 8      5.125 1.600000 2.111111 2.000000
## 9      6.375 2.400000 3.444444 2.428571
## 10     4.875 3.133333 4.555556 3.000000
```

###STEP 3###

Use the mahalanobis() function in R to calculate the distance for each observation:

```
df$mah <- mahalanobis(df, colMeans(df), cov(df))
```

```
head(df, 10)
```

```
##      data.REACT data.SUS  data.SOI data.SMIRB      mah
## 1      5.375 1.066667 0.888889 1.000000 2.4467709
## 2      4.625 1.133333 2.444444 1.857143 1.1614183
## 3      4.625 1.066667 1.888889 2.428571 2.0668241
## 4      5.125 1.533333 1.777778 1.000000 1.5541001
## 5      5.500 1.666667 1.555556 4.142857 8.5213431
## 6      3.750 1.400000 1.111111 1.000000 2.9889809
## 7      1.750 1.400000 1.000000 1.285714 11.0081866
## 8      5.125 1.600000 2.111111 2.000000 0.1157724
## 9      6.375 2.400000 3.444444 2.428571 5.6029036
## 10     4.875 3.133333 4.555556 3.000000 10.9619960
```

###STEP 4###

Calculate p-values for each distance using chi-square

```
df$pvalue <- pchisq(df$mah, df=3, lower.tail = FALSE)
```

```
head(df, 10)
```

```
##      data.REACT data.SUS  data.SOI data.SMIRB      mah      pvalue
## 1      5.375 1.066667 0.888889 1.000000 2.4467709 0.48498762
## 2      4.625 1.133333 2.444444 1.857143 1.1614183 0.76227187
## 3      4.625 1.066667 1.888889 2.428571 2.0668241 0.55865358
## 4      5.125 1.533333 1.777778 1.000000 1.5541001 0.66984151
## 5      5.500 1.666667 1.555556 4.142857 8.5213431 0.03638068
## 6      3.750 1.400000 1.111111 1.000000 2.9889809 0.39332723
## 7      1.750 1.400000 1.000000 1.285714 11.0081866 0.01168169
## 8      5.125 1.600000 2.111111 2.000000 0.1157724 0.98987971
## 9      6.375 2.400000 3.444444 2.428571 5.6029036 0.13261177
## 10     4.875 3.133333 4.555556 3.000000 10.9619960 0.01193316
```

###STEP 5###

Identify cases where $p < .001$ and consider removing these from your data.

```
head(df[order(df$pvalue),], 10)
```

##	data.REACT	data.SUS	data.SOI	data.SMIRB	mah	pvalue
## 43	4.875	1.600000	1.111111	5.285714	20.34492	0.0001439733
## 95	4.500	3.933333	4.555556	3.285714	19.20449	0.0002480300
## 238	1.000	1.066667	1.555556	2.714286	17.49418	0.0005591832
## 16	1.000	1.066667	0.777778	1.857143	16.72444	0.0008052121
## 157	5.375	3.866667	1.888889	4.166667	16.51441	0.0008893148
## 151	1.000	1.200000	1.000000	2.000000	16.15714	0.0010528885
## 210	1.000	1.200000	1.222222	1.000000	15.89539	0.0011913789
## 79	1.000	1.066667	2.111111	1.000000	15.12760	0.0017107888
## 163	5.500	3.800000	2.555556	4.142857	14.77238	0.0020218705
## 204	4.500	3.333333	1.000000	4.142857	14.05326	0.0028335507

```
df_no_multi_outliers <- df[-(which(df$pvalue < .001)),]
```

```
head(df_no_multi_outliers, 10)
```

##	data.REACT	data.SUS	data.SOI	data.SMIRB	mah	pvalue
## 1	5.375	1.066667	0.888889	1.000000	2.4467709	0.48498762
## 2	4.625	1.133333	2.444444	1.857143	1.1614183	0.76227187
## 3	4.625	1.066667	1.888889	2.428571	2.0668241	0.55865358
## 4	5.125	1.533333	1.777778	1.000000	1.5541001	0.66984151
## 5	5.500	1.666667	1.555556	4.142857	8.5213431	0.03638068
## 6	3.750	1.400000	1.111111	1.000000	2.9889809	0.39332723
## 7	1.750	1.400000	1.000000	1.285714	11.0081866	0.01168169
## 8	5.125	1.600000	2.111111	2.000000	0.1157724	0.98987971
## 9	6.375	2.400000	3.444444	2.428571	5.6029036	0.13261177
## 10	4.875	3.133333	4.555556	3.000000	10.9619960	0.01193316