

Table 27-5 Kruskal-Wallis One-Way Analysis of Variance by Ranks: Change in Pain


```
*Nonparametric Tests: Independent Samples.  
NPTESTS  
  /INDEPENDENT TEST (pain) GROUP (group) KRUSKAL_WALLIS (COMPARE=STEPWISE)  
  /MISSING SCOPE=ANALYSIS USERMISSING=EXCLUDE  
  /CRITERIA ALPHA=0.05 CILEVEL=95.
```

Nonparametric Tests

This test was run using Nonparametric Tests > Independent Samples. This method can run multiple comparisons. In SPSS, double clicking on the Hypothesis Test Summary would display further analysis. These results are shown below, including homogeneous subsets.

Hypothesis Test Summary			
	Null Hypothesis	Test	Sig. Decision
1	The distribution of pain is the same across categories of group.	Independent-Samples Kruskal-Wallis Test	.025 Reject the null hypothesis.

Asymptotic significances are displayed. The significance level is .05.



Total N	14
Test Statistic	7.340
Degrees of Freedom	2
Asymptotic Sig. (2-sided test)	.025

1. The test statistic is adjusted for ties.

Homogeneous Subsets based on pain		
	Subset	
	1	2
1	5.400	
Sample ¹ 2	5.800	
3		12.250
Test Statistic	.102	²
Sig. (2-sided test)	.750	.
Adjusted Sig. (2-sided test)	.750	.

Homogeneous subsets are based on asymptotic significances. The significance level is .05.

¹Each cell shows the sample average rank of pain.

²Unable to compute because the subset contains only one sample.

```
NPAR TESTS
  /K-W=pain BY group(1 3)
  /MISSING ANALYSIS.
```

NPar Tests

*This test was run using Nonparametric Tests > Legacy Dialogs > k Independent Samples.
It does not include multiple comparisons.*

Kruskal-Wallis Test

Ranks			
	group	N	Mean Rank
pain	1	5	5.40
	2	5	5.80
	3	4	12.25
	Total	14	

Test Statistics^{a,b}

pain	
Kruskal-Wallis H	7.340
df	2
Asymp. Sig.	.025

- a. Kruskal Wallis Test
- b. Grouping Variable: group