

# CUTANEOUS SQUAMOUS CELL/OTHER CUTANEOUS CARCINOMA STAGING FORM

CLINICAL <i>Extent of disease before any treatment</i>	STAGE CATEGORY DEFINITIONS		PATHOLOGIC <i>Extent of disease through completion of definitive surgery</i>
<input type="checkbox"/> y clinical—staging completed after neoadjuvant therapy but before subsequent surgery	<b>TUMOR SIZE:</b> _____	<b>LATERALITY:</b> <input type="checkbox"/> midline <input type="checkbox"/> left <input type="checkbox"/> right <input type="checkbox"/> bilateral	<input type="checkbox"/> y pathologic – staging completed after neoadjuvant therapy AND subsequent surgery
<input type="checkbox"/> TX <input type="checkbox"/> T0 <input type="checkbox"/> Tis <input type="checkbox"/> T1 <input type="checkbox"/> T2  <input type="checkbox"/> T3 <input type="checkbox"/> T4	<b>PRIMARY TUMOR (T)</b> Primary tumor cannot be assessed No evidence of primary tumor Tis Carcinoma <i>in situ</i> Tumor 2 cm or less in greatest dimension with less than two high risk features** Tumor greater than 2 cm in greatest dimension <i>or</i> Tumor any size with two or more high risk features* Tumor with invasion of maxilla, orbit, or temporal bone Tumor with invasion of skeleton (axial or appendicular) or perineural invasion of skull base * Excludes cSCC of the eyelid – See Chapter 48. **High Risk Features for the Primary Tumor (T) Staging: Depth/Invasion: >2 mm thickness, Clark level ≥ IV, Perineural invasion Anatomic Location: Primary site ear, Primary site hair-bearing lip Differentiation: Poorly differentiated or undifferentiated		<input type="checkbox"/> TX <input type="checkbox"/> T0 <input type="checkbox"/> Tis <input type="checkbox"/> T1 <input type="checkbox"/> T2  <input type="checkbox"/> T3 <input type="checkbox"/> T4
<input type="checkbox"/> NX <input type="checkbox"/> N0 <input type="checkbox"/> N1 <input type="checkbox"/> N2  <input type="checkbox"/> N2a <input type="checkbox"/> N2b  <input type="checkbox"/> N2c  <input type="checkbox"/> N3	<b>REGIONAL LYMPH NODES (N)</b> Regional lymph nodes cannot be assessed No regional lymph node metastasis Metastasis in a single ipsilateral lymph node, 3 cm or less in greatest dimension Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension; or in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension; or in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension Metastasis in a single ipsilateral lymph node, more than 3 cm but not more than 6 cm in greatest dimension Metastasis in multiple ipsilateral lymph nodes, none more than 6 cm in greatest dimension Metastasis in bilateral or contralateral lymph nodes, none more than 6 cm in greatest dimension Metastasis in a lymph node, more than 6 cm in greatest dimension		<input type="checkbox"/> NX <input type="checkbox"/> N0 <input type="checkbox"/> N1 <input type="checkbox"/> N2  <input type="checkbox"/> N2a <input type="checkbox"/> N2b  <input type="checkbox"/> N2c  <input type="checkbox"/> N3
<input type="checkbox"/> M0 <input type="checkbox"/> M1	<b>DISTANT METASTASIS (M)</b> No distant metastasis (no pathologic M0; use clinical M to complete stage group) Distant metastasis		<input type="checkbox"/> M1

HOSPITAL NAME/ADDRESS	PATIENT NAME/INFORMATION

(continued on next page)

# CUTANEOUS SQUAMOUS CELL/OTHER CUTANEOUS CARCINOMA STAGING FORM

## ANATOMIC STAGE • PROGNOSTIC GROUPS

CLINICAL				PATHOLOGIC			
GROUP	T	N	M	GROUP	T	N	M
<input type="checkbox"/> 0	Tis	N0	M0	<input type="checkbox"/> 0	Tis	N0	M0
<input type="checkbox"/> I	T1	N0	M0	<input type="checkbox"/> I	T1	N0	M0
<input type="checkbox"/> II	T2	N0	M0	<input type="checkbox"/> II	T2	N0	M0
<input type="checkbox"/> III	T3	N0	M0	<input type="checkbox"/> III	T3	N0	M0
	T1	N1	M0		T1	N1	M0
	T2	N1	M0		T2	N1	M0
	T3	N1	M0		T3	N1	M0
<input type="checkbox"/> IV	T1	N2	M0	<input type="checkbox"/> IV	T1	N2	M0
	T2	N2	M0		T2	N2	M0
	T3	N2	M0		T3	N2	M0
	T Any	N3	M0		T Any	N3	M0
	T4	N Any	M0		T4	N Any	M0
	T Any	N Any	M1		T Any	N Any	M1

☐ Stage unknown

### PROGNOSTIC FACTORS (SITE-SPECIFIC FACTORS)

#### REQUIRED FOR STAGING:

Tumor thickness (in mm) \_\_\_\_\_  
 Clark's Level \_\_\_\_\_  
 Presence / absence of perineural invasion \_\_\_\_\_  
 Primary site location on ear or hair-bearing lip \_\_\_\_\_  
 Histologic grade \_\_\_\_\_  
 Size of largest lymph node metastasis \_\_\_\_\_

**CLINICALLY SIGNIFICANT:** No additional factors

#### General Notes:

For identification of special cases of TNM or pTNM classifications, the "m" suffix and "y," "r," and "a" prefixes are used. Although they do not affect the stage grouping, they indicate cases needing separate analysis.

**m suffix** indicates the presence of multiple primary tumors in a single site and is recorded in parentheses: pT(m)NM.

**y prefix** indicates those cases in which classification is performed during or following initial multimodality therapy. The cTNM or pTNM category is identified by a "y" prefix. The ycTNM or ypTNM categorizes the extent of tumor actually present at the time of that examination. The "y" categorization is not an estimate of tumor prior to multimodality therapy.

**r prefix** indicates a recurrent tumor when staged after a disease-free interval, and is identified by the "r" prefix: rTNM.

**a prefix** designates the stage determined at autopsy: aTNM.

#### Histologic Grade (G) (also known as overall grade)

##### Grading system

- ☐ 2 grade system  
☐ 3 grade system  
☐ 4 grade system  
☐ No 2, 3, or 4 grade system is available

##### Grade

- ☐ Grade I or 1  
☐ Grade II or 2  
☐ Grade III or 3  
☐ Grade IV or 4

#### ADDITIONAL DESCRIPTORS

**Lymphatic Vessel Invasion (L) and Venous Invasion (V)** have been combined into Lymph-Vascular Invasion (LVI) for collection by cancer registrars. The College of American Pathologists' (CAP) Checklist should be used as the primary source. Other sources may be used in the absence of a Checklist. Priority is given to positive results.

- ☐ Lymph-Vascular Invasion Not Present (absent)/Not Identified  
☐ Lymph-Vascular Invasion Present/Identified  
☐ Not Applicable  
☐ Unknown/Indeterminate

HOSPITAL NAME/ADDRESS

PATIENT NAME/INFORMATION

(continued from previous page)

# CUTANEOUS SQUAMOUS CELL/OTHER CUTANEOUS CARCINOMA STAGING FORM

## Residual Tumor (R)

The absence or presence of residual tumor after treatment. In some cases treated with surgery and/or with neoadjuvant therapy there will be residual tumor at the primary site after treatment because of incomplete resection or local and regional disease that extends beyond the limit of ability of resection.

- ☐ RX Presence of residual tumor cannot be assessed
- ☐ R0 No residual tumor
- ☐ R1 Microscopic residual tumor
- ☐ R2 Macroscopic residual tumor

## General Notes (continued):

**surgical margins** is data field recorded by registrars describing the surgical margins of the resected primary site specimen as determined only by the pathology report.

**neoadjuvant treatment** is radiation therapy or systemic therapy (consisting of chemotherapy, hormone therapy, or immunotherapy) administered prior to a definitive surgical procedure. If the surgical procedure is not performed, the administered therapy no longer meets the definition of neoadjuvant therapy.

☐ Clinical stage was used in treatment planning (describe): \_\_\_\_\_

☐ National guidelines were used in treatment planning ☐ NCCN ☐ Other (describe): \_\_\_\_\_

Physician signature

Date/Time

HOSPITAL NAME/ADDRESS


PATIENT NAME/INFORMATION

(continued on next page)

# CUTANEOUS SQUAMOUS CELL/OTHER CUTANEOUS CARCINOMA STAGING FORM

## Illustration

Indicate on diagram primary tumor and regional nodes involved.



HOSPITAL NAME/ADDRESS

PATIENT NAME/INFORMATION

(continued from previous page)