

# **Machine learning introduction**

**Part I – From neurons to networks**

## **1 – The neuron**

Simon Gay

# Part I-1 : the neuron

- Artificial intelligence aims to mimic several properties of human or animal intelligence
- This intelligence emerges from the most incredible but also mysterious organ : the brain



- Let's analyze it !

# Part I-1 : the neuron

- **Characteristics of human brain:**

- ~80-100 billions neurons
- Equivalent number of *glial cells*
- $\sim 1,5 \times 10^{14}$  connections
- 10.000 to 50.000 neurons and 100 à 500 millions connections by  $\text{mm}^3$




- **Even animals' brains are beyond computer's possibilities:**

- Cat : 760 millions neurons
- Mouse : 71 millions
- frog : 16 millions
- fish : 10 millions
- bee : 960 000 (1 billion connections)
- fly : 250 000 (10 millions connections)


# Part I-1 : the neuron

- Brains are composed of different areas associated to specific functions:

- Cortex :
  - Occipital cortex (vision)
  - Temporal Cortex (hearing and spatial localization)
  - Parietal Cortex (motor coordination and touch)
  - Prefrontal cortex (high level cognitive functions)
- Lymbic system (emotions, navigation, memory, sleeping cycle and hormone control)
- Brain stem (reflexes, body posture, low level vision and sound processing)




The Limbic System




- → The brain is not a neuron tank !

- But all parts rely on neurons



# Part I-1 : the neuron

There are many types (>200) of neurons, with their own working principles





New types are regularly discovered → last one was discovered in 2018 (called rosehip neurons)

# Part I-1 : the neuron

The most common neuron type (and thus the most known) are *multipolar neurons*

- Anatomy of the neuron:
  - A body (also called pericaryon or soma)
  - *Dendrites* (around 7000 per neuron) → inputs
  - A unique *axon* → output
- Neurons communicate through *synapses*
  - Axon to dendrite connection
  - Signal is contactless:
    - The axon emits chemical neurotransmitters
    - The dendrite receives signal through neuroreceptors
  - The strength of the connection depends on the amount of neurotransmitter emitters/receptors in the synapse
- When receiving pulse signals, the neuron tension increase (presence of ions)
  - When reaching -55mV, the neuron emits an electric impulsion of +100mV through its axon, sending a signal to connected neurons
  - Information is encoded in synapses



# Part I-1 : the neuron

## Synaptic learning

- Most common mechanism: Hebb principle (Donald Hebb , 1949)
  - Neurons triggering simultaneously reinforce their synaptic connection
    - When two neurons trigger simultaneously, the number of neurotransmitter emitter/receptor increases to facilitate the signal reception.
    - When two neurons are strongly connected, the signal sent by the first neuron is more likely to trigger the second neuron, that will transmit the signal to next neurons
- Other mechanisms are used by other types of neurons