

Important Notice to Authors

No further publication processing will occur until we receive your response to this proof.

Attached is a PDF proof of your forthcoming article in PRB. Your article has 8 pages and the Accession Code is **BW13806**.

Please note that as part of the production process, APS converts all articles, regardless of their original source, into standardized XML that in turn is used to create the PDF and online versions of the article as well as to populate third-party systems such as Portico, Crossref, and Web of Science. We share our authors' high expectations for the fidelity of the conversion into XML and for the accuracy and appearance of the final, formatted PDF. This process works exceptionally well for the vast majority of articles; however, please check carefully all key elements of your PDF proof, particularly any equations or tables.

Figures submitted electronically as separate files containing color appear in color in the online journal. However, all figures will appear as grayscale images in the print journal unless the color figure charges have been paid in advance, in accordance with our policy for color in print (<https://journals.aps.org/authors/color-figures-print>).

Specific Questions and Comments to Address for This Paper

1 Please update arXiv references with published information, if available.

FQ: This funding provider could not be uniquely identified during our search of the FundRef registry (or no Contract or Grant number was detected). Please check information and amend if incomplete or incorrect.

Q: This reference could not be uniquely identified due to incomplete information or improper format. Please check all information and amend if applicable.

ORCID: Please follow any ORCID links (🔗) after the authors' names and verify that they point to the appropriate record for each author. Requests to add ORCIDs should be sent no later than the first proof revisions. If authors do not subsequently add/authenticate ORCIDs within seven business days, production of the paper will proceed and no further requests to add ORCIDs will be processed. See complete details regarding ORCID requests and ORCID verification at <https://journals.aps.org/authors/adding-orcids-during-proof-corrections>.

NOTE: If this paper is an Erratum or a Reply, the corresponding author's ORCID may be present if previously provided to APS, but no ORCIDs can be added at proof stage.

Crossref Funder Registry ID: Information about an article's funding sources is now submitted to Crossref to help you comply with current or future funding agency mandates. Crossref's Funder Registry (<https://www.crossref.org/services/funder-registry/>) is the definitive registry of funding agencies. Please ensure that your acknowledgments include all sources of funding for your article following any requirements of your funding sources. Where possible, please include grant and award ids. Please carefully check the following funder information we have already extracted from your article and ensure its accuracy and completeness:

Other Items to Check

- Please note that the original manuscript has been converted to XML prior to the creation of the PDF proof, as described above. Please carefully check all key elements of the paper, particularly the equations and tabular data.
- Title: Please check; be mindful that the title may have been changed during the peer-review process.
- Author list: Please make sure all authors are presented, in the appropriate order, and that all names are spelled correctly.
- Please make sure you have inserted a byline footnote containing the email address for the corresponding author, if desired. Please note that this is not inserted automatically by this journal.
- Affiliations: Please check to be sure the institution names are spelled correctly and attributed to the appropriate author(s).
- Receipt date: Please confirm accuracy.
- Acknowledgments: Please be sure to appropriately acknowledge all funding sources.
- Hyphenation: Please note hyphens may have been inserted in word pairs that function as adjectives when they occur before a noun, as in "x-ray diffraction," "4-mm-long gas cell," and "R-matrix theory." However, hyphens are deleted from word pairs when they are not used as adjectives before nouns, as in "emission by x rays," "was 4 mm in length," and "the R matrix is tested."

Note also that Physical Review follows U.S. English guidelines in that hyphens are not used after prefixes or before suffixes: superresolution, quasiequilibrium, nanoprecipitates, resonancelike, clockwise.

- Please check that your figures are accurate and sized properly. Make sure all labeling is sufficiently legible. Figure quality in this proof is representative of the quality to be used in the online journal. To achieve manageable file size for online delivery, some compression and downsampling of figures may have occurred. Fine details may have become somewhat fuzzy, especially in color figures. The print journal uses files of higher resolution and therefore details may be sharper in print. Figures to be published in color online will appear in color on these proofs if viewed on a color monitor or printed on a color printer.
- Please check to ensure that reference titles are given as appropriate.


- Overall, please proofread the entire *formatted* article very carefully. The redlined PDF should be used as a guide to see changes that were made during copyediting. However, note that some changes to math and/or layout may not be indicated.

Ways to Respond

- **Web:** If you accessed this proof online, follow the instructions on the web page to submit corrections.
- **Email:** Send corrections to prbproofs@aptaracorp.com
Subject: **BW13806** proof corrections
- **Fax:** Return this proof with corrections to +1.703.791.1217. Write **Attention:** PRB Project Manager and the Article ID, **BW13806**, on the proof copy unless it is already printed on your proof printout.

Impact ionization processes in a photodriven Mott insulator: Influence of phononic dissipation

Paolo Gazzaneo^{✉,*}, Tommaso Maria Mazzocchi[✉], Jan Lotze[✉], and Enrico Arrigoni[†]
Institute of Theoretical and Computational Physics, Graz University of Technology, 8010 Graz, Austria

 (Received 7 September 2022; revised 7 November 2022; accepted 8 November 2022; published xxxxxxxxx)

We study a model for photovoltaic energy collection consisting of a Mott-insulating layer in presence of acoustic phonons, coupled to two wide-band fermion leads at different chemical potentials and driven into a nonequilibrium steady state by a periodic electric field. We treat electron correlations with nonequilibrium dynamical mean-field theory using the so-called auxiliary master-equation approach as impurity solver and include dissipation by acoustic phonons via the Migdal approximation. For a small hybridization to the leads, we obtain a peak in the photocurrent as a function of the driving frequency which can be associated with impact ionization processes. For larger hybridizations the shallow peak suggests a suppression of impact ionization with respect to direct photovoltaic excitations. Acoustic phonons slightly enhance the photocurrent for small driving frequencies and suppress it at frequencies around the main peak at all considered hybridization strengths.

DOI: [10.1103/PhysRevB.00.005100](https://doi.org/10.1103/PhysRevB.00.005100)

I. INTRODUCTION

The idea of designing photovoltaic devices exploiting the Mott gap to convert electromagnetic radiation into energy has become popular in recent years [1–5]. In particular, it has been suggested that in strongly correlated materials, highly excited charge carriers could use their extra energy to excite additional carriers across the Mott gap via impact ionization (II) [1,4], thus potentially improving their efficiency beyond the Shockley-Queisser limit [6]. Although II is also present in conventional semiconductor devices, the timescales for electron-electron scattering are typically much longer than in correlated materials, so that highly excited electrons will mostly dissipate their energy to phonons.

Experimentally, evidence for fast carrier multiplication processes has been detected via pump-probe experiments in VO₂ [7]. Oxide heterostructures based on LaVO₃/SrTiO₃ have been identified as promising candidates, due to the ideal band gap and the strong polar field, able to separate the excited charge carriers [8]. There are, however, some drawbacks, such as the low mobility of the carriers [5,9], which still put into question these materials' applicability as efficient solar cells. Thus, while a large-scale application of Mott-based solar cells may be difficult to achieve, employing them as photodetectors may be more promising in the long run, due to their high photoresponsivity. From the scientific point of view and for future applications it is thus worthwhile to further investigate the properties of Mott-based photovoltaic materials and the mechanisms behind their inner workings, to explore whether alternative unexpected paths can be opened. In fact, a lot of theoretical work has been done to understand the II process (see, e.g., [4,10–17]).

As a scattering process, II competes with electron-phonon (e-ph) scattering which is the dominant relaxation mechanism in conventional semiconductors. This may not be the case in Mott insulators, at least in certain cases [4]. Therefore, it is essential to understand the influence of e-ph scattering on the photocurrent and II in Mott photovoltaic devices, which is the issue we address in this paper.

For this goal we consider the simplified model shown in Fig. 1, consisting of a Hubbard layer located between two leads. The leads have themselves a wide band, as expected for good conductors, which is effectively narrowed due to the weak coupling to intermediate layers which we take as simplified models for the contacts. Electrons on the Hubbard layer interact locally with acoustic phonons. An external periodic electric field induces a Floquet steady state with a current flowing from the lead with the lower to the lead with the higher chemical potential, thus transferring its energy. We address this periodic problem via nonequilibrium Floquet dynamical mean-field theory (F-DMFT).

Our goal is to investigate the occurrence of II in this photovoltaic setup and how it is affected by the presence of phonons. Specifically, we study the behavior of the photovoltaic current, the spectral properties, as well as the double occupation as a function of the hybridization strength and the properties of the phonons. Our analysis can only provide qualitative results, while quantitative comparisons with experimental systems require a more realistic setup and are beyond the scope of this work.

Our calculations show that a small hybridization to the leads is more favorable for II than a larger one. II therefore plays a crucial role in compensating the smaller charge injection, giving the same order of magnitude in the photocurrent as for the larger hybridization. In all cases, the interaction with phonons slightly enhances the photocurrent for small driving frequencies, while suppressing it in vicinity of the main peak.

The structure of the paper is as follows: In Sec. II we describe the model Hamiltonian. Methods, formalism, and

*paolo.gazzaneo@tugraz.at
 †arrigoni@tugraz.at




FIG. 1. Schematic representation of the setup. A central, correlated layer with local Hubbard interaction U and e-ph coupling g (green) is sandwiched between two noninteracting layers (red) via the hybridization v (black). The latter are in turn coupled to wide-band fermion reservoirs (blue) with different chemical potentials $\mu_{l/r}$, which introduce broadenings $\Gamma_{l/r}$.

observables are presented in Sec. III. We discuss our results in Sec. IV and present our conclusion in Sec. V.

II. MODEL

Our setup introduced in Sec. I and shown in Fig. 1 is described by the Hamiltonian

$$\hat{H}(t) = \varepsilon_c \sum_{i\sigma} \hat{n}_{i\sigma} - \sum_{\sigma} \sum_{(i,j)} t_{ij}(t) \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} + U \sum_i \hat{n}_{i\uparrow} \hat{n}_{i\downarrow} + \hat{H}_{\text{e-ph}} + \hat{H}_{\text{ph}} + \hat{H}_{\text{lead}}. \quad (1)$$

The operator $\hat{c}_{i\sigma}^{\dagger}$ ($\hat{c}_{i\sigma}$) creates (annihilates) an electron with spin $\sigma = \{\uparrow, \downarrow\}$ on the i th lattice site and $\hat{n}_{i\sigma} \equiv \hat{c}_{i\sigma}^{\dagger} \hat{c}_{i\sigma}$ is the particle-number operator. We denote with (i, j) the sum over nearest-neighbor sites and with $\varepsilon_c \equiv -U/2$ the onsite energy. The driving field consists of a time-periodic, homogeneous, and monochromatic electric field with frequency Ω . In the temporal gauge it enters via the Peierls substitution in the time-dependent hopping in Eq. (1) [18]

$$t_{ij}(t) = t_c e^{-i \frac{q}{\hbar} (\mathbf{r}_j - \mathbf{r}_i) \cdot \mathbf{A}(t)}, \quad (2)$$

where t_c is the intralayer hopping, $\mathbf{A}(t)$ the time-dependent vector potential, \hbar Planck's constant, and q the charge of the electrons. Following Refs. [19,20], we choose for simplicity $\mathbf{A}(t) = \mathbf{e}_0 A(t)$, which points along the lattice body diagonal $\mathbf{e}_0 = (1, 1, \dots, 1)$. Therein $A(t) = \frac{\hbar}{qa} \mathcal{A} \sin(\Omega t)$ with $\mathcal{A} = -\frac{qE_0 a}{\hbar \Omega}$, E_0 is the electric field amplitude, and a the lat-

tice spacing [19]. In this temporal gauge the electric field is $\mathbf{E} = -\partial_t \mathbf{A}(t) = \mathbf{e}_0 E_0 \cos(\Omega t)$.

As in Ref. [21], phonons are included in the form of acoustic branches attached to each lattice site and coupled to the electrons via the Hamiltonian

$$\hat{H}_{\text{e-ph}} = g \sum_{i\sigma} \hat{n}_{i\sigma} \hat{x}_i, \quad (3)$$

with g as e-ph coupling and $\hat{x}_i \equiv \frac{1}{\sqrt{2}} (\hat{b}_i^{\dagger} + \hat{b}_i)$, where \hat{b}_i^{\dagger} (\hat{b}_i) creates (annihilates) a phonon of the acoustic branch at site i , with dispersion relation described by \hat{H}_{ph} , discussed in Sec. III B. In contrast to Holstein phonons, they have a finite bandwidth, which is more effective in energy dissipation. Details on the leads are provided in Sec. III B.

For mathematical simplicity [22,23], we consider the correlated central region as a d -dimensional layer and consider the limit of infinite dimensions $d \rightarrow \infty$ by rescaling the hopping as $t_c = t^*/(2\sqrt{d})$. Taking $a = 1$, sums over the crystal momentum \mathbf{k} of a generic quantity χ read as $\sum_{\mathbf{k}} \chi(\omega, \mathbf{k}) \rightarrow \int d\epsilon \int d\bar{\epsilon} \rho(\epsilon, \bar{\epsilon}) \chi(\omega; \epsilon, \bar{\epsilon})$ with $\rho(\epsilon, \bar{\epsilon}) = (1/\pi t^{*2}) \exp[-(\epsilon^2 + \bar{\epsilon}^2)/t^{*2}]$ as the joint density of states [19] and with

$$\begin{aligned} \epsilon &= -2t_c \sum_{i=1}^d \cos(k_i a), \\ \bar{\epsilon} &= -2t_c \sum_{i=1}^d \sin(k_i a). \end{aligned} \quad (4)$$

Within the remainder of this paper we choose our units so that $\hbar = k_B = a = 1 = -q$, $t^* = 2\sqrt{2}$,¹ and $t_{\text{resc}} \equiv \frac{t^*}{2\sqrt{2}} = 1$ as unit of energy.

III. METHOD AND FORMALISM

A. Floquet Green's function method

To correctly describe the nonequilibrium periodic steady state, we use the Floquet generalization of the nonequilibrium Green's function (NEGF) approach [19,24,25]. Every function $G(t, t')$ satisfying the periodicity relation $G(t, t') = G(t + \tau, t' + \tau)$, with $\tau = 2\pi/\Omega$ as the period related to the external driving frequency Ω , may be represented as [19,24,25]

$$\underline{G}_{mn}(\omega) = \int dt_{\text{rel}} \int_{-\tau/2}^{\tau/2} \frac{dt_{\text{av}}}{\tau} e^{i[(\omega + m\Omega)t - (\omega + n\Omega)t']} \underline{G}(t, t'), \quad (5)$$

known as the *Keldysh-Floquet* GF. The integration variables $t_{\text{rel}} = t - t'$ and $t_{\text{av}} = (t + t')/2$ are the relative and average times. Within the rest of this work, we denote a Floquet-represented matrix either as X_{mn} with explicit indices, or use a boldface letter \underline{X} . The underline indicates the overall *Keldysh* structure

$$\underline{G} \equiv \begin{pmatrix} G^R & G^K \\ \mathbf{0} & G^A \end{pmatrix}, \quad (6)$$

¹With such choice for t^* we reproduce the DOS of a 2D layer with $t_c = 1$ in order to compare with Ref. [14].

which contains the *retarded*, *advanced*, and *Keldysh* components $G^{\text{R,A,K}}$, where $G^{\text{A}} = (G^{\text{R}})^{\dagger}$. The *Keldysh* component is defined as $G^{\text{K}} \equiv G^{\text{>}} + G^{\text{<}}$, with $G^{\text{>}}$ as *lesser* and *greater* components [26–29].

B. Dyson equation

The lattice Floquet GF of our setup obeys the Dyson equation

$$\underline{G}^{-1}(\omega_n; \epsilon, \bar{\epsilon}) = \underline{G}_0^{-1}(\omega_n; \epsilon, \bar{\epsilon}) - \underline{\Sigma}(\omega_n; \epsilon, \bar{\epsilon}) - \underline{\Sigma}_{\text{e-ph}}(\omega_n; \epsilon, \bar{\epsilon}), \quad (7)$$

where in our approximation the electron self-energy (SE) $\underline{\Sigma}$ and e-ph SE $\underline{\Sigma}_{\text{e-ph}}$ contribute independently to the total SE.² Therein, the lattice GF of the noninteracting part of the Hamiltonian in Eq. (1) is

$$\underline{G}_{0,mn}^{-1}(\omega_n; \epsilon, \bar{\epsilon}) = \underline{g}_{0,mn}^{-1}(\omega_n; \epsilon, \bar{\epsilon}) - \sum_{\rho \in \{l,r\}} v_{\rho}^2 \underline{g}_{\text{b},\rho}(\omega_n; \epsilon) \delta_{mn} \quad (8)$$

with $v_{l/r}$ as lead-layer hybridization, $\omega_n \equiv \omega + n\Omega$, $n \in \mathbb{Z}$, and

$$\begin{aligned} [g_0^{-1}(\omega_n; \epsilon, \bar{\epsilon})]_{mn}^{\text{R}} &= (\omega_n + i0^+ - \epsilon_c) \delta_{mn} - \epsilon_{mn}(\epsilon, \bar{\epsilon}), \\ [g_0^{-1}(\omega_n; \epsilon, \bar{\epsilon})]_{mn}^{\text{K}} &= 0 \end{aligned} \quad (9)$$

as the noninteracting Green's function of the isolated layer, whose inverse Keldysh component is negligible in the steady state. The Floquet dispersion relation ϵ_{mn} for the periodic field in a hypercubic lattice is [19]

$$\epsilon_{mn}(\epsilon, \bar{\epsilon}) = \begin{cases} \epsilon J_{m-n}(\mathcal{A}) & m - n : \text{even}, \\ i\bar{\epsilon} J_{m-n}(\mathcal{A}) & m - n : \text{odd}, \end{cases} \quad (10)$$

where J_n denotes the n th-order Bessel function of the first kind, with the argument \mathcal{A} defined in Sec. II. The GF $\underline{g}_{\text{b},l/r}$ on the d -dimensional boundary of the decoupled leads, consisting of a single layer coupled to a semi-infinite reservoir in the wide-band limit with broadening $\Gamma_{l/r}$ (cf. Fig. 1), is given by

$$g_{\text{b},l/r}^{\text{R}}(\omega; \epsilon) = \frac{1}{\omega - \epsilon_{l/r}(\epsilon) + i\Gamma_{l/r}}, \quad (11)$$

$$g_{\text{b},l/r}^{\text{K}}(\omega; \epsilon) = 2i \text{Im}[g_{\text{b},l/r}^{\text{R}}(\omega; \epsilon)][1 - 2f(\omega, \mu_{l/r}, \beta)], \quad (12)$$

where $\epsilon_{l/r}(\epsilon) = \epsilon_{l/r} + \frac{\hbar_{l/r}}{\tau} \epsilon$ is the dispersion, $f(\omega, \mu_{l/r}, \beta) = 1/[e^{\beta(\omega - \mu_{l/r})} + 1]$ the Fermi-Dirac distribution function at inverse temperature $\beta \equiv 1/T$, $\epsilon_{l/r}$ the onsite energy, and $\hbar_{l/r}$ the hopping within the d -dimensional layer of the lead.

The electron SE $\underline{\Sigma}$ is obtained from F-DMFT and, therefore, in this approximation, independent of the crystal momentum, i.e., $\underline{\Sigma}(\omega; \epsilon, \bar{\epsilon}) \simeq \underline{\Sigma}(\omega)$. Further details are given in Sec. III C. In the same spirit, the e-ph SE is included locally as $\underline{\Sigma}_{\text{e-ph}}(\omega; \epsilon, \bar{\epsilon}) \simeq \underline{\Sigma}_{\text{e-ph}}(\omega)$. In terms of the Keldysh contour time arguments z and z' , it has the form

$$\Sigma_{\text{e-ph}}(z, z') = ig^2 G(z, z') D_{\text{ph}}(z, z'). \quad (13)$$

The Keldysh components of the noninteracting phonon GF $\underline{D}_{\text{ph}}(t, t')$ are given by [23]

$$\begin{aligned} D_{\text{ph}}^{\text{R}}(t, t') &= -i\theta(t - t') \int d\omega e^{-i\omega(t-t')} A_{\text{ph}}(\omega), \\ D_{\text{ph}}^{\text{>}}(t, t') &= -i \int d\omega e^{-i\omega(t-t')} A_{\text{ph}}(\omega) [1 + b(\omega)], \\ D_{\text{ph}}^{\text{<}}(t, t') &= -i \int d\omega e^{-i\omega(t-t')} A_{\text{ph}}(\omega) b(\omega), \end{aligned} \quad (14)$$

where $b(\omega) = 1/(e^{\beta\omega} - 1)$ is the Bose-Einstein distribution function at inverse temperature β . We consider acoustic phonons, with spectral function $A_{\text{ph}}(\omega) = (\omega/\omega_{\text{ph}}^2) e^{-|\omega|/\omega_{\text{ph}}}$, ω_{ph} being a soft cutoff frequency [30]. The retarded and Keldysh components of the e-ph SE are easily extracted from Eq. (13) and can be found in Ref. [21].

C. Floquet DMFT

We compute the electron SE in the Dyson equation (7) using DMFT [22,31,32], and in particular its nonequilibrium Floquet extension F-DMFT [19,24,25]. In DMFT the crystal momentum dependence of the electron SE is neglected, i.e., $\underline{\Sigma}(\omega, \epsilon, \bar{\epsilon}) \rightarrow \underline{\Sigma}(\omega)$. This allows us to map the original lattice problem onto a single-site impurity model with a bath hybridization function $\underline{\Delta}(\omega)$ encoding the effect of all other lattice sites.

For completeness, we now briefly describe the self-consistency F-DMFT scheme used. (i) We start from an initial guess for the electron SE $\underline{\Sigma}(\omega)$ and set $\underline{\Sigma}_{\text{e-ph}}(\omega) = 0$. (ii) Then we compute the local electron GF as

$$\begin{aligned} \underline{G}_{\text{loc}}(\omega) &= \int d\epsilon \int d\bar{\epsilon} \rho(\epsilon, \bar{\epsilon}) \\ &\times [\underline{G}_0^{-1}(\omega, \epsilon, \bar{\epsilon}) - \underline{\Sigma}(\omega) - \underline{\Sigma}_{\text{e-ph}}(\omega)]^{-1}. \end{aligned} \quad (15)$$

(iii) Using Eq. (13), we obtain the phonon contribution to the e-ph SE $\underline{\Sigma}_{\text{e-ph}}(\omega)$. (iv) The problem is mapped onto a single impurity plus bath, whose hybridization function is given by

$$\underline{\Delta}(\omega) = \underline{g}_{0,\text{site}}^{-1}(\omega) - \underline{G}_{\text{loc}}^{-1}(\omega) - \underline{\Sigma}(\omega), \quad (16)$$

where $\underline{g}_{0,\text{site}}^{-1}(\omega)$ is defined as in Eq. (9) with $\epsilon_{mn}(\epsilon, \bar{\epsilon}) = 0$. (v) The nonequilibrium many-body impurity problem is solved according to the procedure described below, leading to the new $\underline{\Sigma}(\omega)$. (vi) We insert the electron and e-ph SEs into step (ii) and iterate the steps (ii)–(vi) until convergence.

Equation (16) gives a bath hybridization function $\underline{\Delta}(\omega)$ whose periodic time dependence is encoded in its Floquet structure. This means that, in principle, one should solve a time-periodic impurity problem leading to a nondiagonal SE. However, in Ref. [14] it was argued and shown that for the considered electric field amplitudes, off-diagonal terms in the electron SE $\underline{\Sigma}(\omega)$ can be safely neglected with respect to the diagonal ones. We follow this argument and adopt here this Floquet-diagonal self-energy approximation (FDSA) as well. In addition, it turns out that also for the e-ph SE $\underline{\Sigma}_{\text{e-ph}}(\omega)$

²In fact, there is an indirect feedback via the DMFT self-consistency.

this approximation is justified.³ Consequently, we solve a nonequilibrium stationary impurity problem, by considering only the (0,0) Floquet matrix element of all the quantities in Eq. (16). The other diagonal components of the SEs are then reconstructed by using the property $\Sigma_{mm}(\omega) = \Sigma_{00}(\omega + m\Omega)$.

In order to solve the many-body problem in step (v) of the DMFT self-consistent loop, we use the auxiliary master-equation approach (AMEA) [33–36], which we now briefly summarize. Therein, we map the impurity problem onto an auxiliary open quantum system (AOQS) consisting of a finite number of bath sites N_B attached to Markovian reservoirs described by the Lindblad equation. The hybridization function $\Delta_{\text{aux}}(\omega)$ of this AOQS is obtained by fitting the original DMFT one. The key point is that the many-body problem of this AOQS can be solved exactly using standard many-body-diagonalization methods, as long as N_B is small. The accuracy of this solution is set by the difference between Δ_{aux} and Δ , which becomes exponentially small with increasing N_B .

D. Physical quantities

To study direct excitation and II effects in this system, we consider the following time-averaged physical quantities. The photocurrent flowing from the left fermion lead to the right one, passing through the correlated layer, is given by two equivalent expressions, adapted from Ref. [14]:

$$j_{l \rightarrow r} = v^2 \int_{-\Omega/2}^{\Omega/2} \frac{d\omega}{2\pi} \int d\epsilon \int d\bar{\epsilon} \rho(\epsilon, \bar{\epsilon}) \text{ReTr}(\mathbf{J}) \quad (17)$$

$$= v^2 \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi} \int d\epsilon \int d\bar{\epsilon} \rho(\epsilon, \bar{\epsilon}) \text{Re}(J_{00}), \quad (18)$$

where

$$\mathbf{J} = [\mathbf{G}^R(\mathbf{g}_{b,l}^K - \mathbf{g}_{b,r}^K) + \mathbf{G}^K(\mathbf{g}_{b,l}^A - \mathbf{g}_{b,r}^A)]. \quad (19)$$

In our case, it is convenient to use Eq. (18).

The local electronic spectral function (DOS) reads as

$$A(\omega) = -\frac{1}{\pi} \text{Im}[G_{\text{loc},00}^R(\omega)], \quad (20)$$

where $G_{\text{loc},00}^R$ is the time-averaged retarded component of the GF given in Eq. (15). Combining it with the time-averaged Keldysh component gives the occupation function

$$N(\omega) = \frac{1}{4\pi} [\text{Im}(G_{\text{loc},00}^K(\omega)) - 2\text{Im}(G_{\text{loc},00}^R(\omega))]. \quad (21)$$

IV. RESULTS

In order to study II in a Mott-insulating layer, we adjust the bands to achieve the energy scheme shown in Fig. 2(a). We follow Ref. [14] and take $U = 12$ and unless stated otherwise $E_0 = 2$. The fermion leads and the acoustic phonons have temperature $T = 0.02$.⁴ We consider a particle-hole symmet-




FIG. 2. Schematic representation of the processes occurring in the model considered. Red bands on the left and right describe the leads' DOS, where the dark color highlights occupied states and the light color empty states. Green bands are the LHB and UHB of the central layer. The vertical axis represents the energy. Due to the hybridization with the metallic leads only a pseudogap Δ_{pg} is present between the Hubbard bands. (a) Sketches the quantities introduced at the beginning of Sec. IV. (b) Illustrates a direct excitation process, in which an electron is excited by a photon with energy Ω (yellow arrow) to the UHB and escapes into the right lead (red arrow). (c) Displays an II process, where the photoexcited electron in the UHB excites a second electron from LHB to UHB (black arrows) and both escape into the right lead. In the nonequilibrium steady state considered here, only processes recovering the initial configuration are allowed.

ric system with $\Gamma_l = \Gamma_r = 0.37$, $t_l = t_r = 1.7$, $\varepsilon_{l/r} = \mp 6$, and $v_l = v_r = v$. The parameters $t_{l/r}$, $\varepsilon_{l/r}$, and $\Gamma_{l/r}$ are chosen such that the leads' DOS approximately overlaps with the Hubbard bands, i.e., they exhibit a “bandwidth”⁵ $W_b \approx 8$ and are centered at the same position. The chemical potentials are set to

³More specifically, we verified that in the parameter range in which FDSA is justified for the electron SE $\Sigma(\omega)$ (Sec. IV), the off-diagonal terms of the e-ph SE $\Sigma_{\text{e-ph}}(\omega)$ are sufficiently suppressed.

⁴The value chosen for the temperature is much smaller than the other characteristic energy scales of the system, e.g., ω_{ph} , U , and W_b .

⁵Notice that the support of the leads' spectra are formally infinite. W_b is the value at which the spectrum gets suppressed.

$\mu_{l/r} = \mp 1$, so that a current from left to right is produced by taking energy from the driving. Due to the hybridization with the leads, the local DOS of the Hubbard layer only features a pseudogap $\Delta_{pg} \approx 4$. All simulations are carried out with the dimensionless factor $\alpha \equiv t_{\text{resc}} E_0 / \Omega^2 < 0.5$ for which the FDSA is justified.⁶ Whenever the electron-phonon interaction is included, we take $g = 0.8$ and unless stated otherwise $\omega_{ph} = 0.1$.

After some preliminary considerations in Sec. IV A, we discuss the occurrence of II at different v first without (Sec. IV B) and then with coupling to acoustic phonons (Sec. IV C).

A. Qualitative energy considerations and physical processes

To infer the conditions necessary for II, we consider the scheme in Fig. 2. Thereby, we partially follow Ref. [14] for the electron-only (EO) case and extend the analysis to the case with e-ph interaction. In comparison with Ref. [14], here the wide-band leads describe more realistically a metallic structure. This makes the identification of II processes more subtle, as discussed below. As in Ref. [14] we bypass transient behavior and consider directly the Floquet steady state. In our analysis, we neglect higher-order processes such as multiparticle scattering processes.

In order to observe II, the bandwidth of the upper Hubbard band (UHB) has to be at least twice the pseudogap⁷ size Δ_{pg} . Only in this case, the photoexcited electron can acquire enough energy to excite a second one across the pseudogap.

1. EO system

(i) For $\Omega < \Delta_{pg}$, we expect a strong suppression of the current⁸ and we do not expect any current if the DOS of the correlated layer has a true gap.

(ii) For $\Delta_{pg} < \Omega < \Delta_{pg} + 2W_b$ as shown in Fig. 2(b), an electron coming from the left lead into the lower Hubbard band (LHB) is photoexcited to the UHB and can escape directly into the right lead without additional scattering. Such processes are often referred to as direct excitations (DEs).

(iii) For $2\Delta_{pg} < \Omega < \Delta_{pg} + 2W_b$ as shown in Fig. 2(c), a photoexcited electron in the UHB can excite via II a second electron from LHB to UHB, before both escape into the right lead.

(iv) For $\Omega > \Delta_{pg} + 2W_b$, there are no final states available for a photoexcited electron and we do not expect the transition to occur.

Notice that in the energy window $\Delta_{pg} < \Omega < 2\Delta_{pg}$ the only scattering processes taking place are DEs, while for $2\Delta_{pg} < \Omega < \Delta_{pg} + 2W_b$, both DE and II can occur. This is in contrast to the case of Ref. [14], in which the *ad hoc* narrow

⁶In Ref. [14] the factor is defined as $\alpha = t_c E_0 / \Omega^2$ and $t_c = 1$. Setting $t_{\text{resc}} = 1$, we define for consistency α as in the main text.

⁷Due to the hybridization with the metallic leads, only a pseudogap Δ_{pg} is present between the LHB and the UHB.

⁸The remaining leak current originates either from multiple-photon absorption processes which are strongly suppressed, or from the absence of a true gap.




FIG. 3. (a) Time-averaged steady-state current j and (b) double occupancy N_D plotted as a function of the driving frequency Ω . Results are shown for different values of the lead-layer hybridization v as well as without and with e-ph interaction. Default parameters are specified at the beginning of Sec. IV.

leads' bands only allow for II processes in a certain Ω range. For this reason, and due to the fact that the boundaries of these energy ranges are not strict, it is more difficult to disentangle these physical processes in the present case.

2. Inclusion of e-ph interactions

Upon inclusion of e-ph scattering, the outlined scheme remains valid except for the fact that the pseudogap Δ_{pg} is slightly modified. As shown in Sec. IV C, the phonons broaden the DOS of the correlated layer, which slightly fills and shrinks the pseudogap.

B. EO system

We start by analyzing the case without e-ph interaction and address the behavior of the physical quantities as functions of Ω for different values of the lead-layer hybridization.

In Fig. 3(a), the photocurrent j increases as a function of the driving frequency Ω within the range $4 \lesssim \Omega \lesssim 8$. According to the discussion in Sec. IV A 1 (cf. Fig. 2), this Ω range is expected to allow only for DEs. The current in this range increases with increasing v . Similarly, the double occupation shown in Fig. 3(b) increases as a function of the driving frequency.⁹

For $\Omega \gtrsim 8$, the current increases further until it reaches its peak at $\Omega \approx 11$. In this range, both DE and II can occur in principle. The height of the peak is approximately the same for both v , despite the hybridizations differ by a factor of 2. The double occupation in this range shows a similar behavior as the current for $v = 0.4$ with a peak around $\Omega \approx 11$, while it exhibits a plateau for $v = 0.8$. For $v = 0.4$ the current curve is characterized by a substantial increase in slope around $\Omega \approx 7$, which cannot be observed for $v = 0.8$. This, together with the comparable magnitude of the current maxima at $\Omega \approx 11$ and the behavior of the double occupation strongly suggests that for $v = 0.4$ a substantial amount of II processes take place, while these are absent or negligible for $v = 0.8$. Here, the

⁹Notice that the increase of N_D in Fig. 3(b) as compared to Ref. [14] is not as steep and large in magnitude. This is due to our setup, where every photoexcited carrier can escape into the right lead.




FIG. 4. Spectral function $A(\omega)$ (solid line) and occupation function $N(\omega)$ (shaded area) at (a) $\Omega = 5$ and (b) $\Omega = 11$ for $v = 0.4$ and 0.8 for the EO system. Default parameters are specified at the beginning of Sec. IV.

plateau in N_D for $v = 0.8$ suggests that photoexcited electrons arriving in the UHB quickly escape to the right fermionic lead and do not have time to induce Π processes.

The spectral and occupation functions shown in Fig. 4 corroborate this hypothesis. For $\Omega = 5$, in the DE range, the occupation function in the UHB is almost the same for both v , while for $\Omega = 11$ where the peak in the current occurs, the occupation of the UHB for $v = 0.4$ is substantially larger, in line with the behavior of N_D . Further increasing Ω produces excitations near the border of the UHB where the DOS is reduced and thus current and double occupation decrease.¹⁰

This behavior is affected by the electric field amplitude E_0 as illustrated in Fig. 5(a). For $E_0 = 4$ the current develops a sudden increase in slope around $\Omega \approx 7$, as well as a pronounced peak around $\Omega \approx 11$ for the larger hybridization $v = 0.8$. Both features start developing already at $E_0 = 3$. This tendency, corroborated by the behavior of the double occupation displayed in Fig. 5(b), suggests an important role of Π also for larger values of v , provided the field amplitude is strong enough.

¹⁰The current j does not approach zero as expected for $\Omega > 20$ because of the background current as discussed in Ref. [14].




FIG. 5. (a) Time-averaged steady-state current j and (b) double occupancy N_D as a function of Ω , at $v = 0.8$ for $E_0 = 3, 4$ for the EO system. Default parameters are specified at the beginning of Sec. IV.




FIG. 6. Spectral function $A(\omega)$ (solid line) and occupation function $N(\omega)$ (shaded area) at $\Omega = 11$ without and with e-ph interaction, for (a) $v = 0.4$ and (b) $v = 0.8$. Default parameters are specified at the beginning of Sec. IV.

C. Inclusion of e-ph interactions

We now discuss the effect of coupling to acoustic phonons on the results presented thus far. The current j in presence of phonons shown in Fig. 3(a) is slightly larger than the EO one for $\Omega \lesssim 7$. This effect is reduced with increasing Ω until a crossing occurs at $\Omega \approx 7$ where the EO current overtakes. Between $7 \lesssim \Omega \lesssim 14$, the current in the presence of phonons exhibits the same qualitative behavior as the EO one, reaching its maximum at $\Omega \approx 11$. However, the current magnitude is suppressed, especially around the peak. For $\Omega \gtrsim 14$ the current in presence of phonons drops as for the EO case.

The double occupation N_D shown in Fig. 3(b) is larger than and follows the behavior of its EO counterpart within the entire Ω range considered. The spectral and occupation functions shown in Fig. 6 display a redistribution of spectral weight from the peaks towards the edges of the bands, thus reducing the pseudogap. Consequently, the occupation of positive-energy states is slightly shifted to the bottom of the UHB compared to the EO case, as evidenced by the occupation function. The spectral function in presence of phonons is broadened and tends to fill the pseudogap Δ_{pg} as anticipated in Sec. IV A 2. This reduction of the pseudogap explains why the current in Fig. 3(a) is slightly larger in presence of phonons for $\Omega \lesssim 7$. With a smaller pseudogap and more states at its edges, more electrons can be directly excited with a given driving frequency Ω thereby increasing the photocurrent. On the other hand, the suppression of the current by e-ph interaction in the range $7 \lesssim \Omega \lesssim 14$ is due to the dissipation-induced reduction of spectral weight at frequencies $|\omega| \gtrsim 3.5$ (see Fig. 6).

The increase of the double occupation N_D seen in Fig. 3(b) for the case with phonons can already be inferred from the occupation function $N(\omega)$ at positive frequencies depicted in Fig. 6.

Summarizing, phonons slightly enhance the current for almost all driving frequencies $\Omega \lesssim 7$ for which only DEs take place. They suppress the current in the high-frequency range $7 \lesssim \Omega \lesssim 14$. This behavior is qualitatively valid for different values of the soft cutoff phonon frequency ω_{ph} . It can be seen in Fig. 7, which shows that the impact of acoustic phonons




FIG. 7. (a) Time-averaged steady-state current j and (b) double occupancy N_D as a function of Ω , for $v = 0.4, 0.8$, at phonon cutoff frequencies $\omega_{ph} = 0.025, 0.05$. Default parameters are specified at the beginning of Sec. IV.

on current and double occupancy is slightly boosted when decreasing ω_{ph} .¹¹

The results above suggest that the influence of acoustic phonons on the electronic scattering processes does not depend significantly on v . In other words, the effect of phonons does not depend significantly on the rate at which carriers are injected into and removed from the layer. Figure 8 confirms this behavior for intermediate lead-layer hybridizations $v = 0.5, 0.6, 0.7$. Also, the location of the crossing in the current between the EO and e-ph cases is essentially independent of v , as can be seen in Fig. 8(a).

V. CONCLUSION

We investigated the influence of fermion leads and phonon dissipation on electron transport and spectral properties of a Mott-insulating layer driven to the nonequilibrium steady

¹¹Reducing ω_{ph} has two effects. First, the phonon spectral function $A_{ph}(\omega)$ exhibits more weight at low frequencies around $\omega \approx \omega_{ph}$. Second, it restricts the maximum value of the reciprocal lattice vector \mathbf{q}_{max} via $\omega_{ph} = \omega(\mathbf{q}_{max})$ [21], meaning that short-wavelength phonons are suppressed.




FIG. 8. (a) Time-averaged steady-state current j and (b) double occupancy N_D as a function of Ω , for $v = 0.5, 0.6, 0.7$, without and with e-ph interaction. Default parameters are specified at the beginning of Sec. IV.

state by an external periodic electric field. In order to realize a Mott-based photovoltaic setup between metallic leads, we considered a correlated layer coupled to acoustic phonons, located between wide-band fermion leads. We studied the influence of the strength of the coupling to the leads on the scattering process occurring at different driving frequencies and observed how the dissipation by acoustic phonons influences the photocurrent and the double occupancy. We found evidence of a significant amount of II processes leading to a photocurrent peak at small hybridizations, while these are suppressed for larger hybridizations. Dissipation via acoustic phonons slightly boosts the photocurrent for small driving frequencies and suppresses it at larger ones in the vicinity of the main peak. The effect of phonons is not affected significantly by changes in the lead-layer hybridization.

One should comment on the validity of DMFT, which should be accurate for large dimensions only, for the present quasi-two-dimensional (2D) setup. As a matter of fact, the hybridization to the leads introduces a dissipation which suppresses 2D coherence and thus makes nonlocal correlations beyond DMFT less relevant. On the other hand, it has been shown that antiferromagnetic (even short-range) correlations may play an important role in the spreading of photoexcited carriers [37]. Taking into account the effect of such nonlocal correlations (see also [38]) might be an interesting extension, although beyond the scope of the present research.

Experimentally, the electric field amplitude considered in this paper is several orders of magnitude larger than that of the sunlight, therefore, our results are more relevant for photoexcitations produced by intense laser pulses (see also the discussion in [20]) rather than for true photovoltaic systems. We did not consider smaller electric field amplitudes due to the slow F-DMFT convergence for such a choice of parameters. This is also the case for weaker hybridization strengths.

On the other hand, one may try to qualitatively extrapolate the discussion at the end of Sec. IV B in the other direction to smaller electric field amplitudes, for which a smaller hybridization would be necessary in order to sustain II. In a more realistic setup modeling oxide heterostructures [8,13], a regime of effectively small hybridization would be achieved by considering several layers in which photoexcited carriers, separated by an electric field gradient, have more time to induce II processes before escaping into the leads. To address the occurrence of II in such a setup, further extensions such as considering multiple orbitals and impurity scattering should be taken into account. This aspect could be interesting for future studies. Finally, a more realistic description of the effects of the electron-phonon interaction on electronic band structure should take into account the crystal momentum dependence of Σ_{e-ph} [39].

ACKNOWLEDGMENTS

We thank A. Picano and C. Heil for fruitful discussions. This work was supported by the Austrian Science Fund (Grant No. P 33165-N) and by NaWi Graz. The computational results presented have been obtained using the Vienna Scientific Cluster (VSC) and the D-Cluster Graz.

- [1] E. Manousakis, *Phys. Rev. B* **82**, 125109 (2010).
- [2] H. Liang, L. Cheng, X. Zhai, N. Pan, H. Guo, J. Zhao, H. Zhang, L. Li, X. Zhang, X. Wang, C. Zeng, Z. Zhang, and J. G. Hou, *Sci. Rep.* **3**, 1 (2013).
- [3] H.-Z. Guo, L. Gu, Z.-Z. Yang, S.-F. Wang, G.-S. Fu, L. Wang, K.-J. Jin, H.-B. Lu, C. Wang, C. Ge, M. He, and G.-Z. Yang, *Europhys. Lett.* **103**, 47006 (2013).
- [4] J. E. Coulter, E. Manousakis, and A. Gali, *Phys. Rev. B* **90**, 165142 (2014).
- [5] L. Wang, Y. Li, A. Bera, C. Ma, F. Jin, K. Yuan, W. Yin, A. David, W. Chen, W. Wu, W. Prellier, S. Wei, and T. Wu, *Phys. Rev. Appl.* **3**, 064015 (2015).
- [6] W. Shockley and H. J. Queisser, *J. Appl. Phys.* **32**, 510 (1961).
- [7] J. Holleman, M. M. Bishop, C. Garcia, J. S. R. Vellore Winfred, S. Lee, H. N. Lee, C. Beekman, E. Manousakis, and S. A. McGill, *Phys. Rev. B* **94**, 155129 (2016).
- [8] E. Assmann, P. Blaha, R. Laskowski, K. Held, S. Okamoto, and G. Sangiovanni, *Phys. Rev. Lett.* **110**, 078701 (2013).
- [9] M. Jellite, J.-L. Rehspringer, M. Fazio, D. Muller, G. Schmerber, G. Ferblantier, S. Colis, A. Dinia, M. Sugiyama, A. Slaoui, D. Cavalcoli, and T. Fix, *Solar Energy* **162**, 1 (2018).
- [10] M. Eckstein and P. Werner, *Phys. Rev. B* **84**, 035122 (2011).
- [11] M. Eckstein and P. Werner, *Phys. Rev. B* **88**, 075135 (2013).
- [12] P. Werner, K. Held, and M. Eckstein, *Phys. Rev. B* **90**, 235102 (2014).
- [13] F. Petocchi, S. Beck, C. Ederer, and P. Werner, *Phys. Rev. B* **100**, 075147 (2019).
- [14] M. E. Sorantin, A. Dorda, K. Held, and E. Arrigoni, *Phys. Rev. B* **97**, 115113 (2018).
- [15] A. Kauch, P. Worm, P. Prauhart, M. Innerberger, C. Watzenböck, and K. Held, *Phys. Rev. B* **102**, 245125 (2020).
- [16] E. Manousakis, *Sci. Rep.* **9**, 20395 (2019).
- [17] F. Maislinger and H. G. Evertz, *Phys. Rev. B* **105**, 045114 (2022).
- [18] R. Peierls, *Z. Phys. A: Hadrons Nucl.* **80**, 763 (1933).
- [19] N. Tsuji, T. Oka, and H. Aoki, *Phys. Rev. B* **78**, 235124 (2008).
- [20] Y. Murakami and P. Werner, *Phys. Rev. B* **98**, 075102 (2018).
- [21] T. M. Mazzocchi, P. Gazzaneo, J. Lotze, and E. Arrigoni, *Phys. Rev. B* **106**, 125123 (2022).
- [22] A. Georges and G. Kotliar, *Phys. Rev. B* **45**, 6479 (1992).
- [23] H. Aoki, N. Tsuji, M. Eckstein, M. Kollar, T. Oka, and P. Werner, *Rev. Mod. Phys.* **86**, 779 (2014).
- [24] P. Schmidt and H. Monien, *arXiv:cond-mat/0202046*.
- [25] A. V. Joura, J. K. Freericks, and T. Pruschke, *Phys. Rev. Lett.* **101**, 196401 (2008).
- [26] J. Schwinger, *J. Math. Phys.* **2**, 407 (1961).
- [27] L. V. Keldysh, *Sov. Phys.-JETP* **20**, 1018 (1965).
- [28] J. Rammer and H. Smith, *Rev. Mod. Phys.* **58**, 323 (1986).
- [29] H. Haug and A.-P. Jauho, *Quantum Kinetics in Transport and Optics of Semiconductors* (Springer, Heidelberg, 1998).
- [30] A. Picano, J. Li, and M. Eckstein, *Phys. Rev. B* **104**, 085108 (2021).
- [31] W. Metzner and D. Vollhardt, *Phys. Rev. Lett.* **62**, 324 (1989).
- [32] A. Georges, G. Kotliar, W. Krauth, and M. J. Rozenberg, *Rev. Mod. Phys.* **68**, 13 (1996).
- [33] E. Arrigoni, M. Knap, and W. von der Linden, *Phys. Rev. Lett.* **110**, 086403 (2013).
- [34] A. Dorda, M. Nuss, W. von der Linden, and E. Arrigoni, *Phys. Rev. B* **89**, 165105 (2014).
- [35] A. Dorda, M. Ganahl, H. G. Evertz, W. von der Linden, and E. Arrigoni, *Phys. Rev. B* **92**, 125145 (2015).
- [36] A. Dorda, M. Sorantin, W. von der Linden, and E. Arrigoni, *New J. Phys.* **19**, 063005 (2017).
- [37] M. Eckstein and P. Werner, *Phys. Rev. Lett.* **113**, 076405 (2014).
- [38] G. Rohringer, H. Hafermann, A. Toschi, A. A. Katanin, A. E. Antipov, M. I. Katsnelson, A. I. Lichtenstein, A. N. Rubtsov, and K. Held, *Rev. Mod. Phys.* **90**, 025003 (2018).
- [39] F. Giustino, *Rev. Mod. Phys.* **89**, 015003 (2017).