33-445 Formula Sheet

Operator Properties

$$\langle \hat{A} \rangle = \langle \psi \mid \hat{A} \mid \psi \rangle \qquad \Delta \hat{A} = \sqrt{\langle \hat{A}^2 \rangle - \langle \hat{A} \rangle^2} \qquad \hat{U} \hat{U}^\dagger = 1 \qquad [\hat{A}, \hat{B}] = \hat{A} \hat{B} - \hat{B} \hat{A} \qquad \Delta \hat{A} \Delta \hat{B} \geq \langle \frac{[\hat{A}, \hat{B}]}{2i} \rangle \qquad e^{\hat{A}} = \sum_{n=0}^{\infty} \frac{\hat{A}^n}{n!}$$

Vector Properties

$$\langle \psi \mid \psi \rangle = 1 \langle a_i \mid a_j \rangle \qquad = \delta_{ij} \hat{A} \mid \psi \rangle \qquad = \langle \psi \mid^{\dagger} \hat{A}^{\dagger} \mid \psi \rangle \qquad = \sum_n c_n \mid a_n \rangle = \begin{pmatrix} c_1 \\ \vdots \\ c_n \end{pmatrix} \langle \psi \mid \qquad = \sum_n c_n^{\dagger} \langle a_n \mid = \begin{pmatrix} c_1^{\dagger} & \dots & c_n^{\dagger} \end{pmatrix}$$

All spins

 $\vec{n} = \sin\theta\cos\varphi \vec{i} + \sin\theta\sin\varphi \vec{j} + \cos\theta \vec{k}$

$$z:\theta=0, \varphi=0 \qquad x:\theta=\frac{\pi}{2}, \varphi=0 \qquad y:\theta=\frac{\pi}{2}, \varphi=\frac{\pi}{2} \qquad [\hat{S}_{x},\hat{S}_{y}]=i\hbar\hat{S}_{z} \qquad [\hat{S}_{y},\hat{S}_{z}]=i\hbar\hat{S}_{x} \qquad [\hat{S}_{z},\hat{S}_{x}]=i\hbar\hat{S}_{y}$$

$$\hat{J}_{+}=\hat{J}_{x}+i\hat{J}_{y} \qquad \hat{J}_{-}=\hat{J}_{x}-i\hat{J}_{y} \qquad \hat{J}_{+}=\hat{J}_{-}^{\dagger} \qquad \hat{J}_{-}=\hat{J}_{+}^{\dagger}$$

$$\hat{J}_{+}\mid j,m\rangle=\hbar\sqrt{j(j+1)-m(m+1)}\mid j,m+1\rangle \qquad \hat{J}_{-}\mid j,m\rangle=\hbar\sqrt{j(j+1)-m(m-1)}\mid j,m-1\rangle$$

Spin $\frac{1}{2}$ (Electrons)

$$|+n\rangle = \begin{pmatrix} \cos\frac{\theta}{2} \\ \sin\frac{\theta}{2}e^{i\varphi} \end{pmatrix} \qquad |-n\rangle = \begin{pmatrix} \sin\frac{\theta}{2} \\ -\cos\frac{\theta}{2}e^{-i\varphi} \end{pmatrix} \qquad \hat{S}_n = \frac{\hbar}{2} \begin{pmatrix} \cos\theta & \sin\theta e^{-i\varphi} \\ \sin\theta e^{i\varphi} & -\cos\theta \end{pmatrix}$$

Spin 1 (Photons)

$$|X'\rangle = \cos\phi |X\rangle + \sin\phi |Y\rangle \qquad |R\rangle = \frac{1}{\sqrt{2}}(|X\rangle + i|Y\rangle) \qquad \hat{R}\phi\vec{k} |R\rangle = e^{i\phi} |R\rangle \qquad \hat{J}_z |R\rangle = \hbar |R\rangle$$

$$|Y'\rangle = -\sin\phi |X\rangle + \cos\phi |Y\rangle \qquad |L\rangle = \frac{1}{\sqrt{2}}(|X\rangle - i|Y\rangle) \qquad \hat{R}\phi\vec{k} |L\rangle = e^{i\phi} |L\rangle \qquad \hat{J}_z |L\rangle = -\hbar |L\rangle$$

Spin 1 (General)

$$|1,1\rangle_{n} = \begin{pmatrix} e^{-i\varphi} \frac{1+\cos\theta}{\sqrt{2}} \\ \frac{\sin\theta}{\sqrt{2}} \\ e^{i\varphi} \frac{1-\cos\theta}{2} \end{pmatrix} \quad |1,0\rangle_{n} = \begin{pmatrix} -e^{-i\varphi} \frac{\sin\theta}{\sqrt{2}} \\ \cos\theta \\ e^{i\varphi} \frac{\sin\theta}{\sqrt{2}} \end{pmatrix} \quad |1,-1\rangle_{n} = \begin{pmatrix} e^{-i\varphi} \frac{1-\cos\theta}{2} \\ -\frac{\sin\theta}{\sqrt{2}} \\ e^{i\varphi} \frac{1+\cos\theta}{2} \end{pmatrix} \quad \hat{S}_{n} = \hbar \begin{pmatrix} \cos\theta & \frac{e^{-i\varphi} \sin\theta}{\sqrt{2}} & 0 \\ \frac{e^{i\varphi} \sin\theta}{\sqrt{2}} & 0 & \frac{e^{-i\varphi} \sin\theta}{\sqrt{2}} \\ 0 & \frac{e^{i\varphi} \sin\theta}{\sqrt{2}} & -\cos\theta \end{pmatrix}$$

Hamiltonian

$$i\hbar \frac{d}{dt} \mid \psi(t) \rangle = \hat{H} \mid \psi(t) \rangle \qquad \qquad \frac{d\langle \hat{A} \rangle}{dt} = \frac{i}{\hbar} \langle [\hat{H}, \hat{A}] \rangle \qquad \qquad \hat{U}(t) \mid \psi(0) \rangle = \mid \psi(t) \rangle \qquad \qquad \hat{U}(t) = e^{-i\hat{H}\frac{t}{\hbar}}$$

Constant B_0 in the \vec{n} direction: $\hat{H} = -\frac{qB_0}{mc}\hat{S}_n = \omega_0\hat{S}_n$