

- Bowshockpy: A generator of synthetic observations for
- 2 jet-driven bowshocks
- **3 Guillermo Blázquez-Calero** [□] ^{1¶} and et al. ^{1¶}
- 1 Instituto de Astrofísica de Andalucía, CSIC, Glorieta de la Astronomía s/n, E-18008 Granada, Spain ¶
- 5 Corresponding author

DOI: 10.xxxxx/draft

Software

- Review 🗗
- Repository 🗗
- Archive ♂

Editor: Open Journals ♂ Reviewers:

@openjournals

Submitted: 01 January 1970 **Published:** unpublished

License

Authors of papers retain copyrights and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0°_{21}

In partnership with



This article and software are linked with research article DOI 10.3847/xxxxx <- update this with the DOI from AAS once you? know it., published in the Astrophysical Journal <- The name of the AAS journal... 30

Summary

Collimated jets are a common phenomenon in the Universe, appearing in a variety of astrophysical objects with accretion disks, such as black holes, planetary nebulae, and young stellar objects (YSOs). Astrophysical jets sometimes present knotty structures, which have long been suggested to be the result of a velocity variations of the flow within the jet beam (Rees, 1978). In the case of jets from YSOs, interpreting the knots as "internal working surfaces" produced by time-variable ejection, provided an explanation for the multiple bow-shaped shocks typically observed in Herbig-Haro jets: the overpressure in the internal working surface drives material sideways, which interacts with the ambient to form the bowshocks shells (Masson & Chernin, 1993; Raga et al., 1990; Stone & Norman, 1993). Current observations probing the molecular components of jets from YSOs are detecting bowshock shells with sufficient high angular and spectral resolution to reveal in details of their morphology and kinematics. The characterization of these bowshock using models provides valueable information of the interaction of the jet with the environment (Blázquez-Calero et al., under review). Bowshockpy is a Python package that generates synthetic spectral cubes, position-velocity diagrams, and moment images for a simple analytical jet-driven bowshock model, using the prescription for protostellar jets presented in Tabone et al. (2018) and Ostriker et al. (2001).

Statement of need

Jets from YSOs are thought to play a crucial role in the formation of a star by removing angular momentum from the disk and, therefore, allowing accretion onto the forming star. Additionally, jets are invoked in order to explain the low star formation efficiency at both core and cloud scales (Frank et al., 2014). Thus, characterizing the physical properties of protostellar jets and their interaction with their surrounding medium is of major importance in order to understand the star formation process. Recent observations at mm wavelengths with radio interferometers, specially the Atacama Large Milimeter Array (ALMA), enable us to study in great detail the molecular component in jets (mainly in CO and SiO emisson). Sensitive observations at high and spectral resoltion with ALMA reveal the presence of shell-like structures connecting the fast knots in the jet. While knots in the jets are usually interpreted as internal working surfaces (Tafalla et al., 2017) that expel jet material sideways by pressure forces, the shells are thought to be bowshocks that arise from the interaction of this jet material with the surrounding gas (Lee et al., 2017; Plunkett et al., 2015). Characterizing these bowshocks through models an numerical simulations (Ostriker et al., 2001; Tabone et al., 2018) can give information on the interaction of the jet with the surrounding gas, such as quantifying the velocity and mass-rate at which jet material is expeled sideways from the internal working surface, the mass-rate of ambient material incorporated into the bowshock and the ambient density (Blázquez-Calero et al., under review).

Bowshockpy is an implementation of the analytical momentum conserving bowshock model of



Tabone et al. (2018), which has been validated with hydrodynamical numerical simulations.
The software computes the observed column densities of the bowshock shell and, assuming local thermodynamic equilibrium conditions, simulates the CO intensities of low-J rotational transitions, providing mock observations of the CO emission that radio telescopes would be able to detect at milimeter wavelengths. This software provides spectral cubes, PV diagrams, and moment images of the bowshock shell fairly quick in an personal computer, allowing the modelization of observed bowshocks using just a few user-defined model and observer/instrumental parameters. To our knowledge, there is no software publicly available that allows the creation of synthetic observations of an analytical jet-driven bowshock model. We note that, although this software has been initially aimed to be used in jet-driven bowshocks in YSO objects, its applicability would be equally valid for molecular jets from protoplanetary nebulae.

55 Acknowledgements

We acknowledge the significant contribution of Alex Raga to this work, who made a preliminary version of this software in Fortran. G.A., G.B.-C., IdG-M., A.K.D.-R., G.A.F., J.F.G., M.O., acknowledge financial support from grants PID2020-114461GB-I00 and CEX2021-001131-S, funded by MCIN/AEI/10.13039/501100011033. G.B.-C., G.A.F., and M.O. acknowlege financial support from Junta de Andalucía (Spain) grant P20-00880 (FEDER, EU). G.B-C acknowledges support from grant PRE2018-086111, funded by MCIN/AEI/ 10.13039/501100011033 and by 'ESF Investing in your future', and thanks ESO Science Support Discretionary Fund for their finantial support under the 2024 SSDF 06 project.

64 References

- Frank, A., Ray, T. P., Cabrit, S., Hartigan, P., Arce, H. G., Bacciotti, F., Bally, J., Benisty,
 M., Eislöffel, J., Güdel, M., Lebedev, S., Nisini, B., & Raga, A. (2014). Jets and
 Outflows from Star to Cloud: Observations Confront Theory. In H. Beuther, R. S.
 Klessen, C. P. Dullemond, & T. Henning (Eds.), *Protostars and planets VI* (pp. 451–474).
 https://doi.org/10.2458/azu_uapress_9780816531240-ch020
- Lee, C.-F., Ho, Paul. T. P., Li, Z.-Y., Hirano, N., Zhang, Q., & Shang, H. (2017). A rotating protostellar jet launched from the innermost disk of HH 212. *Nature Astronomy*, 1, 0152. https://doi.org/10.1038/s41550-017-0152
- Masson, C. R., & Chernin, L. M. (1993). Properties of Jet-driven Molecular Outflows. 414,
 230. https://doi.org/10.1086/173071
- Ostriker, E. C., Lee, C.-F., Stone, J. M., & Mundy, L. G. (2001). A Ballistic Bow Shock Model for Jet-driven Protostellar Outflow Shells. 557(1), 443–450. https://doi.org/10.1086/321649
- Plunkett, A. L., Arce, H. G., Mardones, D., van Dokkum, P., Dunham, M. M., Fernández-López, M., Gallardo, J., & Corder, S. A. (2015). Episodic molecular outflow in the very young protostellar cluster Serpens South. 527(7576), 70–73. https://doi.org/10.1038/ nature15702
- Raga, A. C., Canto, J., Binette, L., & Calvet, N. (1990). Stellar Jets with Intrinsically Variable Sources. *364*, 601. https://doi.org/10.1086/169443
- Rees, M. J. (1978). The M87 jet: internal shocks in a plasma beam? 184, 61P-65P. https://doi.org/10.1093/mnras/184.1.61P
- Stone, J. M., & Norman, M. L. (1993). Numerical Simulations of Protostellar Jets with Nonequilibrium Cooling. I. Method and Two-dimensional Results. 413, 198. https://doi.org/10.1086/172988



- Tabone, B., Raga, A., Cabrit, S., & Pineau des Forêts, G. (2018). Interaction between a pulsating jet and a surrounding disk wind. A hydrodynamical perspective. *614*, A119. https://doi.org/10.1051/0004-6361/201732031
- Tafalla, M., Su, Y.-N., Shang, H., Johnstone, D., Zhang, Q., Santiago-García, J., Lee, C.-F., Hirano, N., & Wang, L.-Y. (2017). Anatomy of the internal bow shocks in the IRAS 04166+2706 protostellar jet. *597*, A119. https://doi.org/10.1051/0004-6361/201629493

