

Lipid rafts: at a crossroad between cell biology and physics

Ken Jacobson, Ole G. Mouritsen and Richard G. W. Anderson

Membrane lateral heterogeneity is accepted as a requirement for the function of biological membranes and the notion of lipid rafts gives specificity to this broad concept. However, the lipid raft field is now at a technical impasse because the physical tools to study biological membranes as a liquid that is ordered in space and time are still being developed. This has led to a disconnection between the concept of lipid rafts as derived from biochemical and biophysical assays and their existence in the cell. Here, we compare the concept of lipid rafts as it has emerged from the study of synthetic membranes with the reality of lateral heterogeneity in biological membranes. Further application of existing tools and the development of new tools are needed to understand the dynamic heterogeneity of biological membranes.

The origin of the raft hypothesis¹ can be traced to the perplexing discovery that glycosphingolipids cluster in the Golgi apparatus before being sorted to the apical surface of polarized epithelial cells². Subsequent studies established that glycosphingolipid clusters tend to be insoluble in Triton X-100 at 4 °C, forming detergent-resistant membranes (DRM), have a light buoyant density on sucrose gradients and are rich in both cholesterol and glycosylphosphatidyl inositol (GPI)-anchored proteins³. The size of the raft was proposed, on the basis of biophysical experiments, to be in the order of 50 nm⁴. The raft hypothesis was bolstered by the observation that synthetic membranes composed of glycosphingolipids and cholesterol recapitulate the detergent-resistant characteristics of the glycosphingolipid clusters^{5,6}, suggesting that lipid lateral heterogeneity occurs spontaneously as a function of the lipid composition of the membrane.

Two lines of inquiry emerged from these early studies: one exploring lateral heterogeneity in synthetic membranes; and the other addressing the functionality of glycosphingolipid-cholesterol rich domains in biological membranes. Few would doubt that lateral heterogeneity in biological membranes must have a major role in cell and developmental biology, but characterizing this heterogeneity in a compelling way, biochemically or biophysically, has proven to be frustratingly difficult. Indeed, the fact that a continuum of partially tenable models exists⁷ — from lipid ‘shells’⁸ to the idea that the membrane is a collection of contiguous rafts with fluid inclusions⁹ — reflects the difficulty in structurally characterizing the cell membrane¹⁰. A provisional contemporary definition of rafts that begins to embody these complexities emerged at the

2006 Keystone Symposium¹¹: “Membrane rafts are small (10–200 nm), heterogeneous, highly dynamic, sterol- and sphingolipid-enriched domains that compartmentalize cellular processes. Small rafts can sometimes be stabilized to form larger platforms through protein–protein and protein–lipid interactions.” It is our contention that the lipid-raft field is at a technical impasse, largely because the tools to study biological membranes as liquids structured in space and time are rudimentary. As a consequence, contemporary notions of lipid rafts have to be reconciled with the physical realities of biological membranes. This review will discuss the constraints that make biological membranes unusual liquid-like structures, the hierarchical organization of lipid–protein composites in biological membranes and the physical methods available for measuring such dynamically heterogeneous structures. We believe that new technologies and paradigms will be required to move the field forward. Other aspects of raft biology and biophysics have been covered in recent reviews on lipid rafts^{7,12–20} and at an interdisciplinary conference [<http://www.biophysics.org/discussions>].

Biological membranes are different from synthetic membranes

Almost to a fault, the tenets of the lipid raft hypothesis are easily demonstrated in model membrane systems, where domains can range in size from nanoscale^{21–23} to microscale^{5,6} depending on the lipid mixture. These results beg the question why are rafts so ‘elusive or illusive’ in cell membranes¹⁷. The situation in biological membranes is very different from that in artificial bilayer membranes. The biological membrane is a liquid-like structure with some long-range order and short-range order in various lengths and time scales. Short-range order is manifest in a myriad of domains whose composition, size, lifetime and functionality require characterization. For the purposes of discussion, it is useful to divide the length scales characterizing membrane-dynamic structure into shells, complexes and clusters, nanodomains and microdomains (Fig. 1a). Small dynamic structures (such as shells and clusters) and the larger nanodomains may have properties different than more stable

Ken Jacobson is in the Department of Cell and Developmental Biology & Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, 121 Taylor Hall CB#7090, Chapel Hill, NC 27599, USA. Ole G. Mouritsen is in the MEMPHYS Center for Biomembrane Physics, Physics Department, University of Southern Denmark, Campusvej 55, DK-5230 Odense, Denmark. Richard G. W. Anderson is in the Department of Cell Biology, University of Texas Southwestern Medical Center, Dallas, TX 75390–9039, USA. e-mail: frap@med.unc.edu




Figure 1 Domain-length scales and the biomembrane as a protein–lipid composite material. **(a)** Length scales of domains in biomembranes. Shells, complexes and nanoclusters range from 1–10 nm, whereas nanodomains such as caveolae can be as large as 100 nm. **(b)** A schematic representation of the biomembrane as a composite of lipids and proteins. Estimates of lateral protein concentration are about 30,000 per μm^2 based on rhodopsin in the rod outer segment^{28,29} and transmembrane proteins in the baby hamster kidney (BHK) cell membrane²⁷. Lipids were assumed to occupy a surface area of $\sim 0.68 \text{ nm}^2$ (diameter $\sim 0.93 \text{ nm}$) and an α -helix $\sim 1 \text{ nm}^2$ (diameter $\sim 1.1 \text{ nm}$). A $30 \times 30 \text{ nm}^2$ section of membrane is depicted with 32 lipids on a side, 35 transmembrane proteins with 15 single-span, 12 tetraspan and eight heptaspan α -helical proteins, having assumed crosssectional areas in

the plane of the membrane of 1 nm^2 , 4.5 nm^2 and 8 nm^2 , respectively. Taking into account the area excluded by the proteins, the numerical lipid : protein ratio is ~ 50 . For a single-span helix with a diameter of $\sim 1.1 \text{ nm}$, there are about seven lipids in the first boundary layer; for a tetraspan protein with a diameter of $\sim 2.4 \text{ nm}$, there are about 11 lipids in the first boundary layer; for a heptaspan protein (such as rhodopsin) with a diameter of $\sim 3.2 \text{ nm}$, there would be about 14 lipids in the first boundary layer. Such first-boundary layer lipids are shown in white, whereas the second layer is shown in red. All other lipids are shown in yellow. Lipid-binding proteins and adaptors linking transmembrane proteins to membrane proximate cytoskeletal filaments are also depicted as different coloured structures beneath the plane of the membrane, but ectodomains of the membrane proteins are omitted for clarity.

thermodynamic phases that, for practical purposes, have dimensions in the micrometre range (where the interfacial energy associated with the phase boundary is small compared to the free energy of the bulk phase). In addition, microemulsions may form in complex liquid mixtures that can have various characteristic sizes. They may be characterized by a lateral segregation of components, for example, in the form of two-dimensional nanoscopic ‘droplets’ embedded in a connected surrounding membrane. Microemulsions can be stabilized or destabilized by amphiphilic components that act as line surfactants at the boundaries of the droplet and can be changed by minute alterations in composition or environment.

One type of short-range spatial and temporal order, found in entities of the order of 10 nm or less, form because of significant fluctuations in local density and composition. This order is driven by cooperative molecular interactions between membrane constituents that are characteristic of liquids. These fluctuations are characterized by correlation times or the time that a fluctuation persists (that is, a lifetime) and correlation lengths, or the range over which membrane constituents, through direct and indirect interactions, can ‘sense’ each other, thereby providing a measure of domain size. A second type of order consists of more stable, disperse nanoscale structures, including shells and clusters.

In general, biomembranes have a much higher concentration of protein than most artificial membranes that have been studied in the past four decades, although our mental picture of most biological membranes, beginning with the Singer-Nicolson model²⁴, is one where the protein concentration is far lower than is actually found in cell membranes^{25,26}. Quinn *et al.* have calculated the density of integral membrane proteins in the endoplasmic reticulum and Golgi apparatus to be $\sim 30,000$ molecules per μm^2 ²⁷. The density of rhodopsin in the rod outer segment is estimated to be 30,000 per μm^2 ^{28,29}, whereas the density of the GPI-anchored folate receptor in caveolae following the addition of crosslinking antibodies has been calculated to be 32,000 per μm^2 ³⁰. At these protein concentrations, there is only enough room in the bilayer for a few rings of lipid between individual proteins (Fig. 1b).

Biological membranes also have an associated cytoskeleton and adaptor proteins that link the cytoskeleton to specific sets of membrane proteins and lipids. The combination of membrane proximate cytoskeleton adaptors, together with regulatory proteins³¹ that transiently bind to the inner leaflet surface, suggests that the region subjacent to the plasma membrane also has a very high protein concentration³², which should be considered as part of the composite material. In addition to binding membrane proteins, cytoskeletal proteins also interact with specific lipids³³ — for example, spectrin interacts with phosphatidylserine³⁴ and

Table 1 Selected evidence for membrane domains

Probe/cell ¹	Method	Membrane location	Induced ?	Domain size	Cholesterol involved	Citation
Thy-1/RBL cells	TEM-immunogold	OL	No	<100 nm	Not done	50
Thy-1 & PrP/neurons	TEM-immunogold	OL	No	<100 nm	Yes	51
DC-SIGN/ dendritic cells	TEM-immunogold	TM	No	<200 nm	Yes	95
GFP-Ras targeting domains	TEM-immunogold	IL	No	H-Ras (inactive) 44 nm	Yes	96
				K-Ras ~30 nm	No	
HA/HAb2/fibroblasts	TEM & FRET (fixed)	TM	No	Non-randomness from 6–900 nm	Yes	44
Perfringolysin-O/(cholesterol) platelets	Cytochemistry	TM	Yes — by spreading on fibrinogen or by TRAP	Filopodia several μ m long	Yes	94
Folate receptor and GFP-GPIAP/CHO cells	Homo-FRET (live)	OL	No	~4–5 nm	Yes	41
MHC class I & II and other molecules/B lymphoma cells	FRET (by flow cytometry)	TM	No	Cluster detection only	Not done	97
GFP-arginine transporter/yeast	Confocal (live)	TM	No	~300 nm	No	58
GFP-GPI & other components/Jurkat and HL60 cells	Confocal (live)	OL	Yes — by chemotractant	Several μ m	Yes	98
Laurdan/macrophages	Generalized polarization (live)	TM	No	μ m-sized	Yes	60
DMPE-Cy5/HASM	Single domain imaging (live)	OL	No	~700 nm (0.2–2 μ m)	No	59
Various fluorescent lipid analogues	Confocal (live)	OL	Yes (by cholesterol removal)	~ μ m	Yes	99
Lck, LAT, CD2/Jurkat T-cells	Single-molecule tracking (live)	IL, TM	Yes, by TCR X-linking	<250 nm	No	45
CYF-YFP acylated or prenylated probes/MDCK cells	FRET (live)	IL	No	Cluster detection only	M-P/Yes ----- G-G/No	100
EYFP-H-Ras-targeting domain/3T3 cells	SMT (live)	IL	No	~200 nm	No	101

CHO, chinese hamster ovary; OL, outer leaflet; IL, inner leaflet; TM, Transmembrane or both leaflets; G-G, geranylgeranylated; M-P, myristoylated-palmitoylated; SMT, single-molecule tracking.

other lipids³⁵, whereas adaptors containing pleckstrin homology (PH), FYVE and PX motifs bind phosphorylated derivatives of phosphatidylinositol. Indeed, modulation of PtdIns(4,5)P₂ exposure has dramatic effects on membrane–cytoskeletal interaction³⁶. Therefore, the membrane-associated cytoskeleton and its adaptors, as well as regulatory cytosolic proteins, are likely to influence the organization of membrane microdomains through both protein and lipid anchorage points on the inner leaflet. The effect of these interactions, and the high overall protein concentration in membranes, has not been fully incorporated into contemporary models of regulated lipid lateral heterogeneity in biological membranes. Thus, the membrane should be viewed as a lipid–protein composite, rather than a dilute solution of protein in a lipid solvent. Moreover, domains will rarely be isolated islands in a lipid sea, as often depicted in cartoons. The lifetimes of these domains will depend on their size and factors that may stabilize or destabilize them. These include not only lipid–lipid interactions but also protein–lipid and protein–protein interactions that occur both in the plane of the membrane, and between membrane components and elements of the cytoskeleton and pericellular matrix adjacent to the membrane²⁰.

A hierarchy of lipid–protein composites in biological membranes

Apart from the well-known membrane domains in differentiated cells (such as the apical–basolateral specializations and junctional complexes

in epithelial cells and clathrin-coated pits), considerable evidence, both experimental and conceptual, can be marshalled for the existence of other domains of various sizes in diverse biological settings. A number of these structures exhibit lipid specificity (Table 1 and Fig. 2).

The shell model⁸ proposes that specific classes of plasma-membrane proteins (such as certain transmembrane proteins and proteins anchored by lipid tails), bind to preassembled complexes of cholesterol and sphingolipid³⁷ or they may laterally organize specific lipids, including cholesterol, as in the case of peripheral^{31,38} or transmembrane proteins³⁹. Although the shell is envisioned to be stably associated with protein, component lipids can exchange with the ‘bulk’ bilayer lipids. The dynamics of exchange may range from the short-lived classical boundary layer lipids (~1–10 s)⁴⁰, in which shell lipids may rapidly interchange with non-shell lipids, to long-lived lipids that are tightly bound through specific lipid–protein interactions³⁹ and are regulated by ligands. The concept of shelled proteins, existing alone or in nanoclusters, is also supported by homo-fluorescence resonance energy transfer (FRET)⁴¹ (between like donor and acceptor) and hetero-FRET⁴² (between different donor and acceptor) measurements, which can be interpreted to show that DRM molecules are largely monomeric or coexist, but are not in an equilibrium, with a small fraction of ‘nanoclusters’ (~4–5 nm in dimension) containing four or less GPI-anchored proteins. In addition, the neurokinin-1 receptor, a G-protein-coupled receptor, was found to exist in very small (~10 nm) clusters that are cholesterol sensitive⁴³.




Figure 2 Examples of lipid and protein domains in cell membranes. (a) Single domains, enriched in the fluorescent lipid analogue DMPE-Cy5, imaged in human coronary artery smooth muscle cells⁵⁹ (reprinted with permission from the EMBO Journal). (b) Cholesterol-rich domains in platelet processes⁹⁴ visualized using a non-cytolytic derivative of biotinylated perfringolysin O followed by indirect immunofluorescence microscopy with an anti-biotin monoclonal antibody (reprinted with permission from the International Society on Thrombosis and Haemostasis). (c) Lipid domains with greater relative order than the bulk membrane, visualized in living macrophages with the fluorescent probe, Laurdan (6-Lauroyl-2-dimethylaminopthalene), where the warmer pseudo-colours represent more ordered regions⁶⁰ (reprinted with permission of the National Academy of Sciences). (d) Domains formed by the proton-arginine symporter transporter (Can1p-GFP) in *Saccharomyces cerevisiae*⁶⁸ (reprinted from Molecular Biology of the Cell with permission from The American Society for Cell Biology). The scale bars represent 1 μm in **a**, and 5 μm in **c** and **d**.

In spite of criticisms¹⁹, many of which are treated in the original postulates of the hypothesis but require proof, a virtue of the shell hypothesis is that it can accommodate seemingly disparate sets of recent data. For example, the detergent-resistant viral haemeagglutinin does not partition into lipid rafts in which it is miscible, but instead self-assembles into lipid-protein complexes exhibiting order from the nanometre to micrometre scales⁴⁴. Shelled proteins could mediate this 'self-assembly' process by enhancing the attraction between haemeagglutinin trimers. Several recent single-molecule studies also suggest that protein-protein complexes of sizes <250 nm (the resolution limit for conventional light microscopy) are fundamental units that carry out signal transduction — as opposed to these same components partitioning into lipid rafts to convey signals. Single-molecule microscopy shows that components of the immunological synapse, including Lck (a member of the Src family of protein-tyrosine kinases), linker of T-cell activation (LAT) and the coreceptor cell differentiation antigen, CD2, but not the transmembrane phosphatase CD45, exist in very small clusters that seem to be cholesterol independent⁴⁵. In this case, phosphorylated LAT functions as a scaffold mediating downstream protein-protein interactions. In the case of IgE-triggered signalling, cross-correlation microscopy has demonstrated that activated Lyn, a tyrosine kinase, and the Fc receptor form a complex, but other putative inner-leaflet raft components do not suggest a specific association

between the Fc receptor and Lyn following receptor ligation⁴⁶. In both cases, lipid shells surrounding the protein components could help promote the formation of particular complexes.

Simons and Ikonen originally proposed that lipid "...lateral organization probably results from preferential packing of sphingolipids and cholesterol into moving platforms, or rafts, onto which specific proteins attach within the bilayer."¹ Caveolae possess many of the essential properties of the proposed lipid raft; for example, they contain clusters of glycosphingolipids, GPI-anchored proteins and a high concentration of cholesterol. The caveola is a reasonably well characterized 50–100-nm nanodomain that is detergent resistant and can be readily identified in most cells by the marker caveolin-1 (ref. 47). Moreover, we know a lot about the function of caveolae in endocytosis and cell signalling. What is unclear, however, is whether the structure and function of caveolae depends on lateral interactions between cholesterol and sphingolipids alone, or on additional protein-lipid interactions. Nevertheless, caveolae fit the original definition of a lipid raft and they may be used as a provisional standard against which other lipid rafts can be judged.

The strongest evidence for non-caveolae nanodomains (50–200 nm in dimension in the outer leaflet of the plasma membrane) comes from immunogold electron microscopy of components of, for example, the T cell receptor (TCR)^{18,48} in fixed cells (Table 1), where labelled proteins are detected in clusters. The GPI-anchored protein, Thy-1, is observed in such clusters both on rat basophilic leukaemia (RBL) cells^{49,50} and on neurons⁵¹. However, Thy-1, ganglioside 1 (GM-1), LAT or the Fc receptor show little colocalization, even though substantial amounts of these components are enriched in the Triton X-100 DRM fraction⁴⁹, indicating that detergent resistance reduces the inherent lateral complexity of the membrane to one simplified, and therefore misleading, parameter⁵². Part of the difficulty in having full confidence that such nanoscopic domains exist, however, is that measurements in living cells are more consistent with a high percentage of these proteins being monomeric. Fluorescence recovery after photobleaching (FRAP) measurements of inner- and outer-leaflet proteins give diffusion coefficients that correlate with the type of membrane anchorage (GPI, transmembrane and acylation or prenylation), not whether the proteins are detergent insoluble⁵³, whereas in the simple raft concept all raft molecules should have similar diffusion coefficients as they belong to similar domains. Moreover, single-molecule tracing of the transmembrane histocompatibility antigen is consistent with monomer diffusion⁵⁴. Although some of the components involved in TCR signalling do exhibit very low diffusion coefficients, consistent with immobile clusters, others do not and there is no correlation with their presence in the Triton X-100 insoluble fraction⁴⁵. In addition, FRAP measurements of inner-leaflet components show they exhibit rapid, complete recovery^{53,55} after bleaching. This can be reconciled with the inner-leaflet clusters of the small GTPase, Ras, revealed by transmission electron microscopy (TEM) on fixed cells if it is assumed that activated Ras drives the formation of very transient nanoclusters from which downstream signalling ensues⁵⁶. Single-molecule tracking and FRET studies of activated Ras are consistent with this proposal⁵⁷.

Lipid and protein reporters imaged at the light and electron microscope level by conventional and advanced means provide evidence for the existence of stable domains approaching a micrometre in dimension, or greater, in various cells (Fig. 2). Microdomains of detergent-resistant transporters were stable in growing yeast cells for more than ten minutes⁵⁸, although the exchange time of individual proteins was not determined. Microdomains in smooth muscle cells⁵⁹ and macrophages⁶⁰ were stable

Table 2 Some methods for detecting and characterizing membrane domains in live cells

Method	Primary observables	Spatial/temporal resolution	Comments	Representative citations
FCS and ICS	Fluorophore translational mobility & dynamics of lateral heterogeneity	~250 nm/ ~1 μ s	Highly sensitive to clustering, single and multiple colours	46, 102–105
FRET	Donor-acceptor proximity	~5–10 nm separation detected (no size)/<1 s	Cluster identification at nm scale	41–43, 74, 106
SPT	Translational trajectory of particle (including 40 nm gold particles & quantum dots)	~250 nm (≤ 10 nm) ^a ≤ 25 μ s	Single particle sensitivity detects domains as confinement zones	13, 107, 108
SMFI	Single-molecule fluorescence	~250 nm (20 nm) ^a / ≤ 30 ms	Single molecule sensitivity, detects clusters by intensity & confinement	14, 45, 101, 109, 110
TOCCSL	Clusters or monomers	~250 nm/<500 ms	Detects whether molecules entering photobleached region are singles or multimers	69
Thermal noise imaging	Laser-trapped particle attached to membrane component	~10 nm/~ μ s	Explores positions sampled by the particle while it is trapped by laser	111

FCS, fluorescence correlation spectroscopy; ICS, image correlation spectroscopy; SPT, single-particle tracking; SMFI, single-molecule fluorescence imaging; TOCCSL, thinning out clusters while conserving stoichiometry of labelling. ^aPrecision in detecting centroid of particle image.

for tens of seconds; moreover, in muscle they were observed as cohesive entities that rotate as units, whereas individual lipid-probe molecules had a domain residence time of ~10 ms⁵⁹. Thus, it seems that stable, larger domains can exist in various cells but the stabilizing factors²⁰ are largely unknown. In some instances, proteins from the four-transmembrane domain tetraspanin superfamily⁶¹ may organize special microdomains as in microvillae⁶² and HIV may induce tetraspanin-enriched domains as export portals in infected cells⁶³. Caveolae may also form micrometre sized clusters⁶⁴ in some cells. The importance of the lipid phase in developing longer-range order in the plasma membrane has been illustrated by two recent studies. Electron-spin resonance results from several intact living-cell types indicate that a large fraction of outer-leaflet phospholipids is in the liquid-ordered state at temperatures near 5 °C and decreases as temperature increases to 37 °C⁶⁵. Although the individual ordered units have a lower size limit of ~2 nm, the authors suggest that the liquid-ordered state may actually be the contiguous (percolating) phase in the outer leaflet⁹. A similar view has been advanced based on interpretation of FRAP results on the apical surface of polarized epithelial cells, but not fibroblasts, at temperatures ~<25 °C; at 37 °C, the non-raft, liquid-disordered phase becomes the percolating phase⁶⁶.

Biophysical tools to study membrane domains in biological membranes

To be biologically significant, domains should have a minimum size of a few protein diameters and a minimum lifetime corresponding to a short enzyme turnover time of ~microseconds. When domains are stable and of sufficient size, detection is possible using both traditional and advanced imaging techniques (Fig. 2). But, when domains are small and transient, the biophysical challenges of measuring them are similar to those facing physicists a half-century ago when they attempted to describe the structure of liquids. Predictably, the most controversial area of membrane lateral organization is on the nanoscale level, where technology with sufficient simultaneous spatial and temporal resolution is not available. Thus, refining existing methods and developing new ones is now key to progress^{67,68}. As the goal is to study dynamic membrane structure in living cells, applicable techniques, for the most part, rely on

fluorescence microscopy because of the high sensitivity of these methods and their facile application to single, living cells.

In Table 2, we enumerate several techniques with the potential to most directly detect and characterize dynamic heterogeneity in living cells, together with all-important estimates of temporal and spatial resolution. It is likely that advances will be made in characterizing dynamic domain structure using both the single molecule and particle techniques, and advanced fluorescence correlation microscopy. The latter should be useful in characterizing domains that are not visible by ordinary fluorescence microscopy but can be detected using other specialized techniques^{41,59,69}. Moreover, there are several emerging technologies that 'beat' the resolution limit of fluorescence microscopy⁷⁰. There are also a number of developing techniques with great potential to provide complementary information. These include secondary ion mass spectrometry on the 100-nm scale^{71,72} to quantitatively determine composition of microdomains *in situ* with submicrometre resolution, and atomic force microscopy (AFM) and near field scanning microscopy (NSOM) for membrane topography and microcomposition^{73–75}. Such techniques can complement fluorescence microscopy, but are limited in that they either operate on quick-frozen specimens (nano-SIMS) or they lack the time resolution of the fluorescence techniques (AFM and NSOM). Provided that adequate contrast between structures can be obtained, the ultimate tools to determine the length scale of dynamic clusters, nanodomains and rafts in the nanometre range are scattering techniques, such as neutron and X-ray scattering^{76,77}. However, because of difficulties in sample preparation, these techniques have so far only been used in model-membrane studies.

Other challenges

As current biophysical tools mature and new methods emerge, we must not lose sight of the fact that energy must constantly be supplied to biological membranes⁷⁸, both directly through the expenditure of high-energy intermediates and indirectly through intracellular and extracellular transport processes that constantly add and remove lipids and proteins^{79,80}. This leads to creation of an active region^{7,12} with scale-dependent organization and function that depends on mutual interactions between lipids and proteins^{46,81}.

The caveolae nanodomain illustrates the challenge of understanding the role of energy dissipation in thermodynamically open systems⁸². In addition to whatever lateral forces are required for lipid organization in a caveola, there must also be a constant flow of cholesterol through the domain for it to function⁸³. The mechanism of cholesterol supply may involve the movement of caveolin-1 from the endoplasmic reticulum⁸⁴ and lipid droplets^{85,86} to caveolae, although the precise details remain unknown. Shutting off the supply of cholesterol causes caveolae to malfunction, both during endocytosis⁸⁷ and signal transduction⁸⁶. Studies of caveolin-1-expressing cells shows that cholesterol rapidly moves from its site of synthesis in the endoplasmic reticulum across the caveolae bilayer into contiguous membrane, as well as to extracellular lipidic acceptors like high-density lipoprotein (HDL), with a half-time of ~6 min⁸⁴. Even though cholesterol flux seems to be critical, biophysical calculations fail to explain why. Given the large number of animal genes devoted to moving cholesterol around in cells and between tissues, transmembrane cholesterol flux most likely is common, and therefore important for membrane organization. In fact, the energy-dependent turnover of lipids in general is probably critical for the normal function of membranes and membrane domains. The challenge is to incorporate these parameters into future models of nanodomain structure and function, determine how lipid fluxes contribute to nanodomain order and develop new methods to detect lipid fluxes in cell membranes.

Because of the dynamic composition of local membrane structures, there may not be a consensus non-caveolae nanodomain. In addition to the lipid and protein composition, the characteristics of nanodomains may be strongly dependent on available soluble and pericellular matrix-bound ligands for membrane receptors, as well as on transmembrane potential and mechanical stress. We also need to determine whether microdomains are pre-existent structures into which certain components partition⁸⁸, or whether they are preassembled protein-lipid complexes that are actively inserted into the membrane^{7,20,44} but induced to cluster into domains by specific stimuli. Indeed, the field has been moving toward the concept that lipid-shelled proteins and protein nanoclusters can be triggered to aggregate by external influences, such as ligands, to produce larger functional domains^{8,13,89,90}.

We need to assess the degree to which appropriate artificial membranes, at equilibrium or near equilibrium, can be used to study aspects of the biological membrane. In this regard, it will be important to ask how the principles apparent in relatively simple model lipid-membrane systems^{19,88}, many of which are quite intriguing⁹¹, are reflected in the complexity of cell membranes. However, investigators seeking more relevant model membranes will need to deal with the issue of protein concentrations and the natural lipid asymmetry of biomembranes¹⁹, as well as energetics. Model membranes with high protein concentration are used in nuclear magnetic resonance (NMR)⁹², electron microscopy and diffraction investigations. Such membranes may be useful in revisiting the boundary layer and shell issue by using a more expanded set of proteins and lipids than in the original studies. Ultimately, we will need to develop novel lipid-protein model systems with functioning proteins (such as channels, multienzymes systems, etc.) that have some of the properties of biological membranes, but that are also sufficiently well-defined to study domain properties such as formation, size, lifetime and morphology. Finally, there is a clear need for the development of new computational models and algorithms (for example, based on dissipative particle dynamics and molecular dynamics) that will allow

improved computer simulations of the large assemblies of molecules found in domains⁹³. The complementary expertise of biophysicists and cell biologists will be required to make meaningful progress in understanding membrane lateral heterogeneity. □

ACKNOWLEDGEMENTS

K.J. thanks A. G. Lee, G. Schutz and H. Stockinger for stimulating comments. We also thank A. Diehl and B. Yang for artwork and assistance in preparing the manuscript. This work was supported by the National Institutes of Health (NIH) grant GM 41402 (K.J.), the Kenan Distinguished Professorship (K.J.), NIH grant HL 20948 (R.A.), NIH grant GM 52016 (R.A.), the Perot Family Foundation (R.A.) and the Cecil H. Green Distinguished Chair in Cellular and Molecular Biology (R.A.). MEMPHYS-Center for Biomembrane Physics is supported by the Danish National Research Foundation (O.G.M.).

1. Simons, K. & Ikonen, E. Functional rafts in cell membranes. *Nature* **387**, 569–572 (1997).
2. Simons, K. & van Meer, G. Lipid sorting in epithelial cells. *Biochemistry* **27**, 6197–6202 (1988).
3. Brown, D. GPI-anchored proteins and detergent-resistant membrane domains. *Braz. J. Med. Biol. Res.* **27**, 309–315 (1994).
4. Pralle, A., Florin, E.-L., Simons, K. & Horber, J. K. H. Sphingolipid-cholesterol rafts diffuse as small entities in the plasma membrane of mammalian cells. *J. Cell Biol.* **148**, 997–1007 (2000).
5. Dietrich, C. *et al.* Lipid rafts reconstituted in model membranes. *Biophys. J.* **80**, 1417–1428 (2001).
6. Dietrich, C., Volovyk, Z. N., Levi, M., Thompson, N. L. & Jacobson, K. Partitioning of Thy-1, GM1, and cross-linked phospholipids into lipid rafts reconstituted in supported model membrane monolayers. *Proc. Natl Acad. Sci. USA* **98**, 10642–10647 (2001).
7. Mayor, S. & Rao, M. Rafts: scale-dependent, active lipid organization at the cell surface. *Traffic* **5**, 231–240 (2004).
8. Anderson, R. G. & Jacobson, K. A role for lipid shells in targeting proteins to caveolae, rafts, and other lipid domains. *Science* **296**, 1821–1825 (2002).
9. Maxfield, F. R. Plasma membrane microdomains. *Curr. Opin. Cell Biol.* **14**, 483–487 (2002).
10. Mouritsen, O. G. *Life as a matter of fat. The emerging science of lipidomics*. (Springer-Verlag, Heidelberg, 2005).
11. Pike, L. J. Rafts defined: a report on the Keystone symposium on lipid rafts and cell function. *J. Lipid Res.* **47**, 1597–1598 (2006).
12. Edidin, M. The state of lipid rafts: From model membranes to cells. *Annual Rev. Biophys. Biomol. Struct.* **32**, 257–283 (2003).
13. Kusumi, A. *et al.* Paradigm shift of the plasma membrane concept from the two-dimensional continuum fluid to the partitioned fluid: high-speed single-molecules tracking of membrane molecules. *Annu. Rev. Biophys. Biomol. Struct.* **34**, 351–378 (2005).
14. Lommerse, P. H. M., Spaik, H. P. & Schmidt, T. *In vivo* plasma membrane organization: results of biophysical approaches. *Biochim. Biophys. Acta* **2004**, 119–131 (2004).
15. McMullen, T., Lewis, R. & McElhaney, R. Cholesterol-phospholipid interactions, the liquid-ordered phase and lipid rafts in model and biological membranes. *Curr. Opin. Coll. Inter. Sci.* **8**, 459–468 (2004).
16. Mukherjee, S. & Maxfield, F. R. Membrane domains. *Annu. Rev. Cell Dev. Biol.* **20**, 839–866 (2004).
17. Munro, S. Lipid rafts: elusive or illusive? *Cell* **115**, 377–388 (2003).
18. Parton, R. G. & Hancock, J. F. Lipid rafts and plasma membrane microorganization: insights from Ras. *Trends Cell Biol.* **14**, 141–147 (2004).
19. Simons, K. & Vaz, W. Model systems, lipid rafts, and cell membranes. *Ann. Rev. Biophys. Biomol. Struct.* **33**, 269–295 (2004).
20. Veréb, G. *et al.* Dynamic, yet structured: The cell membrane the decades after the Singer-Nicolson model. *Proc. Natl Acad. Sci. USA* **100**, 8053–8058 (2003).
21. de Almeida, R. F., Loura, L., Fedorov, A. & Prieto, M. Lipid rafts have different sizes depending on membrane composition: A time-resolved fluorescence resonance energy transfer study. *J. Mol. Biol.* **346**, 1109–1120 (2005).
22. Hsueh, Y. W., Gilbert, K., Trandum, C., Zuckermann, M. & Thewalt, J. The effect of ergosterol on dipalmitoylphosphatidylcholine bilayers: a deuterium NMR and calorimetric study. *Biophys. J.* **88**, 1799–1808 (2005).
23. Veatch, S. L., Polozov, I. V., Gawrisch, K. & Keller, S. L. Liquid domains in vesicles investigated by NMR and fluorescence microscopy. *Biophys. J.* **86**, 2910–2922 (2004).
24. Singer, S. J. & Nicolson, G. L. The fluid mosaic model of the structure of cell membranes. *Science* **175**, 720–731 (1972).
25. Engleman, D. M. Membranes are more mosaic than fluid. *Nature* **438**, 578–580 (2005).
26. Ryan, T. A., Myers, J., Holowka, D. A., Baird, B. A. & Webb, W. W. Molecular crowding on the cell surface. *Science* **239**, 61–64 (1988).
27. Quinn, P., Griffiths, G. & Warren, G. Density of newly synthesized plasma membrane proteins in intracellular membranes II. Biochemical studies. *J. Cell Biol.* **98**, 2142–2147 (1984).
28. Liang, Y. *et al.* Organization of the G protein-coupled receptors rhodopsin and opsin in native membranes. *J. Biol. Chem.* **278**, 21655–21662 (2003).
29. Lieberman, P. A. & Entine, G. Lateral diffusion of visual pigment in photoreceptor disk membranes. *Science* **185**, 457–459 (1974).

30. Rothberg, K. G., Ying, Y. S., Kamen, B. A. & Anderson, R. G. Cholesterol controls the clustering of the glycosphospholipid-anchored membrane receptor for 5-methyltetrahydrofolate. *J. Cell Biol.* **111**, 2931–2938 (1990).
31. McLaughlin, S. & Murray, D. Plasma membrane phosphoinositide organization by protein electrostatics. *Nature* **438**, 605–611 (2005).
32. Sheetz, M. P. Cell control by membrane-cytoskeleton adhesion. *Nature Rev. Mol. Cell Biol.* **2**, 392–396 (2001).
33. Janmey, P. A. & Lindberg, U. Cytoskeletal regulation: rich in lipids. *Nature Rev. Mol. Cell Biol.* **5**, 658–666 (2004).
34. Maksymiuk, R., Sui, S. -F., Gaub, H. & Sackmann, E. Electrostatic coupling of spectrin dimers to phosphatidylserine containing lipid lamellae. *Biochemistry* **26**, 2983–2990 (1987).
35. Grzybek, M. *et al.* Spectrin-phospholipid interactions. Existence of multiple kinds of binding sites? *Chem. Phys. Lipids* **141**, 133–141 (2006).
36. Kwik, J. *et al.* Membrane cholesterol, lateral mobility, and the phosphatidylinositol 4,5-bisphosphate-dependent organization of cell actin. *Proc. Natl Acad. Sci. USA* **100**, 13964–13969 (2003).
37. McConnell, H. & Radhakrishnan, A. Condensed complexes of cholesterol and phospholipids. *Biochim. Biophys. Acta* **1610**, 159–173 (2003).
38. McLaughlin, S. How clusters of basic/hydrophobic residues on proteins interact with lipids in membranes. *Biophys. J.* **SP23** (2004).
39. Lee, A. G. Lipid-protein interactions in biological membranes: a structural perspective. *Biochim. Biophys. Acta* **1612**, 1–40 (2003).
40. Marsh, D. & Pali, T. Lipid conformation in crystalline bilayers and in crystals of trans-membrane proteins. *Chem. Phys. Lipids* **141**, 48–65 (2006).
41. Sharma, P. *et al.* Nanoscale organization of multiple GPI-anchored proteins in living cell membranes. *Cell* **116**, 577–589 (2004).
42. Kenworthy, A. K. & Edidin, M. Distribution of a glycosylphosphatidylinositol anchored protein at the apical surface of MDCK cells examined at a resolution of <100Å using imaging fluorescence resonance energy transfer. *J. Cell Biol.* **142**, 69–84 (1998).
43. Meyer, B. H. *et al.* FRET imaging reveals that functional neurokinin-1 receptors are monomeric and reside in membrane microdomains of live cells. *Proc. Natl. Acad. Sci. USA* **103**, 2138–2143 (2006).
44. Hess, S. T. *et al.* Quantitative electron microscopy and fluorescence spectroscopy of the membrane distribution of influenza hemagglutinin. *J. Cell Biol.* **169**, 965–976 (2005).
45. Douglas, A. D. & Vale, R. D. Single-molecule microscopy reveals plasma membrane microdomains created by protein-protein networks that exclude or trap signaling molecules in T cells. *Cell* **121**, 937–950 (2005).
46. Larson, D. R., Gosse, J. A., Holowka, D. A., Baird, B. A. & Webb, W. W. Temporally resolved interactions between antigen-stimulated IgE receptors and Lyn kinase on living cells. *J. Cell Biol.* **171**, 527–536 (2005).
47. Parton, R. G. Caveolae—from ultrastructure to molecular mechanisms. *Nature Rev. Mol. Cell Biol.* **4**, 162–167 (2003).
48. Schade, A. E. & Levine, A. D. Lipid raft heterogeneity in human peripheral blood T lymphoblasts: A mechanism for regulating the initiation of TCR signal transduction. *J. Immunol.* **168**, 2233–2239 (2002).
49. Wilson, B. *et al.* Markers for detergent-resistant lipid rafts occupy distinct and dynamic domains in native membranes. *Mol. Biol. Cell* **15**, 2580–2592 (2004).
50. Wilson, B. S., Pfeiffer, J. R. & Oliver, J. M. Observing FcεRI signaling from the inside of the mast cell membrane. *J. Cell Biol.* **149**, 1131–1142 (2000).
51. Bruegger, B. *et al.* The membrane domains occupied by glycosylphosphatidylinositol-anchored prion protein and Thy-1 differ in lipid composition. *J. Biol. Chem.* **279**, 7530–7536 (2004).
52. Lichtenberg, D., Goni, F. M. & Heerklotz, H. Detergent-resistant membranes should not be identified with membrane rafts. *Trends Biochem. Sci.* **30**, 430–436 (2005).
53. Kenworthy, A. K. *et al.* Dynamics of putative raft-associated proteins at the cell surface. *J. Cell Biol.* **165**, 735–746 (2004).
54. Vrljic, M., Nishimura, S. Y., Brasselet, S., Moerner, W. E. & McConnell, H. M. Translational diffusion of individual class II MHC membrane proteins in cells. *Biophys. J.* **83**, 2681–2692 (2002).
55. Niv, H., Gutman, O., Kloog, Y. & Henis, Y. I. Activated K-Ras and H-Ras display different interactions with saturable nonraft sites at the surface of live cells. *J. Cell Biol.* **157**, 865–872 (2002).
56. Plowman, S. J., Muncke, C., Parton, R. G. & Hancock, J. F. H-Ras, K-Ras, and inner plasma membrane raft proteins operate in nanoclusters with differential dependence on the actin cytoskeleton. *Proc. Natl Acad. Sci. USA* **102**, 15500–15505 (2005).
57. Murakoshi, H. *et al.* Single-molecule imaging analysis of Ras activation in living cells. *Proc. Natl Acad. Sci. USA* **101**, 7317–7322 (2004).
58. Malinska, K., Malinsky, J., Opekarova, M. & Tanner, W. Visualization of protein compartmentation within the plasma membrane of living yeast cells. *Mol. Biol. Cell* **14**, 4427–4436 (2003).
59. Schutz, G. J., Kada, G., Pastushenko, V. P. & Schindler, H. Properties of lipid microdomains in a muscle cell membrane visualized by single molecule microscopy. *EMBO J.* **19**, 892–901 (2000).
60. Gaus, K. *et al.* Visualizing lipid structure and raft domains in living cells with two-photon microscopy. *Proc. Natl Acad. Sci. USA* **100**, 15554–15559 (2003).
61. Hemler, M. E. Tetraspanin functions and associated microdomains. *Nature Rev. Mol. Cell Biol.* **6**, 801–811 (2005).
62. Roper, K., Corbett, D. & Huttner, W. B. Retention of prominin in microvilli reveals distinct cholesterol-based lipid microdomains in the apical plasma membrane. *Nature Cell Biol.* **2**, 582–592 (2000).
63. Nydegger, S., Khurana, S., Kremensov, D. N., Foti, M. & Thali, M. Mapping of tetraspanin-enriched microdomains that can function as gateways for HIV-1. *J. Cell Biol.* **173**, 795–807 (2006).
64. Rothberg, K. G. *et al.* Caveolin, a protein component of caveolae membrane coats. *Cell* **68**, 673–682 (1992).
65. Swamy, M. J. *et al.* Coexisting domains in the plasma membranes of live cells characterized by spin-label ESR spectroscopy. *Biophys. J.* **90**, 4452–4465 (2006).
66. Meder, D., Moreno, M. J., Verkade, P., Vaz, W. L. & Simons, K. Phase coexistence and connectivity in the apical membrane of polarized epithelial cells. *Proc. Natl Acad. Sci. USA* **103**, 329–334 (2006).
67. Lagerholm, B. C., Weinreb, G. E., Jacobson, K. & Thompson, N. L. Detecting microdomains in intact cell membranes. *Annu. Rev. Phys. Chem.* **56**, 309–336 (2005).
68. Marguet, D., Lenne, P. F., Rigneault, H. & He, H. T. Dynamics in the plasma membrane: how to combine fluidity and order. *EMBO J.* **25**, 3446–3457 (2006).
69. Moertelmaier, M., Brameshuber, M., Linimeier, M., Schuetz, G. & Stockinger, H. Thinning out clusters while conserving stoichiometry of labeling. *Appl. Phys. Lett.* **87**, 263903 (2005).
70. Egner, A. & Hell, S. W. Fluorescence microscopy with super-resolved optical sections. *Trends Cell Biol.* **15**, 207–215 (2005).
71. Marxer, C. G., Kraft, M. L., Weber, P. K., Hutcheon, I. D. & Boxer, S. G. Supported membrane composition analysis by secondary ion mass spectrometry with high lateral resolution. *Biophys. J.* **88**, 2965–2975 (2005).
72. Winograd, N. The magic of cluster SIMS. *Anal. Chem.* **77**, 142A–149A (2005).
73. Koopman, M. *et al.* Near-field scanning optical microscopy in liquid for high resolution single molecule detection on dendritic cells. *FEBS Lett.* **573**, 6–10 (2004).
74. Krishnan, R. V., Varma, R. & Mayor, S. Fluorescence methods to probe nanometre-scale organization of molecules in living cell membranes. *J. Fluor.* **11**, 211–226 (2001).
75. Lesniewska, E., Milhiet, P. E., Gicondi, M. C. & Le Grimmellec, C. Atomic force microscope imaging of cells and membranes. *Methods Cell Biol.* **68**, 51–65 (2002).
76. Nagle, J. Could diffuse scattering of x-rays and/or neutrons provide structural information about rafts? *Biophys. J.* **SP28** (2004).
77. Pencer, J. *et al.* Detection of sub-micron raft-like domains in membranes by small-angle neutron scattering. *Eur. Phys. J.* **18**, 447–458 (2005).
78. Morowitz, H. *Energy Flow in Biology* (Ox Bow Press, Woodbridge, CT; 1968).
79. Pfeffer, S. Membrane domains in the secretory and endocytic pathways. *Cell* **112**, 507–517 (2003).
80. Gheber, L. A. & Edidin, M. A model for membrane patchiness: lateral diffusion in the presence of barriers and vesicle traffic. *Biophys. J.* **77**, 3163–3175 (1999).
81. Sabra, M. C. & Mouritsen, O. G. Steady-state compartmentalization of lipid membranes by active proteins. *Biophys. J.* **74**, 745–752 (1998).
82. Nicolis, G. & Prigogine, I. *Exploring Complexity* (W. H. Freeman, New York, 1989).
83. Anderson, R. G. The caveolae membrane system. *Annu. Rev. Biochem.* **67**, 199–225 (1998).
84. Smart, E. J., Ying, Y., Donzell, W. C. & Anderson, R. G. A role for caveolin in transport of cholesterol from endoplasmic reticulum to plasma membrane. *J. Biol. Chem.* **271**, 29427–29435 (1996).
85. Pol, A. *et al.* Dynamic and regulated association of caveolin with lipid bodies: modulation of lipid body motility and function by a dominant negative mutant. *Mol. Biol. Cell* **15**, 99–110 (2004).
86. Roy, S. *et al.* Dominant-negative caveolin inhibits H-Ras function by disrupting cholesterol-rich plasma membrane domains. *Nature Cell Biol.* **1**, 98–105 (1999).
87. Chang, W. J., Rothberg, K. G., Kamen, B. A. & Anderson, R. G. Lowering the cholesterol content of MA104 cells inhibits receptor-mediated transport of folate. *J. Cell Biol.* **118**, 63–69 (1992).
88. Silvius, J. R. Partitioning of membrane molecules between raft and non-raft domains: Insights from model-membrane studies. *Biochim. Biophys. Acta* **1746**, 193–202 (2005).
89. Simons, K. & Toomre, D. Lipid rafts and signal transduction. *Nature Rev. Mol. Cell Biol.* **1**, 31–39 (2000).
90. Chen, Y., Thelin, W., Yang, B., Milgram, S. & Jacobson, K. Transient anchorage of cross-linked glycosyl-phosphatidylinositol-anchored proteins depends on cholesterol, Src family kinases, caveolin, and phosphoinositides. *J. Cell Biol.* **175**, 169–178 (2006).
91. Hammond, A. T. *et al.* Crosslinking a lipid raft component triggers liquid ordered/liquid disordered phase separation in model plasma membranes. *Proc. Natl Acad. Sci. USA* **102**, 6320–6325 (2005).
92. De Angelis, A. A. *et al.* NMR experiments on aligned samples of membrane proteins. *Methods Enzymol.* **394**, 350–382 (2005).
93. Nicolau, D. V., Jr., Burrage, K., Parton, R. G. & Hancock, J. F. Identifying optimal lipid raft characteristics required to promote nanoscale protein-protein interactions on the plasma membrane. *Mol. Cell Biol.* **26**, 313–323 (2006).
94. Heijnen, H. F. G. *et al.* Concentration of rafts in platelet filopodia correlates with recruitment of c-Src and CD63 to these domains. *J. Thromb. Haemos.* **1**, 1161–1173 (2003).
95. Cambi, A. *et al.* Microdomains of the C-type lectin DC-SIGN are portals for virus entry into dendritic cells. *J. Cell Biol.* **164**, 145–155 (2004).
96. Prior, I., Muncke, C., Parton, R. & Hancock, J. Direct visualization of Ras proteins in spatially distinct cell surface microdomains. *J. Cell Biol.* **160**, 165–170 (2003).
97. Szollosi, J., Horejsi, V., Bene, L., Angelisova, P. & Damjanovich, S. Supramolecular complexes of MHC class I, MHC class II, CD20, and tetraspanin molecules (CD53, CD81, and CD82) at the surface of a B cell line JY. *J. Immunol.* **157**, 2939–2946 (1996).
98. Gomez-Mouton, C. *et al.* Dynamic redistribution of raft domains as an organizing platform for signaling during cell chemotaxis. *J. Cell Biol.* **164**, 759–768 (2004).
99. Hao, M., Mukherjee, S. & Maxfield, F. R. Cholesterol depletion induces large scale domain segregation in living cell membranes. *Proc. Natl Acad. Sci. USA* **98**, 13072–13077 (2001).

100. Zacharias, D. A., Violin, J. D., Newton, A. C. & Tsien, R. Y. Partitioning of lipid-modified monomeric GFPs into membrane microdomains of live cells. *Science* **296**, 913–916 (2002).
101. Lommerse, P. H. *et al.* Single-molecule imaging of the H-ras membrane-anchor reveals domains in the cytoplasmic leaflet of the cell membrane. *Biophys. J.* **86**, 609–616 (2004).
102. Herbert, B., Constantino, S. & Wiseman, P. W. Spatiotemporal image correlation spectroscopy (STICS) theory, verification, and application to protein velocity mapping in living CHO cells. *Biophys. J.* **88**, 3601–3614 (2005).
103. Sanchez, S. A. & Gratton, E. Lipid–protein interactions revealed by two-photon microscopy and fluorescence correlation spectroscopy. *Acc. Chem. Res.* **38**, 469–477 (2005).
104. Wawrezinieck, L., Rigneault, H., Marguet, D. & Lenne, P.-F. Fluorescence correlation spectroscopy diffusion laws to probe the submicron cell membrane organization. *Biophys. J.* **89**, 4029–4042 (2005).
105. Wiseman, P. W. *et al.* Spatial mapping of integrin interactions and dynamics during cell migration by image correlation microscopy. *J. Cell Sci.* **117**, 5521–5534 (2004).
106. Acasandrea, M. A., Dale, R. E., van de Ven, M. & Ameloot, M.. Twodimensional Förster resonance energy transfer (2-D FRET) and the membrane raft hypothesis. *Chem. Phys. Lett.* **419**, 469–473 (2006).
107. Dahan, M. *et al.* Diffusion dynamics of glycine receptors revealed by singlequantum dot tracking. *Science* **302**, 442–445 (2003).
108. Dietrich, C., Yang, B., Fujiwara, T., Kusumi, A. & Jacobson, K. Relationship of lipid rafts to transient confinement zones detected by single particle tracking. *Biophys. J.* **82**, 274–284 (2002).
109. Cherry, R. J. *et al.* Measurements of associations of cell-surface receptors by single-particle fluorescence imaging. *Biochem. Soc. Trans.* **31**, 1028–1031 (2003).
110. Vrljic, M., Nishimura, S. Y., Moerner, W. E. & McConnell, H. M. Cholesterol depletion suppresses the translational diffusion of class II major histocompatibility complex proteins in the plasma membrane. *Biophys. J.* **88**, 334–347 (2005).
111. Lukic, B. *et al.* Direct observation of nondiffusive motion of a Brownian particle. *Phys. Rev. Lett.* **95**, 160601 (2005).