

Methods of Homotopy Theory in Algebraic Geometry from the Viewpoint of Cohomology Operations

Tongtong Liang

Southern University of Science and Technology, China (SUSTech)

July 25, 2022

Outline

- 1 Background
- 2 Power operations in topology
- 3 Power operations in algebraic geometry
- 4 Questions for further investigation

Motivation: methods of homotopy theory


the study of objects in geometry and topology

methods of homotopy theory



the study of related homotopy classes

Motivation: methods of homotopy theory



Examples of methods of homotopy theory

Theorem (Steenrod 1951)

Let X be a paracompact space and G be a topological group, then

$$\mathcal{B}un_G(X) \cong [X, BG]$$

Examples of methods of homotopy theory

Theorem (Steenrod 1951)

Let X be a paracompact space and G be a topological group, then

$$\mathcal{B}un_G(X) \cong [X, BG]$$

Theorem (Thom 1954)

Let G be a subgroup of $GL(F, k)$ for $F = \mathbb{R}, \mathbb{C}$, or \mathbb{H} . Let X be a manifold, then

$$\{\text{cobordism classes of } G\text{-submanifolds in } X\} \cong [X, MG]$$

Motivation: computational tools


the study of objects in geometry and topology

methods of homotopy theory


capture sufficient geometric features

the study of related homotopy classes

Motivation: computational tools



Motivation: computational tools



How to process homotopy classes

(Multiplicative) cohomology theory $E^*: X \mapsto E^*(X)$ a \mathbb{Z} -graded module (algebra). (contravariant functors)

$$[X, Y] \xrightarrow{E^*} \mathbf{Maps}(E^*Y, E^*X)$$

graded modules (algebras)

How to process homotopy classes

(Multiplicative) cohomology theory $E^*: X \mapsto E^*(X)$ a \mathbb{Z} -graded module (algebra). (contravariant functors)

Cohomology operation $Q_n: E^* \rightarrow E^{*+n}$. (natural transformations)

$$[X, Y] \xrightarrow{E^*} \mathbf{Maps}(E^* Y, E^* X)$$

graded modules (algebras)

How to process homotopy classes

(Multiplicative) cohomology theory $E^*: X \mapsto E^*(X)$ a \mathbb{Z} -graded module (algebra). (contravariant functors)

Cohomology operation $Q_n: E^* \rightarrow E^{*+n}$. (natural transformations)

The graded algebra of cohomology operations E^*E .

$$[X, Y] \xrightarrow{E^*} \mathbf{Maps}(E^*Y, E^*X)$$

graded modules (algebras)

How to process homotopy classes

(Multiplicative) cohomology theory $E^*: X \mapsto E^*(X)$ a \mathbb{Z} -graded module (algebra). (contravariant functors)

Cohomology operation $Q_n: E^* \rightarrow E^{*+n}$. (natural transformations)

The graded algebra of cohomology operations E^*E .

$$\begin{array}{ccc}
 [X, Y] & \xrightarrow{E^*} & \mathbf{Maps}(E^*Y, E^*X) \xrightarrow{\text{finer structure}} \mathbf{Maps}(E^*Y, E^*X) \\
 & \text{graded modules (algebras)} & \text{graded } E^*E\text{-modules} \\
 & & \text{compute it by homological methods!}
 \end{array}$$

Example: mod-2 ordinary cohomology theory

Let $H\mathbb{Z}/2$ be the mod-2 ordinary cohomology theory.

Theorem (Steenrod 1950s)

*There exists a **unique** sequence of cohomology operations Sq^i called **Steenrod squares** on $H\mathbb{Z}/2$ such that*

Example: mod-2 ordinary cohomology theory

Let $H\mathbb{Z}/2$ be the mod-2 ordinary cohomology theory.

Theorem (Steenrod 1950s)

*There exists a **unique** sequence of cohomology operations Sq^i called **Steenrod squares** on $H\mathbb{Z}/2$ such that*

- $Sq^i : H^n(X; \mathbb{Z}/2) \rightarrow H^{n+i}(X; \mathbb{Z}/2);$

Example: mod-2 ordinary cohomology theory

Let $H\mathbb{Z}/2$ be the mod-2 ordinary cohomology theory.

Theorem (Steenrod 1950s)

*There exists a **unique** sequence of cohomology operations Sq^i called **Steenrod squares** on $H\mathbb{Z}/2$ such that*

- $Sq^i : H^n(X; \mathbb{Z}/2) \rightarrow H^{n+i}(X; \mathbb{Z}/2);$
- $Sq^0 = \text{id}$ and Sq^1 is the mod-2 Bockstein operation;

Example: mod-2 ordinary cohomology theory

Let $H\mathbb{Z}/2$ be the mod-2 ordinary cohomology theory.

Theorem (Steenrod 1950s)

*There exists a **unique** sequence of cohomology operations Sq^i called **Steenrod squares** on $H\mathbb{Z}/2$ such that*

- $Sq^i: H^n(X; \mathbb{Z}/2) \rightarrow H^{n+i}(X; \mathbb{Z}/2);$
- $Sq^0 = \text{id}$ and Sq^1 is the mod-2 Bockstein operation;
- $Sq^i(u) = u^2$, if $i = \dim u$;

Example: mod-2 ordinary cohomology theory

Let $H\mathbb{Z}/2$ be the mod-2 ordinary cohomology theory.

Theorem (Steenrod 1950s)

*There exists a **unique** sequence of cohomology operations Sq^i called **Steenrod squares** on $H\mathbb{Z}/2$ such that*

- $Sq^i: H^n(X; \mathbb{Z}/2) \rightarrow H^{n+i}(X; \mathbb{Z}/2);$
- $Sq^0 = \text{id}$ and Sq^1 is the mod-2 Bockstein operation;
- $Sq^i(u) = u^2$, if $i = \dim u$;
- $Sq^i(u) = 0$, if $i > \dim u$;

Example: mod-2 ordinary cohomology theory

Let $H\mathbb{Z}/2$ be the mod-2 ordinary cohomology theory.

Theorem (Steenrod 1950s)

There exists a **unique** sequence of cohomology operations Sq^i called **Steenrod squares** on $H\mathbb{Z}/2$ such that

- $Sq^i: H^n(X; \mathbb{Z}/2) \rightarrow H^{n+i}(X; \mathbb{Z}/2);$
- $Sq^0 = \text{id}$ and Sq^1 is the mod-2 Bockstein operation;
- $Sq^i(u) = u^2$, if $i = \dim u$;
- $Sq^i(u) = 0$, if $i > \dim u$;
- **Cartan's formula:** $Sq^i(uv) = \sum_{j=0}^i Sq^j(u) \cdot Sq^{i-j}(v).$

The mod-2 Steenrod algebra

Let $\mathcal{A}_2^* := H\mathbb{Z}/2^* H\mathbb{Z}/2$ and it is called the **mod-2 Steenrod algebra**.

Theorem (Adem 1952)

$$Sq^a Sq^b = \sum_{j=0}^{[a/2]} \binom{b-1-j}{a-2j} Sq^{a+b-j} Sq^j, \text{ if } 0 < a < 2b.$$

The mod-2 Steenrod algebra

Let $\mathcal{A}_2^* := H\mathbb{Z}/2^* H\mathbb{Z}/2$ and it is called the **mod-2 Steenrod algebra**.

Theorem (Adem 1952)

$$Sq^a Sq^b = \sum_{j=0}^{[a/2]} \binom{b-1-j}{a-2j} Sq^{a+b-j} Sq^j, \text{ if } 0 < a < 2b.$$

Theorem (Serre 1953)

$\{Sq^I \mid \text{all 2-admissible sequences } I\}$ is a $\mathbb{Z}/2$ -basis of \mathcal{A}_2^ and Adem relations determines the all the relations.*

Example: mod- p ordinary cohomology theory for odd prime p

Theorem (Steenrod 1962)

*There exists a unique sequence of cohomology operations P_p^i called **mod- p power operations** on $H\mathbb{Z}/p$ such that*

Example: mod- p ordinary cohomology theory for odd prime p

Theorem (Steenrod 1962)

*There exists a unique sequence of cohomology operations P_p^i called **mod- p power operations** on $H\mathbb{Z}/p$ such that*

- $P_p^i: H^n(-; \mathbb{Z}/p) \rightarrow H^{n+2i(p-1)}(-; \mathbb{Z}/p);$

Example: mod- p ordinary cohomology theory for odd prime p

Theorem (Steenrod 1962)

*There exists a unique sequence of cohomology operations P_p^i called **mod- p power operations** on $H\mathbb{Z}/p$ such that*

- $P_p^i: H^n(-; \mathbb{Z}/p) \rightarrow H^{n+2i(p-1)}(-; \mathbb{Z}/p);$
- $P_p^0 = \text{id};$

Example: mod- p ordinary cohomology theory for odd prime p

Theorem (Steenrod 1962)

*There exists a unique sequence of cohomology operations P_p^i called **mod- p power operations** on $H\mathbb{Z}/p$ such that*

- $P_p^i: H^n(-; \mathbb{Z}/p) \rightarrow H^{n+2i(p-1)}(-; \mathbb{Z}/p);$
- $P_p^0 = \text{id};$
- $P_p^i(u) = u^p$ if $2i = \dim u;$

Example: mod- p ordinary cohomology theory for odd prime p

Theorem (Steenrod 1962)

*There exists a unique sequence of cohomology operations P_p^i called **mod- p power operations** on $H\mathbb{Z}/p$ such that*

- $P_p^i: H^n(-; \mathbb{Z}/p) \rightarrow H^{n+2i(p-1)}(-; \mathbb{Z}/p);$
- $P_p^0 = \text{id};$
- $P_p^i(u) = u^p$ if $2i = \dim u;$
- $P_p^i(u) = 0$, if $2i > \dim u;$

Example: mod- p ordinary cohomology theory for odd prime p

Theorem (Steenrod 1962)

*There exists a unique sequence of cohomology operations P_p^i called **mod- p power operations** on $H\mathbb{Z}/p$ such that*

- $P_p^i: H^n(-; \mathbb{Z}/p) \rightarrow H^{n+2i(p-1)}(-; \mathbb{Z}/p);$
- $P_p^0 = \text{id};$
- $P_p^i(u) = u^p$ if $2i = \dim u;$
- $P_p^i(u) = 0$, if $2i > \dim u;$
- **Cartan's formula:** $P_p^i(uv) = \sum_{j+k=i} P_p^j(u)P^k(v).$

Adem relations in mod- p ordinary cohomology theory

Let β be the mod- p Bockstein operations. If $a < pb$, then

$$P_p^a P_p^b = \sum_{j=0}^{[a/p]} \binom{(p-1)(b-j)-1}{a-pj} P_p^{a+b-j} P_p^j$$

if $a \leq b$, then

$$\begin{aligned} P_p^a \beta P_p^b &= \sum_{j=0}^{[a/p]} \binom{(p-1)(b-j)-1}{a-pj} \beta P_p^{a+b-j} P_p^j \\ &\quad + \sum_{j=0}^{[(a-1)/p]} (-1)^{a+j-1} \binom{(p-1)(b-j)-1}{a-pj-1} \beta P_p^{a+b-j} P_p^j \end{aligned}$$

The mod- p Steenrod algebra

The mod- p Steenrod operation St_p^i is defined as

$$St_p^i = \begin{cases} P_p^k, & i = 2k(p-1) \\ \beta P_p^k, & i = 2k(p-1) + 1 \\ 0, & \text{otherwise.} \end{cases}$$

Theorem (Cartan-Serre 1950s)

$\{St_p^I \mid \text{all } p\text{-admissible sequences } I\}$ is a \mathbb{Z}/p -basis of \mathcal{A}_p^* and Adem relations determines all the relations.

Applications of the Steenrod operations

Theorem (Borel-Serre 1953)

If $n > 3$, then S^{2n} does not admit an almost complex structure.

Applications of the Steenrod operations

Theorem (Borel-Serre 1953)

If $n > 3$, then S^{2n} does not admit an almost complex structure.

Theorem (Thom 1954)

Any mod-2 homology class of a finite complex K can be realized as a manifold. For any integral homology class y of K , there exists N such that Ny can be realized as an oriented manifold.

The classical Adams spectral sequences

Theorem (Adams 1958)

Given spaces or spectra X and Y , there exists a cohomological spectral sequence $\{E_*^{*,*}\}$ called **Adams spectral sequence** such that

$$E_2^{s,t} = \text{Ext}_{\mathcal{A}_p^*}^{s,t}(H\mathbb{Z}/p^*Y, H\mathbb{Z}/p^*X) \Rightarrow ([X, Y]_{t-s})_p^\wedge$$

where $([X, Y]_{t-s})_p^\wedge$ is the p -completion of the group of stable homotopy classes $\text{colim}_n[\Sigma^{n+t-s}X, \Sigma^n Y]$.

If we let X, Y be points, then it converges to the p -completion of the stable homotopy group of spheres.

How Steenrod constructed mod- p power operations

$$\begin{array}{ccc}
 H^n(X) & \xrightarrow{\mathcal{P}^d} & H_{\Sigma_d}^{nd}(X^d) \xrightarrow{\Delta^*} H^{nd}(B\Sigma_d \times X) \\
 [u] & & [u^d]_{\Sigma_d} \\
 n\text{-cocycle class} & & \Sigma_d\text{-equivariant } nd\text{-cocycle class}
 \end{array}$$

where \mathcal{P}^d is called **the d -external power operation** and $\Delta^*\mathcal{P}^d$ is called **the d -total power operation**.

How Steenrod constructed mod- p power operations

$$\begin{array}{ccc}
 H^n(X) & \xrightarrow{\mathcal{P}^d} & H_{\Sigma_d}^{nd}(X^d) \xrightarrow{\Delta^*} H^{nd}(B\Sigma_d \times X) \\
 [u] & & [u^d]_{\Sigma_d} \\
 n\text{-cocycle class} & & \Sigma_d\text{-equivariant } nd\text{-cocycle class}
 \end{array}$$

where \mathcal{P}^d is called **the d -external power operation** and $\Delta^*\mathcal{P}^d$ is called **the d -total power operation**.

How Steenrod constructed mod- p power operations

$$\begin{array}{ccc}
 H^n(X) & \xrightarrow{\mathcal{P}^d} & H_{\Sigma_d}^{nd}(X^d) \xrightarrow{\Delta^*} H^{nd}(B\Sigma_d \times X) \\
 [u] & & [u^d]_{\Sigma_d} \\
 n\text{-cocycle class} & & \Sigma_d\text{-equivariant } nd\text{-cocycle class}
 \end{array}$$

where \mathcal{P}^d is called **the d -external power operation** and $\Delta^*\mathcal{P}^d$ is called **the d -total power operation**.

Given $\alpha \in H_i(B\Sigma_d)$, then we have the cohomology operation derived from α is $[u] \mapsto \Delta^*\mathcal{P}^d([u]) \cap \alpha \in H^{nd-i}(X)$.

How Steenrod constructed mod- p power operations

$$\begin{array}{ccc}
 H^n(X) & \xrightarrow{\mathcal{P}^d} & H_{\Sigma_d}^{nd}(X^d) \xrightarrow{\Delta^*} H^{nd}(B\Sigma_d \times X) \\
 [u] & & [u^d]_{\Sigma_d} \\
 n\text{-cocycle class} & & \Sigma_d\text{-equivariant } nd\text{-cocycle class}
 \end{array}$$

where \mathcal{P}^d is called **the d -external power operation** and $\Delta^*\mathcal{P}^d$ is called **the d -total power operation**.

Given $\alpha \in H_i(B\Sigma_d)$, then we have the cohomology operation derived from α is $[u] \mapsto \Delta^*\mathcal{P}^d([u]) \cap \alpha \in H^{nd-i}(X)$.

If we replace Σ_d by \mathbb{Z}/p and let $i = p$, then we get mod- p power operations.

Outline

- 1 Background
- 2 Power operations in topology
- 3 Power operations in algebraic geometry
- 4 Questions for further investigation

Generalized cohomology theories and spectra

Definition (Spectra)

A spectrum $E = \{E_n, \varepsilon_n\}_{n \in \mathbb{Z}}$ is a sequence of pointed topological spaces E_n with basepoint-preserving maps $\varepsilon_n: \Sigma E_n \rightarrow E_{n+1}$. If $\varepsilon_n: E_n \rightarrow \Omega E_{n+1}$ is a weak homotopy equivalence, it is called an Ω -spectrum.

Generalized cohomology theories and spectra

Definition (Spectra)

A spectrum $E = \{E_n, \varepsilon_n\}_{n \in \mathbb{Z}}$ is a sequence of pointed topological spaces E_n with basepoint-preserving maps $\varepsilon_n: \Sigma E_n \rightarrow E_{n+1}$. If $\varepsilon_n: E_n \rightarrow \Omega E_{n+1}$ is a weak homotopy equivalence, it is called an Ω -spectrum.

Theorem (Brown 1962)

Each generalized cohomology theory h^* is represented by an Ω -spectrum E_n such that $h^n(X) \cong [X, E_n]$.

Generalized cohomology theories and spectra

Definition (Spectra)

A spectrum $E = \{E_n, \varepsilon_n\}_{n \in \mathbb{Z}}$ is a sequence of pointed topological spaces E_n with basepoint-preserving maps $\varepsilon_n: \Sigma E_n \rightarrow E_{n+1}$. If $\varepsilon_n: E_n \rightarrow \Omega E_{n+1}$ is a weak homotopy equivalence, it is called an Ω -spectrum.

Theorem (Brown 1962)

Each generalized cohomology theory h^* is represented by an Ω -spectrum E_n such that $h^n(X) \cong [X, E_n]$.

Example

$H^n(X; A) = [X, K(A, n)]$. In particular, $(H\mathbb{Z}/p)_n := K(\mathbb{Z}/p, n)$.

Generalized cohomology theories and spectra

Definition

A morphism $f: E \rightarrow F$ between spectra consists of $\{f_n: E_n \rightarrow F_n\}$ compatible with Σ and ε_n .

Given a based space X and a spectrum E , $(E \wedge X)_n := E_n \wedge X$.

We say $f \simeq g: E \rightarrow F$ if there exists a map $h: E \wedge I_+ \rightarrow F$ such that $f = h_0$ and $g = h_1$.

the stable homotopy classes: $[E, F]^n := [E, \Sigma^n F]$.

the associated generalized cohomology: $E^*(X) := [\Sigma^\infty X, E]^*$.

the associated generalized homology: $E_*(X) := [\Sigma^\infty S^0, E \wedge X]^*$.

The stable homotopy categories

The essence is “inverting” S^1 with respect to \wedge by stabilizing it.

Theorem

There exists a closed symmetric monoidal category of spectra such that the sphere spectrum \mathbb{S} is a unit.

The stable homotopy categories

The essence is “inverting” S^1 with respect to \wedge by stabilizing it.

Theorem

There exists a closed symmetric monoidal category of spectra such that the sphere spectrum \mathbb{S} is a unit.

The construction of such categories is very complicated!

The stable homotopy categories

The essence is “inverting” S^1 with respect to \wedge by stabilizing it.

Theorem

There exists a closed symmetric monoidal category of spectra such that the sphere spectrum \mathbb{S} is a unit.

The construction of such categories is very complicated!

There are three popular constructions, we choose the category of \mathbb{S} -modules (EKMM) in this presentation.

The algebra of cohomology operations

Proposition

*By Yoneda lemma and Brown's representability theorem, the algebra of cohomology operations on E is $E^*E := [E, E]^*$, the stable homotopy classes from E^* to itself.*

In particular, $\mathcal{A}_p^* = H\mathbb{Z}/p^* H\mathbb{Z}/p$.

The algebra of cohomology operations

Proposition

*By Yoneda lemma and Brown's representability theorem, the algebra of cohomology operations on E is $E^*E := [E, E]^*$, the stable homotopy classes from E^* to itself.*

In particular, $\mathcal{A}_p^* = H\mathbb{Z}/p^* H\mathbb{Z}/p$.

Question

Given a ring spectrum E , how to determine power operations on E ?

Extended powers and H_∞ -structures

Given an \mathbb{S} -module E , the j th **extended power** of E is defined to be $D_j E = (E \Sigma_j)_+ \wedge E^j / \Sigma_j$.

Definition

An H_∞ -ring spectrum is a \mathbb{S} -module M together with $\xi_j: D_j M \rightarrow M$ for $j \geq 0$ satisfying some homotopy coherence conditions.

Extended powers and H_∞ -structures

Given an \mathbb{S} -module E , the j th **extended power** of E is defined to be $D_j E = (E \Sigma_j)_+ \wedge E^j / \Sigma_j$.

Definition

An H_∞ -ring spectrum is a \mathbb{S} -module M together with $\xi_j: D_j M \rightarrow M$ for $j \geq 0$ satisfying some homotopy coherence conditions.

Example

HR , KU and MU are H_∞ -ring spectra.

H_∞ -structures give rise to power operations

Let E be an H_∞ -ring spectrum.

$$\begin{array}{ccccccc}
 E^*(X) & \xrightarrow{\quad \mathcal{P}_j \quad} & & & E^*(B\Sigma_{j+} \wedge X) \\
 \cong \downarrow & & & & \downarrow \cong \\
 [\Sigma^\infty X, E] & \xrightarrow{D_j} [D_j(\Sigma^\infty X), D_j E] & \xrightarrow{\circ \xi_j} & [\Sigma^\infty(D_j X), E] & \xrightarrow{\Delta^*} & [\Sigma^\infty(B\Sigma_{j+} \wedge X), E]
 \end{array}$$

Then we can derive power operations from $E_*(B\Sigma_j)$.

The generalized Adams spectral sequences

Theorem (Adams spectral sequences)

Given spaces or spectra X and Y and a cohomology theory E^ , there exists a cohomological spectral sequence $\{E_*^{*,*}\}$ such that*

$$E_2^{s,t} = \text{Ext}_{E^*E}^{s,t}(E^*Y, E^*X) \Rightarrow [X, Y]_{t-s}^E$$

where $[X, Y]_{t-s}^E$ is the set of stable homotopy classes from X to Y in an E -localization shifting $t - s$.

If E is an H_∞ -ring spectra, then the induced power operations appear in the E_2 -page.

Outline

- 1 Background
- 2 Power operations in topology
- 3 Power operations in algebraic geometry**
- 4 Questions for further investigation

The construction of motivic homotopy theory

Construction (Morel-Voevodsky 1990s)

Let S be a qcqs Noetherian scheme of finite dimension and let Sm/S be the category of smooth schemes of finite type over S . Let $\Delta^{op}\mathbf{Shv}_{Nis}(\mathrm{Sm}/S)$ be the category of **Nisnevich sheaves** of simplicial sets with **projective model structure**. The unstable motivic homotopy category is

$$\mathcal{H}(S) := L_{\mathbb{A}^1}\Delta^{op}\mathbf{Shv}_{Nis}(\mathrm{Sm}/S)$$

where $L_{\mathbb{A}^1}$ is the **Bousfield localization with respect to the class generated by natural projections** $X \times_S \mathbb{A}^1 \rightarrow X$ for all $X \in \mathrm{Sm}/S$.

Spheres in motivic homotopy category

Definition (Spheres in motivic homotopy category)

Simplicial circle S_s^1 (or denote it $S^{1,0}$): the constant sheaf valued at the $\Delta^1/\partial\Delta^1$.

Tate circle S_t^1 (or denote it $S^{1,1}$): the sheaf represented by \mathbb{G}_m .
Given a, b two non-negative integers with $a \geq b$, the **bigraded motivic sphere** $S^{a,b} := (S_t^1)^{\wedge b} \wedge (S_s^1)^{\wedge a-b}$.

Spheres in motivic homotopy category

Definition (Spheres in motivic homotopy category)

Simplicial circle S_s^1 (or denote it $S^{1,0}$): the constant sheaf valued at the $\Delta^1/\partial\Delta^1$.

Tate circle S_t^1 (or denote it $S^{1,1}$): the sheaf represented by \mathbb{G}_m .
Given a, b two non-negative integers with $a \geq b$, the **bigraded motivic sphere** $S^{a,b} := (S_t^1)^{\wedge b} \wedge (S_s^1)^{\wedge a-b}$.

Proposition

$$S^{2n,n} \simeq \mathbb{P}^n / \mathbb{P}^{n-1} \simeq \mathbb{A}^n / (\mathbb{A}^n - 0)$$

The motivic stable homotopy category

Construction

Recall that we obtain classical stable homotopy category by "inverting" the circle S^1 from $\mathbf{h}(\mathcal{S}\text{paces})$, we obtain motivic stable homotopy category $\mathcal{SH}(S)$ over S by "inverting" $\mathbb{P}^1 \simeq S_t^1 \wedge S_s^1$ from $\mathcal{H}(S)$, whose objects are called motivic spectra.

The motivic stable homotopy category

Construction

Recall that we obtain classical stable homotopy category by "inverting" the circle S^1 from $\mathbf{h}(\mathcal{S}\text{paces})$, we obtain motivic stable homotopy category $\mathcal{SH}(S)$ over S by "inverting" $\mathbb{P}^1 \simeq S_t^1 \wedge S_s^1$ from $\mathcal{H}(S)$, whose objects are called motivic spectra.

cohomology theory	classical spectrum	motivic spectrum
singular cohomology	$H\mathbb{Z}$	$H\mathbb{Z}_{\text{mot}}$
K-theory	KU	KGL
cobordism theory	MU	MGL

Table: Cohomology theories and spectra in classical setting and motivic setting

How motivic homotopy theory captures arithmetic data

Theorem (Morel 2004)

If k is a perfect field (with $\text{char } k \neq 2$), then we have an isomorphism between graded rings

$$K_*^{MW}(k) \cong [S^0, S_t^1]_{\mathbb{P}^1}$$

How motivic homotopy theory captures arithmetic data

Theorem (Morel 2004)

If k is a perfect field (with $\text{char } k \neq 2$), then we have an isomorphism between graded rings

$$K_*^{MW}(k) \cong [S^0, S_t^1]_{\mathbb{P}^1}$$

Theorem (Morel 2004)

If k is a perfect field (with $\text{char } k \neq 2$), then we have an isomorphism between rings

$$GW(k) \cong [S^0, S^0]_{\mathbb{P}^1}$$

The motivic Steenrod operations

Theorem (Voevodsky 2003)

There exists $P_\ell^i: H^{,*}(X; \mathbb{Z}/\ell) \rightarrow H^{*+2i(\ell-1), *+i(\ell-1)}(X; \mathbb{Z}/\ell)$ and $B_\ell^i: H^{*,*}(X; \mathbb{Z}/\ell) \rightarrow H^{*+2i(\ell-1)+1, *+i(\ell-1)}(X; \mathbb{Z}/\ell)$ such that*

The motivic Steenrod operations

Theorem (Voevodsky 2003)

There exists $P_\ell^i: H^{*,*}(X; \mathbb{Z}/\ell) \rightarrow H^{*+2i(\ell-1), *+i(\ell-1)}(X; \mathbb{Z}/\ell)$ and $B_\ell^i: H^{*,*}(X; \mathbb{Z}/\ell) \rightarrow H^{*+2i(\ell-1)+1, *+i(\ell-1)}(X; \mathbb{Z}/\ell)$ such that

1 $P_\ell^0 = \text{id}$ and $P_\ell^n(u) = u^n$ if $u \in H^{2n, n}$;

The motivic Steenrod operations

Theorem (Voevodsky 2003)

There exists $P_\ell^i: H^{*,*}(X; \mathbb{Z}/\ell) \rightarrow H^{*+2i(\ell-1), *+i(\ell-1)}(X; \mathbb{Z}/\ell)$ and $B_\ell^i: H^{*,*}(X; \mathbb{Z}/\ell) \rightarrow H^{*+2i(\ell-1)+1, *+i(\ell-1)}(X; \mathbb{Z}/\ell)$ such that

1 $P_\ell^0 = \text{id}$ and $P_\ell^n(u) = u^n$ if $u \in H^{2n,n}$;

2 **Cartan formula:** if $\ell \neq 2$,

$$P_\ell^i(uv) = \sum_{j=0}^i P_\ell^j(u) P_\ell^{i-j}(v)$$

$$B_\ell^i(uv) = \sum_{j=0}^i B_\ell^j(u) P_\ell^{i-j}(v) + (-1)^{\deg(u)} P_\ell^j(u) B_\ell^{i-j}(v)$$

The motivic Steenrod operations

Theorem (Voevodsky 2003)

If $\ell = 2$, let $Sq^{2i} = P_2^i$, $Sq^{2i+1} = B_2^i$, τ be the generator of $H^{0,1}(K; \mathbb{Z}/2)$, and $\rho \in H^{1,1}(k; \mathbb{Z}/2)$ be the class of -1 , then

$$Sq^{2i}(uv) = \sum_{j=0}^i Sq^{2j}(u)Sq^{2i-2j}(v) + \tau \sum_{s=0}^{i-1} Sq^{2s+1}(u)Sq^{2i-2s-1}(v)$$

$$\begin{aligned} Sq^{2i+1}(uv) &= \sum_{j=0}^i (Sq^{2j+1}(u)Sq^{2i-2j}(v) + Sq^{2j}(u)Sq^{2i-2j-1}(v)) \\ &\quad + \rho \sum_{s=0}^{i-1} Sq^{2s+1}(u)Sq^{2i-2s-1}(v) \end{aligned}$$

The Milnor conjecture and the Bloch-Kato conjecture

Voevodsky used motivic Steenrod operations to prove the following two theorems:

Theorem (Milnor conjecture, Voevodsky 2003)

Let k be a field of characteristic not equal to 2, then the norm residue homomorphisms $K_n^M(k)/2 \rightarrow H_{\text{ét}}^n(k; \mathbb{Z}/2)$ are isomorphisms for all $n \geq 0$.

The Milnor conjecture and the Bloch-Kato conjecture

Voevodsky used motivic Steenrod operations to prove the following two theorems:

Theorem (Milnor conjecture, Voevodsky 2003)

Let k be a field of characteristic not equal to 2, then the norm residue homomorphisms $K_n^M(k)/2 \rightarrow H_{\text{ét}}^n(k; \mathbb{Z}/2)$ are isomorphisms for all $n \geq 0$.

Theorem (Bloch-Kato conjecture, Voevodsky 2010)

Let k be a field of characteristic not equal to a prime ℓ , then the norm residue homomorphisms $K_n^M(k)/\ell \rightarrow H_{\text{ét}}^n(k; \mathbb{Z}/\ell)$ are isomorphisms for all $n \geq 0$.

The motivic Steenrod algebras

Theorem (Voevodsky 2003, Voevodsky 2011)

Let k be field and ℓ be a prime coprime to $\text{char}(k)$, and k contains a primitive ℓ th root of unity. Then the motivic cohomology

$$\mathbb{M}_\ell := H^{*,*}(k; \mathbb{Z}/\ell) \cong \frac{K_*^M(k)}{\ell}[\tau]$$

where $K_^M(k)/\ell$ has degree (n, n) and τ is of degree $(0, 1)$.*

The motivic Steenrod algebras

Theorem (Voevodsky 2003, Voevodsky 2011)

Let k be field and ℓ be a prime coprime to $\text{char}(k)$, and k contains a primitive ℓ th root of unity. Then the motivic cohomology

$$\mathbb{M}_\ell := H^{*,*}(k; \mathbb{Z}/\ell) \cong \frac{K_*^M(k)}{\ell}[\tau]$$

where $K_^M(k)/\ell$ has degree (n, n) and τ is of degree $(0, 1)$.*

Theorem (Voevodsky 2003)

The bigraded motivic Steenrod algebra $\mathcal{A}_\ell^{,*}$ on mod- ℓ motivic cohomology is generated by P_ℓ^i and B_ℓ^i over \mathbb{M}_ℓ and is characterized by motivic Adem relations.*

The motivic Adams spectral sequences

Theorem (Dugger-Isaksen 2010, Hu-Kriz-Ormsby 2011, Kylling-Wilson 2019)

Let k be a field of characteristic not equal to a prime ℓ , let $\mathbb{M}_\ell := H^{,*}(k; \mathbb{Z}/\ell)$, there is spectral sequence called **motivic Adams spectral sequence** such that*


$$E_2 = \mathrm{Ext}_{\mathcal{A}_\ell^{*,*}}(\mathbb{M}_\ell, \mathbb{M}_\ell) \Rightarrow [\Sigma_{s,t}^\infty \mathrm{Spec}(k), \Sigma_{s,t}^\infty \mathrm{Spec}(k)]_{*,*}^{\mathbb{A}_k^1}$$

Outline

- 1 Background
- 2 Power operations in topology
- 3 Power operations in algebraic geometry
- 4 Questions for further investigation**


How the Adams spectral sequences detect information

We summarize Bruner's mechanism in the following diagram.




How the Adams spectral sequences detect information


We summarize Bruner's mechanism in the following diagram.




What hides behind the power operations




What hides behind the power operations



What hides behind the power operations



How motivic extended powers emerge in the motivic Adams spectral sequences



Question & Answer

Thank you!