

# Introduction to linear regression

Gabriel Campos

The Human Freedom Index is a report that attempts to summarize the idea of “freedom” through a bunch of different variables for many countries around the globe. It serves as a rough objective measure for the relationships between the different types of freedom - whether it’s political, religious, economical or personal freedom - and other social and economic circumstances. The Human Freedom Index is an annually co-published report by the Cato Institute, the Fraser Institute, and the Liberales Institut at the Friedrich Naumann Foundation for Freedom.

In this lab, you’ll be analyzing data from Human Freedom Index reports from 2008-2016. Your aim will be to summarize a few of the relationships within the data both graphically and numerically in order to find which variables can help tell a story about freedom.

## The data

The data we’re working with is in the `openintro` package and it’s called ‘`hfi`’, short for Human Freedom Index.

1.

What are the dimensions of the dataset?

**ANSWER:** *Rows : 1,458 Columns : 123*

```
glimpse(hfi)
```


2.

What type of plot would you use to display the relationship between the personal freedom score, 'pf\_score', and one of the other numerical variables? Plot this relationship using the variable 'pf\_expression\_control' as the predictor. Does the relationship look linear? If you knew a country's 'pf\_expression\_control', or its score out of 10, with 0 being the most, of political pressures and controls on media content, would you be comfortable using a linear model to predict the personal freedom score?

**ANSWER:** I would use a scatter plot, which shows there is some linearity to the points. I do not feel comfortable using a linear model for predicting the personal freedom score, the data is too scattered with outliers, and I would be unsure of what type of plot to use.


### Scatterplot

Personal freedom score and personal freedom expression control (Linear Model 'lm')



### Scatterplot

Personal freedom score and personal freedom expression control (Local Estimated Scatterplot Smoothing LEOSS)



If the relationship looks linear, we can quantify the strength of the relationship with the correlation coefficient.

```
## # A tibble: 1 x 1
##   'cor(pf_expression_control, pf_score, use = "complete.obs")'
##                                                                    <dbl>
## 1                                                                    0.796
```

Here, we set the `use` argument to “complete.obs” since there are some observations of NA.

## Sum of squared residuals

In this section, you will use an interactive function to investigate what we mean by “sum of squared residuals”. You will need to run this function in your console, not in your markdown document. Running the function also requires that the `hfi` dataset is loaded in your environment.

Think back to the way that we described the distribution of a single variable. Recall that we discussed characteristics such as center, spread, and shape. It’s also useful to be able to describe the relationship of two numerical variables, such as `pf_expression_control` and `pf_score` above.

### 3.


Looking at your plot from the previous exercise, describe the relationship between these two variables. Make sure to discuss the form, direction, and strength of the relationship as well as any unusual observations.

**ANSWER:** From the bottom left to the top right, it’s clear to see a positive correlation between the two variables visually and the correlation coefficient of 0.796 supports the relationship is a strong one. There are however a great deal of plots, including outliers to consider regarding the data.

Just as you've used the mean and standard deviation to summarize a single variable, you can summarize the relationship between these two variables by finding the line that best follows their association. Use the following interactive function to select the line that you think does the best job of going through the cloud of points.

```
#this isn't working, wondering if we should just remove rows with NA  
plot_ss(x = pf_expression_control, y = pf_score, data = hfi)
```

### Plot\_ss with NA removed



After running this command, you'll be prompted to click two points on the plot to define a line. Once you've done that, the line you specified will be shown in black and the residuals in blue. Note that there are 30 residuals, one for each of the 30 observations. Recall that the residuals are the difference between the observed values and the values predicted by the line:

$$e_i = y_i - \hat{y}_i$$

The most common way to do linear regression is to select the line that minimizes the sum of squared residuals. To visualize the squared residuals, you can rerun the plot command and add the argument `showSquares = TRUE`.

```
plot_ss(x = pf_expression_control, y = pf_score, data = 'hfi', showSquares = TRUE)
```


Note that the output from the `plot_ss` function provides you with the slope and intercept of your line as well as the sum of squares.

4.

Using 'plot\_ss', choose a line that does a good job of minimizing the sum of squares. Run the function several times. What was the smallest sum of squares that you got? How does it compare to your neighbors?

**Stefano Biguzzi 9:50 AM**

*For lab 8 question 4 it asks to compare your answer to your neighbors. The smallest Sum of Least squares I got was 989.955*



```
## Click two points to make a line.
## Call:
## lm(formula = y ~ x, data = pts)
##
## Coefficients:
## (Intercept)          x
##      4.6171      0.4914
##
## Sum of Squares:  952.153
```

## The linear model

It is rather cumbersome to try to get the correct least squares line, i.e. the line that minimizes the sum of squared residuals, through trial and error. Instead, you can use the `lm` function in R to fit the linear model (a.k.a. regression line).

```
m1 <- lm(pf_score ~ pf_expression_control, data = hfi)
```

The first argument in the function `lm` is a formula that takes the form `y ~ x`. Here it can be read that we want to make a linear model of `pf_score` as a function of `pf_expression_control`. The second argument specifies that R should look in the `hfi` data frame to find the two variables.

The output of `lm` is an object that contains all of the information we need about the linear model that was just fit. We can access this information using the summary function.

```
summary(m1)
```

Let's consider this output piece by piece. First, the formula used to describe the model is shown at the top. After the formula you find the five-number summary of the residuals. The "Coefficients" table shown next is key; its first column displays the linear model's y-intercept and the coefficient of `pf_score`. With this table, we can write down the least squares regression line for the linear model:

$$\hat{y} = 4.61707 + 0.49143 \times pf\_expression\_control$$

One last piece of information we will discuss from the summary output is the Multiple R-squared, or more simply,  $R^2$ . The  $R^2$  value represents the proportion of variability in the response variable that is explained by the explanatory variable. For this model, 63.42% of the variability in runs is explained by 'pf\_expression\_control'.

5.

Fit a new model that uses 'pf\_expression\_control' to predict 'hf\_score', or the total human freedom score. Using the estimates from the R output, write the equation of the regression line. What does the slope tell us in the context of the relationship between human freedom and the amount of political pressure on media content?

**ANSWER:** The slope indicates a positive relationship with a small incremental increase or mild steepness.


$$\hat{y} = 5.153687 + 0.349862 \times pf\_expression\_control$$

```
m5 <- lm(hf_score ~ pf_expression_control, data = hfi)
summary(m5)
```

## Prediction and prediction errors

Let's create a scatterplot with the least squares line for m1 laid on top.

```
ggplot(data = hfi, aes(x = pf_expression_control, y = pf_score)) +
  geom_point() +
  stat_smooth(method = "lm", se = FALSE)
```



Here, we are literally adding a layer on top of our plot. `geom_smooth` creates the line by fitting a linear model. It can also show us the standard error `se` associated with our line, but we'll suppress that for now.

This line can be used to predict  $y$  at any value of  $x$ . When predictions are made for values of  $x$  that are beyond the range of the observed data, it is referred to as *extrapolation* and is not usually recommended. However, predictions made within the range of the data are more reliable. They're also used to compute the residuals.

6.

If someone saw the least squares regression line and not the actual data, how would they predict a country's personal freedom school for one with a 6.7 rating for 'pf\_expression\_control'? Is this an overestimate or an underestimate, and by how much? In other words, what is the residual for this prediction?

**ANSWER:** You could substitute  $\hat{y}$  with 6.7 which is a relatively close result since the pf\_scores at pf\_expression\_control 7.5~ is between 7-8.5

$$\hat{y} = 5.153687 + 0.349862 \times 6.7$$


$$\hat{y} = 7.497762$$

## Model diagnostics

To assess whether the linear model is reliable, we need to check for (1) linearity, (2) nearly normal residuals, and (3) constant variability.

**Linearity:** You already checked if the relationship between `pf_score` and 'pf\_expression\_control' is linear using a scatterplot. We should also verify this condition with a plot of the residuals vs. fitted (predicted) values.

```
ggplot(data = m1, aes(x = .fitted, y = .resid)) +  
  geom_point() +  
  geom_hline(yintercept = 0, linetype = "dashed") +  
  xlab("Fitted values") +  
  ylab("Residuals")
```




Notice here that `m1` can also serve as a data set because stored within it are the fitted values ( $\hat{y}$ ) and the residuals. Also note that we're getting fancy with the code here. After creating the scatterplot on the first layer (first line of code), we overlay a horizontal dashed line at  $y = 0$  (to help us check whether residuals are distributed around 0), and we also rename the axis labels to be more informative.

7.

Is there any apparent pattern in the residuals plot? What does this indicate about the linearity of the relationship between the two variables?


**ANSWER:** No real pattern outside of appearing to be nearly normally distributed or possibly linear

**Nearly normal residuals:** To check this condition, we can look at a histogram



or a normal probability plot of the residuals.





Note that the syntax for making a normal probability plot is a bit different than what you're used to seeing: we set `sample` equal to the residuals instead of `x`, and we set a statistical method `qq`, which stands for “quantile-quantile”, another name commonly used for normal probability plots.

8.

Based on the histogram and the normal probability plot, does the nearly normal residuals condition appear to be met?

**ANSWER:** Yes it appears nearly normal

**Constant variability:**

9.

Based on the residuals vs. fitted plot, does the constant variability condition appear to be met?

**ANSWER:** Yes constant variability condition appears to be met.

## More Practice


10.

Choose another freedom variable and a variable you think would strongly correlate with it.. Produce a scatterplot of the two variables and fit a linear model. At a glance, does there seem to be a linear relationship?


**ANSWER:** New Variable is pf\_movements. Their appears to be a somewhat linear and positive relationship.

### Scatterplot

Personal freedom score and personal freedom movement



Source: Human Freedom Index



11

How does this relationship compare to the relationship between 'pf\_expression\_control' and 'pf\_score'? Use the  $R^2$  values from the two

model summaries to compare. Does your independent variable seem to predict your dependent one better? Why or why not?

**ANSWER:**  $R^2$ -value for my 'movement' model is 61.02% compared to my original percentage of 63.42% which is a slight decrease making the original model better since it will be accounting for a larger variation.

```
hfi%>%  
summarise(cor(pf_movement, pf_movement, use = "complete.obs"))  
summary(m10)  
summary(m1)
```

12

What's one freedom relationship you were most surprised about and why? Display the model diagnostics for the regression model analyzing this relationship.

**ANSWER:** I was not surprised the personal freedom movement was as closely tied to personal freedom control as it was, I feel those two variables are like seeing different sides of the same coin.

---

This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.