

Lecture 3:

Local Descent

Optimization T2023

Màster de Fonaments de Ciència de Dades



UNIVERSITAT_{DE}
BARCELONA

$$f(\mathbf{x}) \rightarrow \min, \quad \mathbf{x} \in D \subseteq \mathbb{R}^n, \quad n \geq 1, \quad f \text{ is smooth}$$

Goal: Iteratively find a sequence $\mathbf{x}^{(1)}, \mathbf{x}^{(2)}, \dots \rightarrow \mathbf{x}^*$,
where \mathbf{x}^* is a solution of the optimization problem
(local or global minimum), realizing the descent

$$f(\mathbf{x}^{(1)}) > f(\mathbf{x}^{(2)}) > \dots$$

(for all or most* of the iterates)

Recall that
 $\nabla f(\mathbf{x}^*) = 0$

General descent method.

given a starting point $\mathbf{x}^{(1)} \in D$

repeat

1. Determine *descent direction* $\mathbf{p}^{(k)}$ (often, $\|\mathbf{p}^{(k)}\| = 1$)
2. Determine *step size/learning rate* $\alpha^{(k)}$
3. Update $\mathbf{x}^{(k+1)} := \mathbf{x}^{(k)} + \alpha^{(k)} \mathbf{p}^{(k)}$

until *stopping criterion* is satisfied

III. Descent direction?

II. Step size?

I. Stopping criterion?

Digression: Why gradient?

Recall that from the Taylor formula

$$\begin{aligned} f(\mathbf{x} + \mathbf{v}) &= \text{up to H.O.T. } f(\mathbf{x}) + \mathbf{v}^T \cdot \nabla f(\mathbf{x}) \\ &= f(\mathbf{x}) + \nabla_{\mathbf{v}} f(\mathbf{x}) \end{aligned}$$

Directional
derivative



We want it to
be negative

Theorem:

Let $f: D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ be a differentiable function, $\mathbf{a} \in D$, $\mathbf{d} \in \mathbb{R}^n$ with $\|\mathbf{d}\| = 1$. If θ is the angle between \mathbf{d} and $\nabla f(\mathbf{a})$. Then

$$\nabla_{\mathbf{d}} f(\mathbf{a}) = \mathbf{d}^T \cdot \nabla f(\mathbf{a}) = \|\nabla f(\mathbf{a})\| \cos \theta$$

In particular, the vector $-\nabla f(\mathbf{a})$ gives the maximum descent direction of f at the point \mathbf{a} .



I. Stopping criteria/termination conditions

- **Maximum iterations:** repeat until $k \leq k_{max}$

- **Absolute improvement:** repeat until

$$f(\mathbf{x}^{(k)}) - f(\mathbf{x}^{(k+1)}) < \epsilon_a$$

- **Relative improvement:** repeat until

$$f(\mathbf{x}^{(k)}) - f(\mathbf{x}^{(k+1)}) < \epsilon_r |f(\mathbf{x}^{(k)})|$$

- **Gradient magnitude:** repeat until

$$\|\nabla f(\mathbf{x}^{(k+1)})\| < \epsilon_g$$

-
- ✓ One or more termination conditions can be used
 - ✓ If there are several local minima, one can add **random restart** with $\mathbf{x}^{(1),new}$ sampled randomly from D

II. Step size/learning rate

Suppose $x = x^{(k)}$ and $p = p^{(k)}$ is given. How to find $\alpha = \alpha^{(k)}$?

Methods:

1. Exact line search
2. Approximate line search
3. Trust region methods

Exact line search

$$\text{minimize}_{\alpha} f(\mathbf{x} + \alpha \mathbf{p})$$

- This is univariate optimization problem for $\phi(\alpha) := f(\mathbf{x} + \alpha \mathbf{p}) \rightarrow$


→ Find a **bracket** for the optimal solution α^*

(α^* is characterized by $\phi(\alpha^*) < \phi(\alpha)$ for all α near α^*)

→ Use univariate optimization methods to find an approximate of α^* by successively shrinking the bracket. Methods include:

- Dyadic/binary search
- Fibonacci search
- Quadratic fit search
- Shubert–Piyavskii method
- Bisection method

Definition: A *bracket* is an interval $[\alpha', \alpha'']$ containing α^*







Digression: some univalent optimization methods

[KW, Ch.3]


- **Dyadic/binary search**: subdivide interval 'in half' at each step
- **Fibonacci search**: max reduction of interval size for given number of function evaluations
- **Quadratic fit search**
- **Shubert-Piyavskii method**: assuming ϕ is Lipschitz, e.g.

$$|\phi(x) - \phi(y)| \leq \ell \cdot |x - y|, \quad \forall x, y \in [\alpha', \alpha'']$$

- **Bisection method**: solve $\phi'(\alpha) = 0$ instead



Dyadic/binary and Fibonacci search




Assumption ★:

ϕ is **unimodal**, that is, ϕ has a unique minimum on (α', α'')


ϕ is decreasing on $[\alpha', \alpha^*]$ and ϕ is increasing on $[\alpha^*, \alpha'']$

ϕ is convex on $[\alpha', \alpha'']$
($\Leftrightarrow \phi'' > 0$)

Basic Splitting Step: for a pair $x_1 < x_2$ of points in the starting bracket




New interval if
 $\phi(x_2) > \phi(x_1)$



New interval if
 $\phi(x_2) < \phi(x_1)$

Exercise: Check that, under Assumption ★, for all $x_1 < x_2$ after the Basic Splitting Step the new interval contains α^* (hence, is a bracket).

Basic Splitting Step in “almost” two parts:
do the basic splitting step for x_1 and $x_1 + \epsilon$, where $\epsilon > 0$ is small



- Each Basic Splitting Step requires 2 evaluations of the function at x_1 and x_2 .
- In general, i.e., if Assumption ★ is violated, the Basic Splitting Step doesn't work!

Exercise: Give an example

Dyadic/binary search:
under **Assumption** ★

given the desired size $\epsilon > 0$ of the bracket

choose $\delta < \epsilon$ (usually much smaller)

repeat

1. Pick the midpoint $x_1 = \frac{\alpha' + \alpha''}{2}$
2. Do the **Basic Splitting Step in 'almost' two parts** using x_1 and $x_1 + \delta$
3. Update $[\alpha', \alpha'']$ with the new bracket from step 2 above

until $|\alpha'' - \alpha'| < \epsilon$

Exercise: How many evaluations of the function ϕ is required in the dyadic search in order to shrink the bracket by a factor of 100?

Fibonacci search (under Assumption ★)

- Fibonacci numbers are given by the recursive relation $F_{n+2} = F_{n+1} + F_n$, with starting condition $F_1 = F_2 = 1$.
- This generates the sequence 1, 1, 2, 3, 5, 8, 11, ...
- This sequence grows as $F_n \sim \frac{1}{\sqrt{5}} \varphi^n$, for n large, where $\varphi = \frac{1+\sqrt{5}}{2} \approx 1.6$ is the Golden ratio.

given the number of steps N

for $i = N, N - 1, \dots, 1$ **do**

if $i \neq 1$,

 1. Compute $x_1, x_2 \in [\alpha', \alpha'']$ such that

$$\frac{\alpha'' - x_1}{\alpha'' - \alpha'} = \frac{F_i}{F_{i+1}} \text{ and } \frac{x_2 - \alpha'}{\alpha'' - \alpha'} = \frac{F_i}{F_{i+1}}$$

 2. Do the **Basic Splitting Step** using x_1 and x_2

 3. Update $[\alpha', \alpha'']$ with the new bracket from step 2 above

otherwise


 Do the **Basic Splitting Step in 'almost' two parts** using $\frac{\alpha' + \alpha''}{2}$ and $\frac{\alpha' + \alpha''}{2} + \epsilon$

Observe that after this step, the length of the new bracket is proportional to the length of the previous bracket as F_i to F_{i+1}

Key Advantage: Fibonacci search uses significantly smaller evaluations of the function than the dyadic search because it re-uses some evaluation points! (see example on the next slide)

Exercise: How many evaluations of the function ϕ is required in the Fibonacci search in order to shrink the bracket by a factor of 100? Compare it to the corresponding result of the dyadic search.

Fibonacci search (under Assumption ★): an example



Consider using Fibonacci search with five function evaluations to minimize $f(x) = \exp(x-2) - x$ over the interval $[a, b] = [-2, 6]$. The first two function evaluations are made at $\frac{F_5}{F_6}$ and $1 - \frac{F_5}{F_6}$, along the length of the initial bracketing interval:

$$f(x^{(1)}) = f\left(a + (b-a)\left(1 - \frac{F_5}{F_6}\right)\right) = f(1) = -0.632$$

$$f(x^{(2)}) = f\left(a + (b-a)\frac{F_5}{F_6}\right) = f(3) = -0.282$$

The evaluation at $x^{(1)}$ is lower, yielding the new interval $[a, b] = [-2, 3]$. Two evaluations are needed for the next interval split:

$$x_{\text{left}} = a + (b-a)\left(1 - \frac{F_4}{F_5}\right) = 0$$

$$x_{\text{right}} = a + (b-a)\frac{F_4}{F_5} = 1$$

A third function evaluation is thus made at x_{left} , as x_{right} has already been evaluated:

$$f(x^{(3)}) = f(0) = 0.135$$

The evaluation at $x^{(1)}$ is lower, yielding the new interval $[a, b] = [0, 3]$. Two evaluations are needed for the next interval split:

$$x_{\text{left}} = a + (b-a)\left(1 - \frac{F_3}{F_4}\right) = 1$$

$$x_{\text{right}} = a + (b-a)\frac{F_3}{F_4} = 2$$

A fourth functional evaluation is thus made at x_{right} , as x_{left} has already been evaluated:

$$f(x^{(4)}) = f(2) = -1$$

The new interval is $[a, b] = [1, 3]$. A final evaluation is made just next to the center of the interval at $2 + \epsilon$, and it is found to have a slightly higher value than $f(2)$. The final interval is $[1, 2 + \epsilon]$.

Quadratic fit search



The method is based on the following observations:

- 'close' to the minima functions look like quadratic functions
- we can explicitly find minima of quadratic functions:

Lemma:

There exists a unique parabola that passes through any triple of distinct points (a, y_a) , (b, y_b) , (c, y_c) . This parabola has its extremum at

$$x^* = \frac{1}{2} \frac{y_a(b^2 - c^2) + y_b(c^2 - a^2) + y_c(a^2 - b^2)}{y_a(b - c) + y_b(c - a) + y_c(a - b)}$$



Exercise*: Show that the algorithm described on the next slide converges to a local minimum (assuming the function is smooth)

Quadratic fit search

given a triple $a < b < c$ where $[a, c]$ is a bracket of ϕ and $\phi(b) < \phi(a)$, $\phi(b) < \phi(c)$

repeat


1. Compute the critical point x^* of the parabola passing through a, b, c
2. If $x^* \in [b, c]$, then
 - Check which value is larger, $\phi(x^*)$ or $\phi(b)$:
 - i. If $\phi(b) > \phi(x^*)$, update the triple (a, b, c) with (b, x^*, c)
 - ii. If $\phi(b) < \phi(x^*)$, update the triple (a, b, c) with (a, b, x^*)
3. Otherwise
 - Again, check which value is larger, $\phi(x^*)$ or $\phi(b)$:
 - i. If $\phi(b) > \phi(x^*)$, update the triple (a, b, c) with (a, x^*, b)
 - ii. If $\phi(b) < \phi(x^*)$, update the triple (a, b, c) with (x^*, b, c)

until the $|a - c| < \epsilon$




That is, when $x^* \in [a, b)$
because of condition #

Or any other stopping criterion
based on variation of the function

#



In these examples, parabola at the current step is in orange; parabola at the next step is in blue



Bisection method

The method is based on the following observations:


- Instead of looking for a local minimum of ϕ , we can look for a solution of $\phi' = 0$
- We assume that $[\alpha', \alpha'']$ is a bracket for ϕ , and hence there exists a solution of $\phi' = 0$ on this interval

given an interval $[a, b]$ such that $\phi'(a) \cdot \phi'(b) < 0$


repeat

1. Compute the midpoint $c = \frac{a+b}{2}$
2. If $\phi'(a) \cdot \phi'(c) < 0$, update interval $[a, b]$ with $[a, c]$
3. If $\phi'(b) \cdot \phi'(c) < 0$, update interval $[a, b]$ with $[c, b]$

until $|a - b| < \epsilon$



- If $[\alpha', \alpha'']$ doesn't satisfy the condition $\phi'(\alpha') \cdot \phi'(\alpha'') < 0$, then one can try iteratively shrink this interval by a constant factor (say 2), until the condition is fulfilled. However, it might not always work (see an example of the function on the left where the bisection method can fail; this is the situation of a local minimum in a 'deep valley'). More sophisticated methods should be used instead.



Exercise: Let $\phi(x) = \frac{x^2}{2} - x$. Apply the bisection method to find an interval containing the minimizer of ϕ starting with the interval $[0, 1000]$. Execute 3 steps of the algorithm.

End of digression

Exact line search

$$\text{minimize}_{\alpha} f(\mathbf{x} + \alpha \mathbf{p})$$


Consider conducting a line search on $f(x_1, x_2, x_3) = \sin(x_1 x_2) + \exp(x_2 + x_3) - x_3$ from $\mathbf{x} = [1, 2, 3]$ in the direction $\mathbf{d} = [0, -1, -1]$. The corresponding optimization problem is:

$$\text{minimize}_{\alpha} \sin((1 + 0\alpha)(2 - \alpha)) + \exp((2 - \alpha) + (3 - \alpha)) - (3 - \alpha)$$

which simplifies to:

$$\text{minimize}_{\alpha} \sin(2 - \alpha) + \exp(5 - 2\alpha) + \alpha - 3$$

The minimum is at $\alpha \approx 3.127$ with $\mathbf{x} \approx [1, -1.126, -0.126]$.



Approximate line search

Find $\alpha^{(k)}$ approximately and move on with the descent method

For simplicity, $x_k = \mathbf{x}^{(k)}$, $p_k = \mathbf{p}^{(k)}$, $\alpha_k = \alpha^{(k)}$

We impose the following condition for α_k

$$\phi(\alpha_k) := f(x_k + \alpha_k p_k) < f(x_k) + c_1 \alpha_k (\nabla f(x_k))^T p_k, \quad c_1 \in (0, 1).$$

The condition is called (sufficient decrease condition).


Remarks.

- $\ell(\alpha_k) := f(x_k) + c_1 \alpha_k \nabla f^T(x_k) p_k$ is a linear function.
- For small values of $\alpha_k > 0$ we have $\phi(\alpha_k) < \ell(\alpha_k)$. This is so because $c_1 \in (0, 1)$ and then

$$\phi'(0) = (\nabla f(x_k))^T p_k < c_1 (\nabla f(x_k))^T p_k = \ell'(0) < 0.$$

Approximate line search

Sufficient decrease. We ask for a decrease proportional to α and $\phi'(0) = (\nabla f(x_k))^T p_k$. Usually $c_1 \approx 0.1$.




Approximate line search

Curvature condition. Since the previous condition is always satisfied for small values of α_k we need to add further conditions for termination. We use the so called **curvature condition**

$$(\nabla f(x_k + \alpha_k p_k))^T p_k \geq c_2 (\nabla f(x_k))^T p_k, \quad c_2 \in (c_1, 1)$$

In other words if $\phi'(\alpha_k)$ is not **negative enough** we terminate the k -step.



Approximate line search

Wolfe conditions

Definition. The conditions (together) to terminate the k -step given by

$$\begin{aligned} f(x_k + \alpha_k p_k) &< f(x_k) + c_1 \alpha_k (\nabla f(x_k))^T p_k, \\ (\nabla f(x_k + \alpha_k p_k))^T p_k &\geq c_2 (\nabla f(x_k))^T p_k, \end{aligned}$$

with $0 < c_1 < c_2 < 1$ are usually called **Wolfe conditions**.


Definition. The conditions (together) to terminate the k -step given by (we do not allow $\phi'(\alpha_k)$ to be too positive).

$$\begin{aligned} f(x_k + \alpha_k p_k) &< f(x_k) + c_1 \alpha_k (\nabla f(x_k))^T p_k, \\ |(\nabla f(x_k + \alpha_k p_k))^T p_k| &\leq |c_2 (\nabla f(x_k))^T p_k|, \end{aligned}$$

with $0 < c_1 < c_2 < 1$ are usually called **strong Wolfe conditions**.

Approximate line search

Wolfe conditions



Approximate line search

Wolfe conditions: existence

Lemma. Suppose $f : D \subset \mathbb{R}^n \rightarrow \mathbb{R}$ be a \mathcal{C}^1 function. Let p_k a descent direction at the point $x_k \in D$ and assume $f|_{L_{p_k}}$ is bounded below where $L_{p_k} = \{x \in \mathbb{R}^n \mid x = x_k + \alpha p_k, \alpha > 0\}$. Then if $0 < c_1 < c_2 < 1$ there exist intervals of step lengths satisfying the (strong) Wolfe conditions

Proof. Since $\ell'(\alpha_k) < 0$ (and constant) there exists a first intersection, $\hat{\alpha}_k > 0$, between $\ell(\alpha_k)$ and $\phi(\alpha_k)$:

$$f(x_k + \hat{\alpha}_k p_k) = f(x_k) + c_1 \hat{\alpha}_k (\nabla f(x_k))^T p_k. \quad (1)$$

The sufficient decrease condition it is satisfied for all $\alpha_k \in [0, \hat{\alpha}_k]$. By the Mean Value Theorem we have that there exists $\tilde{\alpha}_k \in [0, \hat{\alpha}_k]$ such that

$$f(x_k + \hat{\alpha}_k p_k) - f(x_k) = \hat{\alpha}_k (\nabla f(x_k + \tilde{\alpha}_k p_k))^T p_k$$

All together imply

$$(\nabla f(x_k + \tilde{\alpha}_k p_k))^T p_k = c_1 \hat{\alpha}_k (\nabla f(x_k))^T p_k > c_2 \hat{\alpha}_k (\nabla f(x_k))^T p_k.$$

Therefore $\tilde{\alpha}_k$ satisfies the Wolfe conditions and smoothness gives the desired interval.

Approximate line search

Convergence

Remark. Until this moment we just consider the **definition of the process**, that is the election of p_k and α_k . But we need to study if the process converge to **somewhere**.

Let p_k be a descent direction, and let θ_k the angle of p_k and $-\nabla f(x^*)$

$$\cos(\theta_k) = -\frac{1}{\|\nabla f(x_k)\| \|p_k\|} (\nabla f(x_k))^T p_k$$

Theorem. Assume notation above with p_k a descent direction and α_k satisfying Wolfe's conditions. Suppose f is \mathcal{C}^2 and bounded below in \mathbb{R}^n . Then

$$\sum_{k=0}^{\infty} \cos^2(\theta_k) \|\nabla f(x_k)\| < \infty. \quad (2)$$

Approximate line search

Convergence

Corollary. Under the above notation and assumptions we have

$$\cos^2(\theta_k) \|\nabla f(x_k)\| \rightarrow 0$$

Moreover if there exists $\delta > 0$ such that $\cos(\theta) > \delta$ then

$$\lim_{k \rightarrow \infty} \|\nabla f(x_k)\| = 0 \quad (\text{globally convergent algorithms})$$

Remark. The final δ -condition basically means that p_k do not get arbitrarily **orthogonal** to the gradient vector. This is, for instance, the case of the **steepest descent method**.