Parameterization, Curvature


1. Given the parameterization

$$\mathbf{r}(t) = \langle 2\cos(t), 3\sin(t) \rangle, \quad 0 \le t \le 2\pi.$$

(a) Show that x(t) and y(t) satisfy the equation for the ellipse

$$\frac{x^2}{4} + \frac{y^2}{9} = 1. \qquad \frac{x}{4}^2 + \frac{y^2}{9} = \frac{4\cos^2 t}{4} + \frac{9\sin^2 t}{4}$$

(b) Draw this ellipse on the axes below.



(c) Find
$$\mathbf{v} = \mathbf{r}'(0)$$
 and $\mathbf{a} = \mathbf{r}''(0)$ and find $\kappa = \frac{|\mathbf{v} \times \mathbf{a}|}{|\mathbf{v}|^3}$ at $t = 0$.,

$$\vec{v}(t) = \langle -2 \sin t, 3 \cos t \rangle \qquad \vec{v}(0) = \langle 0, 3 \rangle$$

$$\vec{a}(t) = \langle -2 \cos t, -3 \sin t \rangle \qquad \vec{a}(0) = \langle -2, 0 \rangle$$

$$1 = \frac{|\langle 0, 3 \rangle \times \langle -2, 0 \rangle|}{|\langle 0, 3 \rangle|^3} = \frac{6}{27} = \frac{2}{9}$$

2. Given the parameterization for a spiral,

$$\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle.$$

(a) Find the unit tangent vector **T** and the unit normal vector **N** at time t = 1. [Hint: you can find the normal direction by taking $\mathbf{v} \times \mathbf{a} \times \mathbf{v}$.]

$$\vec{v} = \langle -\sin t, \cos t, 1 \rangle \qquad \vec{a} = \langle -\cos t, -\sin t, o \rangle$$

$$\vec{T} = \frac{\vec{v}}{|\vec{v}|} = \frac{\langle -\sin t, \cos t, 1 \rangle}{\sqrt{1+1}} = \frac{\vec{b}}{\sqrt{2}} \langle -\sin t, \cos t, 1 \rangle$$

$$\vec{n} = \vec{\nabla} \times \vec{a} \times \vec{v}$$

$$= \langle t \cdot s \cdot t, cost, l \rangle \times \langle -s \cdot t, cost, l \rangle$$

$$= \langle -cost, 2s \cdot s \cdot t, o \rangle = \vec{N} = \langle -cost, s \cdot s \cdot t, o \rangle$$
(b) Find an equation for a line tangent to the spiral at time t_0 .

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -\sin t \\ \cos t \\ 1 \end{pmatrix} + \begin{pmatrix} \cos t \\ \sin t \\ t \end{pmatrix}$$

(c) Given $t_0 = \pi$, find the point at which the tangent line intersects the plane z = 0.

2

$$\begin{pmatrix} \dot{x} \\ \dot{x} \\ \dot{x} \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix} + 4 \begin{pmatrix} -1 \\ 0 \\ \pi \end{pmatrix}$$

$$Z=0$$
 => $0=t+\Pi$
 $t=-\Pi$

- 3. Give examples of the following.
 - (a) An equation for a cylinder so that the point (1,2,3) is on its surface.

(b) The point P on the plane 3x+2y+z=6 that is closest to the point Q(7,7,-1).

Find intersection

of & and plane:

$$3x + 2y + 2 = 3(3t+7) + 2(2t+7) + 1(t-1)$$
 $= 14t + 34 = 6$
 $t = -2$
 $(x, y, z) = (1, 3, -3)$

(c) A paraboloid that opens downwards in the z direction and intersects the xyplane in the ellipse $4x^2 + 9y^2 = 36$.

$$72 = 0$$
 $4x^{2} + 9y^{2} = 36 - 7$

(d) Two parallel lines L_1 and L_2 in the parallel planes x + 2y + 2z = 3 and x + 2y + 2z = 6 so that the distance between L_1 and L_2 is 3.

Choose
$$L_1 := \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ -1 \end{pmatrix} + \begin{pmatrix} 6 \\ 0 \\ 0 \end{pmatrix}$$

$$By \text{ aspection,}$$

$$L_2 := \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} + \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}$$

- 4. Given the points A(1, 2, 3) and B(5, 4, -2).
 - (a) Find both the parametric and symmetric equations of the straight line connecting A and B.

$$\vec{v} = 8 - A = \left(\frac{4}{7}, \frac{7}{-5} \right)$$

$$\left(\frac{x}{7} \right) = \left(\frac{4}{7}, \frac{7}{-5} \right) + \left(\frac{1}{2} \right)$$

$$\frac{x-1}{4} = \frac{y-7}{2} = \frac{z-3}{-5}$$

(b) Find the plane perpendicular to this line that goes through the point (1, 2, 4).

$$\vec{n} = \begin{pmatrix} 4 \\ \frac{2}{5} \end{pmatrix}$$

$$4x + 2y - 5z = 4(1) + 2(2) - 5(4)$$

$$4x + 2y - 5z = -12$$

(c) Find a plane that this line does not intersect.

$$\vec{n}$$
 must be \vec{L} to \vec{v}
 \Rightarrow choose $\vec{n} = (1, -2, 0)$
 $x - 2y + 0z = 0$
 $\boxed{x - 2y = 0}$