

Combinatorics of the critical loci of deep linear networks

March 23, 2026

Abstract

We study the critical loci of deep linear networks i.e. the set of parameters at which the gradient of the loss vanishes. The set of vanishing gradients can be described as an algebraic variety described by cyclic vanishing conditions. Using cyclic quiver representation theory we stratify the critical loci into orbits parametrized by cyclic rank patterns which are ranks of partial products of the deep linear networks weights matrices. The strata provides a combinatorial description of critical loci and in particular saddle points of a given rank. This stratification promise to be a useful tool to globally understand how the critical loci shape the training dynamics of stochastic gradient descent in deep linear networks.

1 Introduction

Critical points of deep linear networks were parametrized in Achour et al. [2024], where they were shown to satisfy vanishing cyclic conditions (see the proof of Proposition 10 in Appendix B.5). Furthermore, the stratification of the locus of global minima was obtained in Lehalleur and Rimányi [2024] through the use of affine quiver representation theory. The present work relies heavily on both of these results.

Our starting point is the observation that the cyclic vanishing conditions defining critical points can be interpreted as the vanishing of cyclic products in a representation of a cyclic quiver. This interpretation makes it possible to apply methods from cyclic quiver representation theory to the study of critical loci. In particular, the cyclic vanishing conditions effectively reduce the difficult problem of classifying indecomposable representations of cyclic quivers to that of classifying indecomposable modules over Nakayama algebras, which form a class of finite-dimensional algebras.

Using this framework, and following a strategy similar to that of Lehalleur and Rimányi [2024], we show that the critical locus admits a stratification indexed by a finite set. This provides a combinatorial description of saddle points, in which different classes of saddles are identified by their cyclic rank patterns.

1.1 Related works

1.2 Contributions

2 Setup and background

2.1 Notations

Notation	Meaning
$1, \dots, L$	Indices of the layers of the deep linear network (DLN)
0	Index of the input space
L	Index of the output space
d_l	Width of layer l , for $0 \leq l \leq L$
$d = (d_0, \dots, d_L)$	Dimension vector
Ω	Parameter space of the DLN
$\mathcal{F}$	Function space of the DLN
$W \in \mathcal{F}$	Student matrix (realized end-to-end map)
$\mu : \Omega \rightarrow \mathcal{F}$	Product map of the DLN
$r = \text{rank}(W)$	Rank of the student matrix
$d_r = (d_0, d_1 - r, \dots, d_L - r)$	Reduced dimension vector associated with rank r
W^*	Teacher matrix; equivalently, the global minimizer in function space
d_L	Maximal possible rank
$\Sigma_{YX} = \mathbb{E}[YX^\top]$	Input-output covariance matrix
$\Sigma_X = \mathbb{E}[XX^\top]$	Input covariance matrix
S	Subset of singular directions of the teacher matrix W^*
$U = [U_S \ U_Q]$	Matrix of left singular vectors of W^*
U_S	Left singular vectors of W^* corresponding to the singular values indexed by S
$P_S = U_S U_S^\top$	Orthogonal projector onto the singular subspace indexed by S
$G_d = \prod_{l=0}^L \text{GL}(d_l)$	Full base-change group
$G_{\text{hid}} = \prod_{l=1}^{L-1} \text{GL}(d_l)$	Hidden-layer symmetry group
Σ_d^r	Determinantal variety of matrices of rank r
$\Gamma_{d,\sigma}$	Cyclic variety with dimension vector d and closing arrow σ
$\Gamma_{d,XY}$	Cyclic variety with closing arrow $\sigma = \Sigma_{YX} U_Q$
$\mathcal{C}_{S,d}^r$	Critical locus associated with subset S , rank r , and dimension vector d
$\rho \in \mathcal{R}_d^{\text{orb}}$	Cyclic rank pattern
$m \in \mathcal{M}_d^+$	Cyclic multiplicity pattern
$M[i, j]$	Indecomposable representation of the cyclic quiver with relations, supported from vertex i to vertex j

2.2 Deep linear networks setup

Definition 2.1. A L -layers Deep Linear Network (DLN) is a function:

$$f : \mathbb{R}^{d_0} \times \prod_{l=0}^{L-1} \mathbb{R}^{d_l \times d_{l+1}} \rightarrow \mathbb{R}^{d_L}$$

$$(x, W_1, \dots, W_L) \mapsto W_L \dots W_1 x$$

The dimension d_0 is the dimension of the input data and the dimensions d_l , $1 \leq l \leq L$ are the width of the layers. A Deep Linear Network is a multi-linear function which is linear in each of the weight matrix W_l and the data.

Definition 2.2. Consider the product map:

$$\mu : \prod_{l=0}^{L-1} \mathbb{R}^{d_l \times d_{l+1}} \rightarrow \mathbb{R}^{d_0 \times d_L}$$

$$W := (W_1, \dots, W_L) \mapsto W_L \dots W_1$$

This function μ maps the parameter space $\Omega := \{(W_1, \dots, W_L) \in \prod_{l=0}^{L-1} \mathbb{R}^{d_l \times d_{l+1}}\}$ to the function space $\mathcal{F} \cong \mathbb{R}^{d_0 \times d_L}$ of a Deep Neural Network. A DLN is a matrix-monomial (so non-linear) in the parameter space Ω (the space above) and linear map in the function space $\mathcal{F}$ (the space below).

Definition 2.3. We consider the quadratic loss function of a DLN with training data $\{(x_i, y_i)\}_{i=1}^N$ i.i.d. :

$$\mathcal{L}_N(W) = \frac{1}{N} \sum_{i=1}^N \|W_L \dots W_1 x_i - y_i\|_2^2$$

With the covariance matrices $\Sigma_X = \frac{1}{N} \sum_{i=1}^N x_i x_i^\top$, $\Sigma_{XY} = \frac{1}{N} \sum_{i=1}^N x_i y_i^\top$, $\Sigma_{YX} = \frac{1}{N} \sum_{i=1}^N y_i x_i^\top$ and $\Sigma_Y = \frac{1}{N} \sum_{i=1}^N y_i y_i^\top$ identified with their expectation in the limit of large samples.

Definition 2.4. The teacher matrix (global minimum of the loss function) is defined as $W^* = \Sigma_{YX} \Sigma_X^{-1}$ and the teacher rank is defined as $r_{\max} = \text{rk}(W^*)$. The teacher SVD decomposition writes:

$$W^* = U \Sigma_{YX} \Sigma_X^{-1} V^\top$$

With $U = [U_S \ U_Q]$ where U_S are the left singular vectors of W^* corresponding to the singular values indexed by $S = 1, \dots, r$ and U_Q are the left singular vectors orthogonal to U_S .

Assumptions We will assume that Σ_X is invertible, that Σ_{XY} has full rank d_L . Furthermore we assume that the matrix: $\Sigma_{YX} \Sigma_X^{-1}$ has distinct singular values (non degenerate data). Furthermore the DLN has no bottleneck i.e. $d_i \geq d_L$ for all i . These are the same assumptions as in Achour et al. [2024].

2.3 Determinantal and cyclic varieties

Definition 2.5 (Residual space $\mathcal{Z}$). Fix $r \leq \min_i d_i$. Define the space of residuals:

$$\mathcal{Z}_d := \text{Mat}_{d_1, d_0} \times \prod_{\ell=2}^L \text{Mat}_{d_\ell, d_{\ell-1}}.$$

We will be interested in the space of residuals with dimension vector $d_r := (d_0, d_1 - r, \dots, d_L - r)$:

$$\mathcal{Z}_{d_r} := \text{Mat}_{d_1-r, d_0-r} \times \prod_{\ell=2}^L \text{Mat}_{d_\ell-r, d_{\ell-1}-r}.$$

Definition 2.6 (Cyclic variety Γ_{d_r}). Let $\Sigma_{XY} \in \text{Mat}_{d_0, d_L}$ and $U_Q \in \text{Mat}_{d_L, d_L-r}$ as defined in 2.4. Define the algebraic variety:

$$\Gamma_{d_r} := \{Z \in \mathcal{Z}_{d_r} \mid Z_{\ell-1} \cdots Z_2 Z_1 \Sigma_{XY} U_Q Z_L \cdots Z_{\ell+1} = 0, \ 1 \leq \ell \leq L \bmod L+1\},$$

Definition 2.7 (Variety Σ_d). Let $d = (d_0, \dots, d_L)$ and $r \leq \min_i d_i$. Define the determinantal variety of rank r matrices:

$$\Sigma_d^r := \{W \in \Omega \mid \text{rk}(W_L \dots W_1) = r\}$$

We can relate Σ_d^r to the residual variety Σ_{d-r}^0 :

$$\Sigma_{d-r}^0 = \{Z \in \mathcal{Z}_d, 1 \leq l \leq L \mid Z_L \dots Z_1 = 0\}$$

via the action of $GL(d) = \prod_{\ell=0}^L GL(d_\ell)$ on Ω :

$$\begin{aligned} \Sigma_d^r &= GL(d) \cdot \iota_r(\Sigma_{d-r}^0) \\ g \cdot W &= (g_L W_L g_{L-1}^{-1}, \dots, g_1 W_1 g_0^{-1}) \\ \iota_r(Z)_\ell &= \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} \end{aligned}$$

We will be interested in the space of residuals with dimension vector $d_r := (d_0, d_1 - r, \dots, d_L - r)$:

$$\Sigma_{d_r}^0 := \{Z \in \mathcal{Z} \mid Z_L \cdots Z_1 = 0\}.$$

Definition 2.8. Let $d_r := (d_0, d_1 - r, \dots, d_L - r)$, define the cyclic variety as an algebraic variety of residuals satisfying cyclic vanishing conditions:

$$\Gamma_{d_r, XY} := \Sigma_{d_r}^0 \cap \Gamma_{d_r}$$

2.4 Combinatorics of the global minima

2.5 Critical loci parametrization

Recall proposition 1 in Achour et al. [2024]:

Proposition 2.9. Let $W = (W_H, \dots, W_1)$ be a first-order critical point of $\mathcal{L}$ and set

$$r = \text{rk}(W_H \cdots W_1) \in \{0, \dots, r_{\max}\}.$$

Then there exists a unique subset $S \subset \{1, \dots, d_L\}$ with $|S| = r$ such that

$$W_H \cdots W_1 = U_S U_S^\top \Sigma_{YX} \Sigma_X^{-1}.$$

Definition 2.10 (Action of G_{hid}). Let $\mathbf{d}_{\text{hid}} = (d_1, \dots, d_{L-1})$. Define a group action of $G_{\text{hid}} = \prod_{\ell=1}^{L-1} GL(d_\ell, \mathbb{R})$ on Ω by:

$$g \cdot W = (W_L g_{L-1}^{-1}, (g_\ell W_\ell g_{\ell-1}^{-1})_{2 \leq \ell \leq L-1}, g_1 W_1)$$

We have the equivalence relation is defined via the definition just above.

Proposition 2.11. From Achour et al. [2024] proposition 9, proposition 10 and proof of proposition 10, appendix B.5 (gradient formulas), we can write the set of rank r critical points of a DLN associated with the subset of S indexing the singular vectors of W^* as the following algebraic variety:

$$\mathcal{C}_{d,S}^r := \{W \in \Omega \mid \exists Z \in \Gamma_{d_r, XY}, W_L \sim [U_S, U_Q Z_L], W_\ell \sim \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix}, W_1 \sim [\Sigma_X^{-1} \Sigma_{XY} U_S, Z_1^\top]^\top\}$$

In other words, $\mathcal{C}_{d,S}^r \subset \Omega$ denote the set of rank- r first-order critical points in a normal form of the DLN associated with S indexing the singular vectors of W^* , i.e. the set described by the existence of residual blocks $Z = (Z_1, \dots, Z_L) \in \Gamma_{d_r, XY}$ such that there exists $g \in G_{\text{hid}}$ with

$$W_L = [U_S, U_Q Z_L] g_{L-1}^{-1}, \quad W_\ell = g_\ell \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} g_{\ell-1}^{-1} \quad (2 \leq \ell \leq L-1), \quad W_1 = g_1 \begin{pmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ Z_1 \end{pmatrix}$$

In particular when computing the value of a DLN at a point in $\mathcal{C}_{d,S}^r$ we recover proposition 2.5. Furthermore, when $r = d_L$, one has $U_Q = 0$ and hence $\sigma = \Sigma_{XY} U_Q = 0$. In this case the cyclic residual variety $\Gamma_{d_r, XY}$ reduces to the residual representation space of an equioriented type- A quiver with dimension vector

$$d_r = (d_0, d_1 - d_L, \dots, d_{L-1} - d_L),$$

i.e. the corresponding stratification reduces to an affine-quiver stratification analogous to the treatment of the global-minimum stratification studied by Lehalleur–Rimányi. In the remaining of the document we will be interested in the case $r < d_L$ where the closing arrow σ is nonzero, i.e. the stratification of saddle points.

3 From the cyclic variety to the critical locus

3.1 Lifting the cyclic variety onto the critical locus

3.2 Stratification of the critical locus from stratifying the cyclic variety

4 Stratification of cyclic quivers with finite relation

5 Application: stratification of the critical loci

6 Discussion and future works

References

El Mehdi Achour, François Malgouyres, and Sébastien Gerchinovitz. The loss landscape of deep linear neural networks: a second-order analysis. *Journal of Machine Learning Research*, 25(242): 1–76, 2024.

Simon Pepin Lehalleur and Richárd Rimányi. Geometry of fibers of the multiplication map of deep linear neural networks. *arXiv preprint arXiv:2411.19920*, 2024.

Appendix