Notes on the Geometry of Deep Linear Networks

February 24, 2026

1 Deep Linear Networks

Definition 1.1. A deep linear network (DLN) is parameter-function map defined by a composition of linear transformations. For input space $\mathbb{R}^{d_0}$, output space $\mathbb{R}^{d_L}$, and L layers with weight matrices $W_1 \in \mathbb{R}^{d_1 \times d_0}, W_2 \in \mathbb{R}^{d_2 \times d_1}, \ldots, W_L \in \mathbb{R}^{d_L \times d_{L-1}}$, the DLN is given by:

$$f(x; W_1, W_2, \ldots, W_L) = W_L W_{L-1} \cdots W_1 x,$$

where $x \in \mathbb{R}^{d_0}$ is an input vector.

Note that the DLN f is a polynomial function of the entries of the weight matrices and linear in the input x.

Lemma 1.2. The space of parameter $\Omega = \mathbb{R}^{d_1 \times d_0} \times \mathbb{R}^{d_2 \times d_1} \times \cdots \times \mathbb{R}^{d_L \times d_{L-1}}$ is a a differentiable manifold of dimension $\sum_{l=1}^{L} d_l d_{l-1}$. Furthermore, the space Ω has a scalar product:

$$\langle U, V \rangle := \sum_{l=1}^{L} \langle U_l, V_l \rangle_F = \sum_{l=1}^{L} \mathrm{Tr}(U_l^\top V_l)$$

The differential is induced by the differential on the Euclidean spaces $\mathbb{R}^{d_l \times d_{l-1}}$. The scalar product is induced by the Frobenius scalar product on each of the Euclidean spaces $\mathbb{R}^{d_l \times d_{l-1}}$.

Definition 1.3. The empirical loss function of a DLN is defined as:

$$\mathcal{L}_N(W_1, W_2, \ldots, W_L) := \frac{1}{2N} \sum_{i=1}^{N} \||y_i - f(x_i; W_1, W_2, \ldots, W_L)\||_F^2,$$

where $\{(x_i, y_i)\}_{i=1}^N$ is a dataset of input-output pairs. The population loss is the expected loss over the data distribution. Assuming that the input data is whitened i.e. $x \sim N(0, I)$[1] then the square norm of the gap between the product of the weights and the target linear map.

$$\mathcal{L}(W_1, W_2, \ldots, W_L) := \frac{1}{2}||M - W||_F^2 = \frac{1}{2}||M - W_L W_{L-1} \cdots W_1||_F^2$$

We can derive the differential of the DLN map by using the usual product rule for differentials.

[1] This is a common assumption in the DLN literature

Proposition 1.4. Fix an input $x \in \mathbb{R}^{d_0}$ and write

$$h_{l-1} := W_{<l}x, \qquad W_{>l} := W_L \cdots W_{l+1}.$$

For a displacement $U = (U_l)_l \in \Omega$, the function f has a differential at any point $(W_1, W_2, \ldots, W_L) \in \Omega$ given by:

$$df[U] = \sum_{l=1}^{L} W_{>l} U_l h_{l-1}$$

From the differential, we can derive the Jacobian of the function f, by vectorizing and using the identity $\mathrm{vec}(AUB) = (B^\top \otimes A)\,\mathrm{vec}(U)$,

Lemma 1.5 (Jacobian from the differential)**.** Let $\theta := \big(\mathrm{vec}(W_1)^\top, \ldots, \mathrm{vec}(W_L)^\top\big)^\top \in \mathbb{R}^P$ with $P = \sum_{l=1}^{L} d_l d_{l-1}$, and similarly $\mathrm{vec}(U) := (\mathrm{vec}(U_1)^\top, \ldots, \mathrm{vec}(U_L)^\top)^\top$. By vectorizing the differential, we obtain:

$$\mathrm{vec}\big(df[U]\big) = \sum_{l=1}^{L} \big(h_{l-1}^\top \otimes W_{>l}\big)\,\mathrm{vec}(U_l).$$

By definition of the Jacobian $J(\theta; x) \in \mathbb{R}^{d_L \times P}$ (the unique matrix such that $\mathrm{vec}(df[U]) = J(\theta; x)\,\mathrm{vec}(U)$), we obtain the block form

$$J(\theta; x) = \big[\, J_1(x) \;\mid\; J_2(x) \;\mid\; \cdots \;\mid\; J_L(x) \,\big], \qquad J_l(x) \in \mathbb{R}^{d_L \times (d_l d_{l-1})}.$$

For each layer $l = 1, \ldots, L$,

$$\boxed{J_l(x) \;=\; h_{l-1}^\top \otimes W_{>l}}$$

so that, for any perturbation U_l,

$$\mathrm{vec}\big(W_{>l} U_l h_{l-1}\big) \;=\; J_l(x)\,\mathrm{vec}(U_l).$$

Lemma 1.6 (Differential of the loss)**.** Let

$$\mathcal{L}(W_1, \ldots, W_L) = \mathbb{E}\big[\langle y - f,\; y - f\rangle\big], \qquad r := y - f(W; x),$$

where the differential of f is

$$df[U] = \sum_{k=1}^{L} W_{>k} U_k h_{k-1}, \quad h_{k-1} := W_{<k}x.$$

Then, for any displacement $U = \{U_l\}_{l=1}^{L}$, the differential of $\mathcal{L}$ is

$$\boxed{d\mathcal{L}[U] = \sum_{l=1}^{L} \big\langle U_l,\; -\mathbb{E}\big[\, W_{>l}^\top (y - f)\, h_{l-1}^\top \big]\big\rangle_F .}$$

Equivalently, the gradient of $\mathcal{L}$ with respect to W_l is

$$\boxed{\nabla_{W_l}\mathcal{L} = -\,\mathbb{E}\big[\, W_{>l}^\top (y - f)\, h_{l-1}^\top \big].}$$

Derivation. Start from the definition of $\mathcal{L}$:

$$\mathcal{L} = \mathbb{E}\big[\langle y - f,\ y - f\rangle\big] = \mathbb{E}\big[\langle r, r\rangle\big], \qquad r := y - f.$$

Taking the differential gives

$$d\mathcal{L}[U] = \mathbb{E}\big[\langle r, dr[U]\rangle\big] = -\,\mathbb{E}\big[\langle r, df[U]\rangle\big],$$

since $dr[U] = -df[U]$. Substituting the expression of $df[U]$,

$$df[U] = \sum_{k=1}^{L} W_{>k} U_k h_{k-1},$$

yields

$$d\mathcal{L}[U] = -\,\mathbb{E}\left[\sum_{k=1}^{L} \langle r,\ W_{>k} U_k h_{k-1}\rangle\right].$$

Using the identity $\langle AUB, v\rangle = \langle U, A^\top v B^\top\rangle_F$ gives

$$d\mathcal{L}[U] = \sum_{k=1}^{L} \big\langle U_k,\ -\mathbb{E}\big[\, W_{>k}^\top r\, h_{k-1}^\top\big]\big\rangle_F\,.$$

Identifying coefficients of U_k yields the stated gradient.

Corollary 1.7 (Population loss case)**.** If $y = Mx$ and $\Sigma_x = \mathbb{E}[xx^\top]$, then with $\Delta := M - W$ we have

$$\boxed{\nabla_{W_l}\mathcal{L} = -\,W_{>l}^\top\, \Delta\, \Sigma_x\, W_{<l}^\top.}$$

In the whitened input case $\Sigma_x = I$,

$$\boxed{\nabla_{W_l}\mathcal{L} = -\,W_{>l}^\top\, \Delta\, W_{<l}^\top.}$$

Derivation. Since $y - f = \Delta x$ and $h_{l-1} = W_{<l}x$,

$$\mathbb{E}\big[(y - f)\, h_{l-1}^\top\big] = \mathbb{E}\big[\Delta x x^\top W_{<l}^\top\big] = \Delta\, \Sigma_x\, W_{<l}^\top,$$

and substituting this identity into the gradient formula of the lemma gives the result.

Remark 1.8 (Use of the Riesz representation theorem)**.** The differential $d\mathcal{L}$ is a linear functional on the parameter space:

$$d\mathcal{L} : \mathbb{R}^P \to \mathbb{R}, \qquad d\mathcal{L}(d\theta) = -\langle r,\ J(x)\, d\theta\rangle.$$

By the Riesz representation theorem in Euclidean space, there exists a unique vector $\nabla_\theta\mathcal{L} \in \mathbb{R}^P$ such that $d\mathcal{L}(d\theta) = \langle \nabla_\theta\mathcal{L}, d\theta\rangle$ for all $d\theta$. Using the adjoint identity $\langle a, Jb\rangle = \langle J^\top a, b\rangle$, we obtain

$$d\mathcal{L}(d\theta) = \langle -J(x)^\top r,\ d\theta\rangle,$$

hence

$$\boxed{\nabla_\theta\mathcal{L} = -J(x)^\top r.}$$

Definition 1.9. A point $(W_1, W_2, \ldots, W_L)$ is a critical point of the loss $\mathcal{L}$ if and only if $\nabla_{W_l}\mathcal{L} = 0$ for all $l = 1, 2, \ldots, L$. There are four kinds of critical points:

- Global minimum: $\mathcal{L}(W^\star) \leq \mathcal{L}(W)\ \forall W \in \Omega$.
- Local minimum: there exist an open set $\mathcal{U}$ of Ω such that $\mathcal{L}(W^\star) \leq \mathcal{L}(W')\ \forall W' \in \mathcal{U}$.
- Saddle points: $\nabla_{W_l}\mathcal{L} = 0\ \forall l$ and W_l is neither a local nor a global minimum.
- A saddle point is **strict** if the Hessian has a strictly negative eigenvalue and **non-strict** otherwise (i.e. there exists at least one zero eigenvalue of the Hessian).

The following lemma simply follows from the definition of the gradient of the loss.

Lemma 1.10. Let $\Delta := M - W$, a point $(W_1, W_2, \ldots, W_L)$ is a critical point of the loss $\mathcal{L}$ if and only if for all $l = 1, 2, \ldots, L$:

$$W_{>l}^\top \Delta W_{<l}^\top = 0$$

Lemma 1.11 (2-differential of the DLN map)**.** Let

$$f(W_1, \ldots, W_L; x) = W_L W_{L-1} \cdots W_1 x,$$

and define the shorthand notation

$$W_{>l} := W_L \cdots W_{l+1}, \quad W_{<l} := W_{l-1} \cdots W_1, \quad a_{l-1} := W_{<l} x.$$

Then the first differential of f in direction $U = \{U_l\}_{l=1}^L$ is

$$df[U] = \sum_{l=1}^{L} W_{>l}\, U_l\, W_{<l}\, x.$$

The second differential of f in directions $U = \{U_l\}$ and $V = \{V_l\}$ is given by the symmetric form:

$$\boxed{d^2 f[U, V] = \sum_{1 \leq i < j \leq L} \Big(W_{>j}\, V_j\, W_{j-1:i+1}\, U_i\, W_{<i} \ + \ W_{>i}\, V_i\, W_{i-1:j+1}\, U_j\, W_{<j} \Big) x.}$$

Derivation: The 2-differential can be computed using the product rule:

$$d^2 f[U, V] = \sum_{l=1}^{L} \left(dW_{>l}[V]\, U_l\, W_{<l} \ + \ W_{>l}\, U_l\, dW_{<l}[V] \right) x,$$

$$\text{where} \quad dW_{>l}[V] = \sum_{j>l} W_{>j}\, V_j\, W_{j-1:l+1}, \qquad dW_{<l}[V] = \sum_{j<l} W_{l-1:j+1}\, V_j\, W_{<j}.$$

Substituting these expressions yields the explicit bilinear form:

$$d^2 f[U, V] = \sum_{l=1}^{L} \sum_{j>l} W_{>j}\, V_j\, W_{j-1:l+1}\, U_l\, W_{<l}\, x \ + \ \sum_{l=1}^{L} \sum_{j<l} W_{>l}\, U_l\, W_{l-1:j+1}\, V_j\, W_{<j}\, x.$$

Reindexing over ordered pairs (i, j) with $1 \leq i < j \leq L$ gives the symmetric form stated in the lemma.

Remark 1.12.

- The map f is linear in each W_l separately, so $\partial^2 f/\partial W_l^2 = 0$.
- All curvature arises from cross-layer interactions between distinct layers $i \neq j$.
- The 2-differential is symmetric, i.e. $d^2 f[U, V] = d^2 f[V, U]$.

Corollary 1.13 (Quadratic variation)**.** For a single infinitesimal perturbation $dW = \{dW_l\}_{l=1}^L$, we have

$$\boxed{d^2 f = \sum_{1 \le i < j \le L} \Big(W_{>j}\, dW_j\, W_{j-1:i+1}\, dW_i\, W_{<i} \;+\; W_{>i}\, dW_i\, W_{i-1:j+1}\, dW_j\, W_{<j} \Big) x.}$$

Lemma 1.14 (Expected squared differential of the DLN map)**.** Let

$$df[U] \;=\; \sum_{l=1}^{L} W_{>l}\, U_l\, h_{l-1}, \qquad h_{l-1} := W_{<l} x, \qquad \Sigma_x := \mathbb{E}[xx^\top],$$

and define

$$G_{lm} := W_{>l}^\top W_{>m} \in \mathbb{R}^{d_l \times d_m}, \qquad C_{ml} := \mathbb{E}[\, h_{m-1} h_{l-1}^\top \,] = W_{<m}\, \Sigma_x\, W_{<l}^\top \in \mathbb{R}^{d_{m-1} \times d_{l-1}}.$$

Then the expected squared norm of the differential is

$$\boxed{\mathbb{E}\langle df, df \rangle = \sum_{l,m=1}^{L} \mathrm{Tr}\big(U_l^\top\, G_{lm}\, U_m\, C_{ml} \big) \;=\; \sum_{l,m=1}^{L} \mathrm{vec}(U_l)^\top \big(C_{ml}^\top \otimes G_{lm} \big)\, \mathrm{vec}(U_m).}$$

Derivation. Starting from $df[U] = \sum_{l=1}^L W_{>l} U_l h_{l-1}$,

$$\begin{aligned}
\langle df, df \rangle &= \Big(\sum_l W_{>l} U_l h_{l-1} \Big)^\top \Big(\sum_m W_{>m} U_m h_{m-1} \Big) \\
&= \sum_{l,m} h_{l-1}^\top U_l^\top\, W_{>l}^\top W_{>m}\, U_m\, h_{m-1} = \sum_{l,m} h_{l-1}^\top U_l^\top\, G_{lm}\, U_m\, h_{m-1}.
\end{aligned}$$

Taking expectation and using $C_{ml} = \mathbb{E}[h_{m-1} h_{l-1}^\top]$,

$$\mathbb{E}\langle df, df \rangle = \sum_{l,m} \mathbb{E}\big[h_{l-1}^\top U_l^\top G_{lm} U_m h_{m-1} \big] = \sum_{l,m} \mathrm{Tr}\big(U_l^\top G_{lm} U_m\, C_{ml} \big),$$

which yields the *trace form*. For the *vectorized form*, use $\mathrm{Tr}(A^\top B) = \mathrm{vec}(A)^\top\, \mathrm{vec}(B)$ and $\mathrm{vec}(AU_mB) = (B^\top \otimes A)\, \mathrm{vec}(U_m)$ to obtain

$$\mathrm{Tr}\big(U_l^\top G_{lm} U_m\, C_{ml} \big) = \mathrm{vec}(U_l)^\top \big(C_{ml}^\top \otimes G_{lm} \big)\, \mathrm{vec}(U_m).$$

Finally, since $h_{k-1} = W_{<k} x$ and $\Sigma_x = \mathbb{E}[xx^\top]$, we have $C_{ml} = W_{<m} \Sigma_x W_{<l}^\top$, completing the derivation.

Lemma 1.15 (Second differential of the squared-loss)**.** Let $f(W;x) = W_L \cdots W_1 x$ with $W_l \in \mathbb{R}^{d_l \times d_{l-1}}$, and write

$$W_{>l} := W_L \cdots W_{l+1} \in \mathbb{R}^{d_L \times d_l}, \quad W_{<l} := W_{l-1} \cdots W_1 \in \mathbb{R}^{d_{l-1} \times d_0}, \quad h_{l-1} := W_{<l} x.$$

Let the residual be $r := y - f(W;x) \in \mathbb{R}^{d_L}$, and define

$$\Sigma_x := \mathbb{E}[xx^\top] \in \mathbb{R}^{d_0 \times d_0}, \qquad S := \mathbb{E}[\, r\, x^\top\,] \in \mathbb{R}^{d_L \times d_0}.$$

For two displacement directions $U = \{U_l\}_{l=1}^L$ and $V = \{V_l\}_{l=1}^L$, the second differential of

$$\mathcal{L}(W) := \tfrac{1}{2}\, \mathbb{E}\big[\|y - f(W;x)\|_2^2\big]$$

is the symmetric bilinear form

$$\boxed{\begin{aligned} d^2\mathcal{L}[U,V] = &\sum_{l,m=1}^{L} \operatorname{Tr}\big(U_l^\top\, G_{lm}\, V_m\, C_{ml}\big) \\ &- \sum_{1 \le i < j \le L} \Big(\operatorname{Tr}\big(S^\top W_{>j} V_j W_{j-1:i+1} U_i W_{<i}\big) + \operatorname{Tr}\big(S^\top W_{>i} V_i W_{i-1:j+1} U_j W_{<j}\big)\Big), \end{aligned}}$$

where

$$G_{lm} := W_{>l}^\top W_{>m} \in \mathbb{R}^{d_l \times d_m}, \qquad C_{ml} := \mathbb{E}[\, h_{m-1} h_{l-1}^\top\,] = W_{<m}\, \Sigma_x\, W_{<l}^\top \in \mathbb{R}^{d_{m-1} \times d_{l-1}}.$$

Equivalently, in block–Kronecker (Hessian) form, if $\theta = (\operatorname{vec}(W_1); \ldots; \operatorname{vec}(W_L))$, then

$$d^2\mathcal{L}[U,V] = \sum_{l,m=1}^{L} \operatorname{vec}(U_l)^\top\, H_{lm} \operatorname{vec}(V_m), \quad H_{lm} = \underbrace{C_{ml}^\top \otimes G_{lm}}_{\text{metric block}} - \underbrace{R_{lm}}_{\text{residual block}},$$

with

$$R_{lm} = \begin{cases} W_{m-1:l+1}^\top \otimes \big(W_{<l} S^\top W_{>m}\big), & l < m, \\ W_{l-1:m+1}^\top \otimes \big(W_{<m} S^\top W_{>l}\big), & l > m, \\ 0, & l = m. \end{cases}$$

In the population case $y = Mx$ with $\Delta := M - W$ we have $S = \Delta\, \Sigma_x$, and in the whitened case $\Sigma_x = I$ this reduces to $S = \Delta$ and $C_{ml} = W_{<m} W_{<l}^\top$. At any global minimizer ($\Delta = 0$), the residual blocks vanish and $d^2\mathcal{L}$ equals the positive semidefinite metric term.

Derivation. *Step 1 (First and second differentials of f).* Since $f(W;x) = W_L \cdots W_1 x$ is linear in each W_l separately,

$$df[U] = \sum_{l=1}^{L} W_{>l}\, U_l\, h_{l-1}, \qquad h_{l-1} = W_{<l} x.$$

By the product rule (Lemma 1.11),

$$d^2 f[U,V] = \sum_{1 \le i < j \le L} \Big(W_{>j} V_j W_{j-1:i+1} U_i W_{<i} + W_{>i} V_i W_{i-1:j+1} U_j W_{<j} \Big) x. \tag{$*$}$$

Step 2 (First differential of $\mathcal{L}$). Let $r := y - f$. Then

$$\mathcal{L} = \tfrac{1}{2}\,\mathbb{E}\langle r, r\rangle, \quad dr[U] = -df[U], \quad d\mathcal{L}[U] = \mathbb{E}\langle r, dr[U]\rangle = -\mathbb{E}\langle r, df[U]\rangle. \tag{1}$$

Step 3 (Master identity for $d^2\mathcal{L}$). Differentiate (**??**) in direction V:

$$\begin{aligned} d^2\mathcal{L}[U,V] &= \mathbb{E}\big[\langle dr[V], dr[U]\rangle\big] + \mathbb{E}\big[\langle r,\, d^2 r[U,V]\rangle\big] \\ &= \mathbb{E}\big[\langle -df[V], -df[U]\rangle\big] + \mathbb{E}\big[\langle r,\, -d^2 f[U,V]\rangle\big] \\ &= \boxed{\mathbb{E}\langle df[U], df[V]\rangle} - \boxed{\mathbb{E}\langle r,\, d^2 f[U,V]\rangle}. \end{aligned} \tag{2}$$

Step 4 (Metric term $\mathbb{E}\langle df[U], df[V]\rangle$). Using $df[U] = \sum_l W_{>l} U_l h_{l-1}$ and $df[V] = \sum_m W_{>m} V_m h_{m-1}$,

$$\langle df[U], df[V]\rangle = \sum_{l,m} h_{l-1}^\top U_l^\top\, W_{>l}^\top W_{>m}\, V_m\, h_{m-1}.$$

Taking expectation and defining

$$G_{lm} := W_{>l}^\top W_{>m}, \qquad C_{ml} := \mathbb{E}[\, h_{m-1} h_{l-1}^\top\,] = W_{<m} \Sigma_x W_{<l}^\top,$$

we obtain

$$\boxed{\mathbb{E}\langle df[U], df[V]\rangle = \sum_{l,m} \mathrm{Tr}\big(U_l^\top\, G_{lm}\, V_m\, C_{ml}\big).} \tag{3}$$

(Use $\mathbb{E}[a^\top B c] = \mathrm{Tr}(B\,\mathbb{E}[c a^\top])$.)

Step 5 (Residual–curvature term $\mathbb{E}\langle r, d^2 f[U,V]\rangle$). From (**??**),

$$\langle r,\, d^2 f[U,V]\rangle = \sum_{i<j} \Big(r^\top W_{>j} V_j W_{j-1:i+1} U_i W_{<i} x + r^\top W_{>i} V_i W_{i-1:j+1} U_j W_{<j} x \Big).$$

Let $S := \mathbb{E}[r\, x^\top]$. Using the identity

$$\mathbb{E}\,[r^\top A x] = \mathrm{Tr}(S^\top A) \quad \text{for any } A \in \mathbb{R}^{d_L \times d_0}, \tag{4}$$

we get

$$\boxed{\mathbb{E}\langle r,\, d^2 f[U,V]\rangle = \sum_{i<j} \Big(\mathrm{Tr}\big(S^\top W_{>j} V_j W_{j-1:i+1} U_i W_{<i}\big) + \mathrm{Tr}\big(S^\top W_{>i} V_i W_{i-1:j+1} U_j W_{<j}\big)\Big).} \tag{5}$$

(Proof of (**??**): $r^\top A x = \mathrm{Tr}(r^\top A x) = \mathrm{Tr}(A x r^\top)$; take expectation and use linearity to get $\mathrm{Tr}(A\,\mathbb{E}[x r^\top]) = \mathrm{Tr}(S^\top A)$.)

Step 6 (Assemble). Subtract (**??**) from (**??**) in (**??**) to obtain the trace formula in the lemma.

Step 7 (Block–Kronecker form). Use the identity

$$\mathrm{Tr}(A\, V\, B\, U) = \mathrm{vec}(U)^\top\, (B^\top \otimes A)\, \mathrm{vec}(V), \tag{6}$$

to convert each trace:

$$\mathrm{Tr}\big(U_l^\top G_{lm} V_m C_{ml}\big) = \mathrm{vec}(U_l)^\top \left(C_{ml}^\top \otimes G_{lm}\right) \mathrm{vec}(V_m).$$

For $i < j$,

$$\mathrm{Tr}\big(S^\top W_{>j} V_j W_{j-1:i+1} U_i W_{<i}\big) = \mathrm{vec}(U_i)^\top \big(W_{j-1:i+1}^\top \otimes (W_{<i} S^\top W_{>j})\big) \mathrm{vec}(V_j),$$

and symmetrically with (i, j) swapped for the other cross term. Grouping all blocks yields H_{lm} stated in the lemma. Symmetry $H_{lm}^\top = H_{ml}$ follows from exchanging $U \leftrightarrow V$ in (**??**) and the definitions of G, C.

□

Proposition 1.16 (Hessian of the squared loss as a block operator)**.** Let $f(W; x) = W_L \cdots W_1 x$ with $W_l \in \mathbb{R}^{d_l \times d_{l-1}}$. For $1 \le l \le L$ set

$$W_{>l} := W_L \cdots W_{l+1} \in \mathbb{R}^{d_L \times d_l}, \quad W_{<l} := W_{l-1} \cdots W_1 \in \mathbb{R}^{d_{l-1} \times d_0}, \quad h_{l-1} := W_{<l} x.$$

Let $r := y - f(W; x)$, $\Sigma_x := \mathbb{E}[xx^\top] \in \mathbb{R}^{d_0 \times d_0}$ and

$$S := \mathbb{E}[\, r\, x^\top \,] \in \mathbb{R}^{d_L \times d_0}, \qquad G_{lm} := W_{>l}^\top W_{>m} \in \mathbb{R}^{d_l \times d_m}, \qquad C_{ml} := \mathbb{E}[\, h_{m-1} h_{l-1}^\top \,] = W_{<m} \Sigma_x W_{<l}^\top.$$

Define the *middle product* between layers l and m by

$$W_{m-1:l+1} := \begin{cases} W_{m-1} \cdots W_{l+1} \in \mathbb{R}^{d_{m-1} \times d_l}, & \text{if } m > l+1, \\ I_{d_l}, & \text{if } m = l+1. \end{cases}$$

(When $l > m$, we will only use $W_{l-1:m+1}$ defined analogously.)

Consider the squared-error population loss

$$\mathcal{L}(W) := \tfrac{1}{2}\, \mathbb{E}\big[\|y - f(W; x)\|_2^2\big].$$

Then the Hessian $\nabla^2 \mathcal{L}(W)$, viewed as a block operator acting on a perturbation $V = \{V_m\}_{m=1}^L$, has blocks

$$\big(\nabla^2 \mathcal{L}(W)\,[V]\big)_l = \sum_{m=1}^{L} \big(\nabla^2_{W_l, W_m} \mathcal{L}\big)[V_m],$$

with the following explicit actions:

$$\boxed{\begin{aligned} \big(\nabla^2_{W_l, W_l} \mathcal{L}\big)[V_l] &= G_{ll}\, V_l\, C_{ll} \;=\; W_{>l}^\top W_{>l}\, V_l\, W_{<l} \Sigma_x W_{<l}^\top, \\ \big(\nabla^2_{W_l, W_m} \mathcal{L}\big)[V_m] &= G_{lm}\, V_m\, C_{ml} \;-\; \underbrace{\big(W_{<l} S^\top W_{>m}\big)\, V_m\, W_{m-1:l+1}}_{\text{residual–curvature coupling}}, \qquad l \neq m. \end{aligned}}$$

Equivalently, in block–Kronecker form, if $\theta = (\mathrm{vec}(W_1); \ldots; \mathrm{vec}(W_L))$,

$$\nabla^2 \mathcal{L}(W) \;\equiv\; H = \big[H_{lm}\big]_{l,m=1}^L, \qquad d^2 \mathcal{L}[U, V] = \sum_{l,m} \mathrm{vec}(U_l)^\top H_{lm}\, \mathrm{vec}(V_m),$$

with

$$\boxed{H_{lm} = \begin{cases} C_{ml}^\top \otimes G_{lm}, & l = m, \\ C_{ml}^\top \otimes G_{lm} \;-\; W_{m-1:l+1}^\top \otimes \big(W_{<l} S^\top W_{>m}\big), & l \neq m. \end{cases}}$$

In the population case $y = Mx$ one has $S = \Delta\,\Sigma_x$ with $\Delta := M - W$, and in the whitened case $\Sigma_x = I$,

$$\big(\nabla^2_{W_l,W_m}\mathcal{L}\big)[V_m] = W_{>l}^\top W_{>m}\, V_m\, W_{<m} W_{<l}^\top \;-\; \Delta\, V_m\, W_{m-1:l+1}, \qquad l \neq m,$$

while the diagonal block remains $\big(\nabla^2_{W_l,W_l}\mathcal{L}\big)[V_l] = W_{>l}^\top W_{>l}\, V_l\, W_{<l} W_{<l}^\top$.

Derivation. *(i) From the 2-differential to the Hessian blocks.)* By Lemma 1.15 (second differential),

$$d^2\mathcal{L}[U,V] = \sum_{l,m} \mathrm{Tr}\big(U_l^\top\, G_{lm}\, V_m\, C_{ml}\big) - \sum_{i<j} \Big(\mathrm{Tr}\big(S^\top W_{>j} V_j W_{j-1:i+1} U_i W_{<i}\big) + \mathrm{Tr}\big(S^\top W_{>i} V_i W_{i-1:j+1} U_j W_{<j}\big)\Big). \tag{1}$$

We will rewrite each trace as a Frobenius inner product in U_l with a linear image of V_m, thereby identifying the operator $(\nabla^2_{W_l,W_m}\mathcal{L})[\cdot]$.

(ii) Metric term. Using $\mathrm{Tr}(U_l^\top A V_m B) = \langle U_l,\ A V_m B\rangle_F$,

$$\sum_{l,m} \mathrm{Tr}\big(U_l^\top\, G_{lm}\, V_m\, C_{ml}\big) = \sum_{l,m} \big\langle U_l,\ G_{lm}\, V_m\, C_{ml}\big\rangle_F. \tag{2}$$

(iii) Residual–curvature term. For the first sum in (1) with $i < j$, cyclicity of trace yields

$$\mathrm{Tr}\big(S^\top W_{>j} V_j W_{j-1:i+1} U_i W_{<i}\big) = \mathrm{Tr}\big(U_i^\top\, (W_{<i} S^\top W_{>j})\, V_j\, W_{j-1:i+1}\big) = \big\langle U_i,\, (W_{<i} S^\top W_{>j})\, V_j\, W_{j-1:i+1}\big\rangle_F. \tag{3}$$

The second sum in (1) (swap i, j) similarly gives, for $i < j$,

$$\mathrm{Tr}\big(S^\top W_{>i} V_i W_{i-1:j+1} U_j W_{<j}\big) = \big\langle U_j,\, (W_{<j} S^\top W_{>i})\, V_i\, W_{i-1:j+1}\big\rangle_F. \tag{4}$$

Combining (3) and (4) and reindexing $(i,j) \mapsto (l,m)$, we obtain

$$\sum_{i<j} \cdots = \sum_{l<m} \big\langle U_l,\, (W_{<l} S^\top W_{>m})\, V_m\, W_{m-1:l+1}\big\rangle_F \;+\; \sum_{m<l} \big\langle U_l,\, (W_{<l} S^\top W_{>m})\, V_m\, W_{m-1:l+1}\big\rangle_F. \tag{5}$$

Thus, for *every* ordered pair (l,m) with $l \neq m$,

$$-\mathbb{E}\langle r,\, d^2 f[U,V]\rangle = -\sum_{l\neq m} \big\langle U_l,\, (W_{<l} S^\top W_{>m})\, V_m\, W_{m-1:l+1}\big\rangle_F. \tag{6}$$

(iv) Identification of the operator blocks. Collecting (2) and (6) into $d^2\mathcal{L}[U,V] = \sum_{l,m}\langle U_l,\ \cdot\,\rangle_F$, we read off the unique linear maps

$$\boxed{\begin{aligned} \big(\nabla^2_{W_l,W_l}\mathcal{L}\big)[V_l] &= G_{ll}\, V_l\, C_{ll}, \\ \big(\nabla^2_{W_l,W_m}\mathcal{L}\big)[V_m] &= G_{lm}\, V_m\, C_{ml} \;-\; (W_{<l} S^\top W_{>m})\, V_m\, W_{m-1:l+1}, \qquad l \neq m, \end{aligned}}$$

since for all test directions U_l,

$$\langle U_l,\ (\nabla^2_{W_l,W_m}\mathcal{L})[V_m]\rangle_F = \mathrm{Tr}\big(U_l^\top G_{lm} V_m C_{ml}\big) - 1_{l\neq m}\,\mathrm{Tr}\big(U_l^\top (W_{<l} S^\top W_{>m}) V_m W_{m-1:l+1}\big).$$

(v) Kronecker form. Using $\mathrm{Tr}(U^\top AVB) = \mathrm{vec}(U)^\top (B^\top \otimes A)\mathrm{vec}(V)$, the bilinear representation $d^2\mathcal{L}[U,V] = \sum_{l,m} \mathrm{vec}(U_l)^\top H_{lm}\mathrm{vec}(V_m)$ follows with

$$H_{lm} = C_{ml}^\top \otimes G_{lm} 1_{l\neq m}\big(W_{m-1:l+1}^\top \otimes (W_{<l} S^\top W_{>m})\big).$$

This H is symmetric ($H_{lm}^\top = H_{ml}$) by construction.

(vi) Population/whitened specializations. If $y = Mx$, then $r = \Delta x$ with $\Delta := M - W$ and $S = \Delta\,\Sigma_x$. For whitened inputs $\Sigma_x = I$,

$$C_{ml} = W_{<m} W_{<l}^\top, \qquad S^\top = \Delta^\top,$$

which yields the stated simplified forms. □

2 DLNs as Quiver Representations

Definition 2.1. A *quiver* is a directed graph $Q = (P, E, s, t)$ where P is the set of vertices, E the set of directed edges, and $s, t : E \to P$ are the source and target maps.

Definition 2.2. A *representation* of a quiver Q over a field k assigns a vector space V_v to each vertex $v \in P$ and a linear map $\phi_e : V_{s(e)} \to V_{t(e)}$ to each directed edge $e \in E$. For a fixed *dimension vector* $d = (d_v)_{v\in P}$ with $d_v = \dim_k(V_v)$, we denote by $\mathrm{Rep}_d(Q)$ the space of d-dimensional representations of Q over k:

$$\mathrm{Rep}_d(Q) \cong \bigoplus_{e\in E} \mathrm{Hom}_k(k^{d_{s(e)}}, k^{d_{t(e)}}).$$

Definition 2.3. The *path algebra* kQ is the k-vector space with basis given by all paths in Q (including a trivial path e_v of length 0 at each vertex v). Multiplication is given by concatenation of paths when composable and 0 otherwise, extended by linearity.

Proposition 2.4. The category of representations of a quiver Q is equivalent to the category of left modules over its path algebra kQ.

Sketch of proof: Given a representation (V_v, ϕ_e), define the module $M = \bigoplus_{v\in P} V_v$ with e_v acting as projection onto V_v and an arrow $a : v \to w$ acting as ϕ_a on V_v and 0 elsewhere. Conversely, for a kQ-module M, define $V_v = e_v M$ and ϕ_a the action of a on M. These constructions define functors that are quasi-inverse to each other.

Definition 2.5. A *deep linear network (DLN)* of depth L is a linear map $\phi \in \mathrm{Hom}_k(k^{d_0}, k^{d_L})$ that factorizes as a composition of linear maps

$$\phi = \phi_L \phi_{L-1} \cdots \phi_1,$$

where $\phi_\ell \in \mathrm{Hom}_k(k^{d_{\ell-1}}, k^{d_\ell})$ for $\ell = 1, \ldots, L$.

Proposition 2.6. A DLN with dimensions $(d_0, \ldots, d_L)$ can be identified with a representation of the *linear quiver* A_{L+1} consisting of $L+1$ vertices connected by L consecutive arrows. In this identification, each layer ϕ_ℓ corresponds to an edge of A_{L+1} and the vector spaces k^{d_ℓ} to the vertices.

Over $k = \mathbb{R}$ (or $\mathbb{C}$) we fix inner products on each $V_v \cong k^{d_v}$ and endow each $\mathrm{Hom}(V_{s(e)}, V_{t(e)})$ with the Frobenius inner product

$$\langle \phi, \psi \rangle = \mathrm{Tr}(\phi^\top \psi).$$

Summing this over all edges defines an inner product on $\mathrm{Rep}_d(Q)$ and hence a Euclidean structure on the representation space.

Definition 2.7. Let $\phi^\star \in \mathrm{Hom}_k(k^{d_0}, k^{d_L})$ be a target linear map. The loss function of a DLN $\phi = \phi_L \cdots \phi_1$ is

$$\mathcal{L}((\phi_\ell)_{\ell=1}^L) = \|\phi^\star - \phi_L \cdots \phi_1\|_F^2,$$

where $\|\cdot\|_F$ denotes the Frobenius norm. We are particularly interested in the fiber $\mathcal{L}^{-1}(0)$.

Proposition 2.8. The representation space $\mathrm{Rep}_d(Q)$ is an affine variety with coordinate ring

$$k[\mathrm{Rep}_d(Q)] = k[x_{ij}^{(\ell)} \mid 1 \le \ell \le L,\, 1 \le i \le d_\ell,\, 1 \le j \le d_{\ell-1}],$$

and hence

$$\mathrm{Rep}_d(Q) \cong \mathbb{A}^{\sum_{\ell=1}^L d_{\ell-1} d_\ell}.$$

Proposition 2.9. The loss $\mathcal{L} : \mathrm{Rep}_d(Q) \to k$ is a polynomial map of degree $2L$. Its graph

$$\{(\Phi, \lambda) \in \mathrm{Rep}_d(Q) \times k \mid \lambda = \mathcal{L}(\Phi)\}$$

is an affine subvariety of dimension $\sum_{\ell=1}^L d_{\ell-1} d_\ell$.

Since $\mathrm{Rep}_d(Q)$ is isomorphic to an affine space, it carries a natural module of Kähler differentials $\Omega_{k[\mathrm{Rep}_d]/k}$, giving an algebraic notion of differential $d\mathcal{L}$. Over $\mathbb{R}$ or $\mathbb{C}$, once the Frobenius inner product is fixed, this identifies with the usual analytic gradient and the second differential with the Hessian of $\mathcal{L}$. This will enable us to study the critical points and local geometry of the DLN loss landscape.

3 Fibers of the multiplication map

This is a summary of Simon Pepin Lehalleur's paper on computing the codimension of the global minimum of the population loss of a deep linear network. The weight matrices of a DLN defines a representation of a Quiver. We have an action of the group $GL_{\bar{d}} := \prod_i GL_{d_i}$ over $Rep_{\bar{d}}$. We can then partition $Rep_{\bar{d}}$ into its orbits under the action of $G_d := GL_{\bar{d}}$. Recall that we have an isomorphism between Rep_d and kQ-modules. The isomorphism classes of kQ-modules of dimension $\bar{d}$ are isomorphic to the orbits of Rep_d under the action of G_d. The loci of zero loss is defined by:

$$\pi^{-1}(0) \subset Rep_{\bar{d}} \text{ with } \pi(\phi) = \phi_L ... \phi_1$$

We are interested by relating the geometry of $\pi^{-1}(0)$ with the combinatorics of the ranks of products of weight matrices. Note that we have:

$$\pi^{-1}(0) = \Sigma_d^0 = \{\phi \in Rep_d | rk(\phi) = 0\}$$

More generally, define:

$$\Sigma_d^r = \{\phi \in Rep_d | rk(\phi) = r\}$$

Given that the map π is equivarient w.r.t to G, we can decompose Σ_d^r into orbits under the action of G. This is true because we have a group action of G induced on $\mathrm{Im}(\pi)$ by the equivarience relation. An important result is that we can parametrize the G-orbits of Rep_d via the multiplicity pattern defined by:

$$\mathcal{M}_d^+ := \{(m_{ij})_{1 \leq i \leq j \leq N} | m_{ij} \in \mathbb{N}, d_k = \sum_{i \leq k \leq j} m_{ij}\}$$

The multiplicity patterns parametrize the isomorphic classes of the of the Dynkin Quiver $\mathbb{A}_n$-modules whose indecomposables representaions are given by (following from Gabriel's Theorem):

$$M_{ij} := 0 \to ... \to 0 \to k \to ... \to k \to 0... \to 0$$

By Krull-Schmidt every Q-module writes:

$$M = \oplus_{1 \leq i \leq j \leq N} m_{ij} M_{ij}$$

The rank patterns is another important object that enables to parametrize the orbits of Rep_d as well in the particular case of the A_n quiver. More specifically we are interested in the following orbit:

$$\mathcal{O}_{\bar{r}} := \{\phi | rk(\phi_j ... \phi_{i+1} = r_{ij}\}$$

With:

$$r_{ij} \in \mathcal{R}_d := \{(r_{ij})_{1 \leq i \leq j \leq N} | r_{ij} \in \mathbb{N}, r_{ij} - r_{i,j+1} - r_{i-1,j} + r_{i-1,j+1} \geq 0, r_{ii} = d_i\}$$

The rank pattern set $\mathcal{R}_d$ is in bijection with the multiplicity patterns and therefore parametrize the orbit of Rep_d as well. In other words, the G-orbits of Rep_d corresponds to rank condition on a composition of weight matrices in a DLN. We can therefore stratify the space Σ_d^r by the orbits $\mathcal{O}_{\bar{r}}$ parametrized by the rank patterns. This allows to relate the codimension of 0-loss with the codimension of the orbits which can be encoded in a power series using Pochammer symbols.

4 Loss landscape of DLNs

4.1 Equivariance of the gradient map

TODO: Recall main results on local minima being global minima, recall 1st and 2nd order analysis of critical points. Consider a DLN. We assume that the teacher matrix has distinct singular values, the output space is lower dimensional than the input space and we are in the overparametrize regime. Input-output covariance matrix is full rank and input covariance matrix is invertible. The paper on the Loss Landscape of Deep Linear Neural Networks parametrize the first order critical points with the subset of the singular values of the teacher matrix.

Proposition 4.1. Let $\phi \in \mathrm{Rep}_d$ be a first order critical point of the loss. Let $M = U\Sigma V^\top$ be the teacher matrix, let $r := rk(\pi(\phi))$, let $P_S = U_S U_S^\top$ be a projector along the top r singular values of the teacher matrix M then there exists a unique subset $S \subset \{1, ..., d_{out}\}$ of size r such that:

$$\pi(\phi) = \phi_L ... \phi_1 = P_S M$$

If ϕ is a first order critical point then it is given by some subset of singular values of the teacher matrix. However, not all $P_S M$ are critical points. So we cannot use $P_S M$ as a parametrization of critical points. In other words we have:

$$\mathrm{Crit}(\mathcal{L}) \subset \pi^{-1}(P_S M)$$

Let $G := G_{d_0} \times G_h \times G_{d_L}$ acting on Rep_d via:

$$g.\phi = (g_1 \phi_1 g_0^{-1}, ..., g_L \phi_L g_{L-1}^{-1})$$

The group $G_h = \prod_{0<i<L} G_{d_i}$ acts on Rep_d as a subgroup of G. Let $\pi_{>l} : Rep_d \to \mathrm{Hom}(k^{d_l}, k^{d_L})$ and $\pi_{<l} : Rep_d \to \mathrm{Hom}(k^{d_0}, k^{d_l})$ be the product maps:

$$\pi_{>l}(\phi) = \phi_L ... \phi_{l+1}$$
$$\pi_{<l}(\phi) = \phi_{l-1} ... \phi_1$$

The multiplication map π is G-equivariant and we can show that the product maps $\pi_{>l}$ and $\pi_{<l}$ are G-equivariant:

$$\pi_{>l}(g.\phi) = g_L \phi_L ... \phi_{l+1} g_l^{-1} = g_L \pi_{>l}(\phi) g_l^{-1} \in \mathrm{Hom}(k^{d_l}, k^{d_L})$$
$$\pi_{<l}(g.\phi) = g_{l-1} \phi_{l-1} ... \phi_1 g_0^{-1} = g_{l-1} \pi_{<l}(\phi) g_0^{-1} \in \mathrm{Hom}(k^{d_0}, k^{d_{l-1}})$$

Let

$$E(\phi) := \pi(\phi^\star) - \pi(\phi)$$

where $\phi^\star$ is a global minimum of the loss in Rep_d. Let $G_{\phi^\star}$ be the stabilizer of the target map $\pi(\phi^\star)$ in $GL_{d_L} \times GL_{d_0}$. In other words,

$$G_{\phi^\star} := \{(g_L, g_0) \in GL_{d_L} \times GL_{d_0} |\ g_L \pi(\phi^\star) g_0^- 1 = \pi(\phi^\star)\}$$

The group $G := G_{\phi^\star} \times G_h$ acting on Rep_d. The error term E is $G_{\phi^\star}$-equivariant because $\pi(\phi)$ is G-equivariant and $\pi(\phi^\star) \in Rep_d$ is $G_{\phi^\star}$-invariant. In other words we have:

$$\pi(g.\phi^\star) = g_L \pi(\phi^\star) g_0^{-1} = \pi(\phi^\star), \text{ for } (g_L, g_0) \in G_{\phi^\star}$$

The gradient at vertex l writes:

$$\nabla_l L(\phi) = \pi_{>l}(\phi)^t E(\phi) \pi_{<l}(\phi)^t$$

In particular we can define the gradient map at vertex l, $\nabla_l \in \mathrm{Hom}(Rep_d, \mathrm{Hom}(k^{d_{l-1}}, k^{d_l}))$:

$$\nabla_l : \phi \mapsto \pi_{>l}(\phi)^t E(\phi) \pi_{<l}(\phi)^t$$

Which induces a gradient map

$$\nabla := (\nabla_l)_l \in \bigoplus_l \mathrm{Hom}(Rep_d, \mathrm{Hom}(k^{d_{l-1}}, k^{d_l})) \cong \mathrm{Hom}(Rep_d, Rep_d)$$

note that Rep_d^* is a representation of the dual quiver with all the orientation of arrows from Q inverted[2]. The space $\mathcal{C}_L$ is the set of critical points i.e. points in Rep_d at which the map ∇ vanishes.

[2]Maybe the double quiver will be useful?

Proposition 4.2. Let $O_i := O(d_i)$ be the orthogonal group acting on vertex i. Let

$$O_{\phi^\star} := G_{\phi^\star} \cap (O_L \times O_0)$$

and let $OG := O_{\phi^\star} \times G_h$ be a group acting on Rep_d. The map ∇ is equivariant under the action of OG.

Proof We use the OG-equivariance of both the multiplication map π and the error map E and find that:

$$\nabla_l(L(g.\phi)) = g_l^{-t}\pi_{>l}(\phi)^t g_L^t g_L E(\phi) g_0^{-1} g_0^{-t}\pi_{<l}(\phi)^t g_{l-1}^t$$

Since $g_L \in O_L$ and $g_0 \in O_0$ then we have: $g_L^t g_L = 1,\ g_0^t g_0 = 1$ and thus:

$$\begin{aligned}
\nabla_l(L(g.\phi)) &= g_l^{-t}\nabla_l L(\phi) g_{l-1}^t, \forall l \\
\nabla L(g.\phi) &= g * \nabla L(\phi) \quad \text{Action of OG defined by:} \\
g * \nabla L(\phi) &= (g_l^{-t}\nabla_l L(\phi) g_{l-1}^t)_l
\end{aligned}$$

In particular thanks to the latter equality we see that if $\phi \in Rep_d$ is a critical point of L then for $g \in OG$ we have that $g.\phi$ is also a critical point of L. This defines a group action of OG over $\mathcal{C}_L$. Since we have a group action of OG over $\mathcal{C}_L$ there is a stratification of the critical points of L into orbits of OG:

$$\mathcal{C}_L := \bigsqcup_{\phi \in \mathcal{C}_L} \mathcal{O}_\phi$$

We want to find a parametrization of these orbits that could enable us to study the geometry of critical points (for example computing their codimension and studying their irreducible components) similar to what has been done previously on the fibers of ϕ (Simon's work).

4.2 Quasi-Parameterization of saddle points

Recall proposition 9 and 10 from the paper The Loss Landscape of Deep Linear Neural Networks: a Second-order Analysis.
Assumptions:

- Input data dimension d_x and output data dimension d_y are such that $d_y \leq d_x \leq m$
- Input data covariance Σ_X is invertible and input-output data covariance Σ_{YX} is full rank d_y
- The singular values of $\sigma = \Sigma_{YX}\Sigma_X^{-1}X$ are all distinct

Proposition 9 (reformulated): Let ϕ be a critical point. Let $S \subset \{1, ...d_y\}$ be a set indexing the singular values of the target map. Let $r := |S|$ be the rank of ϕ, proposition 9 ensures that ϕ is conjugate to the point $\tilde{\phi}$ in Rep_d with:

$$\begin{aligned}
\pi(\tilde{\phi}) &= \tilde{\phi}_L ... \tilde{\phi}_1 = U_S U_S^\top \Sigma_{YX}\Sigma_X^{-1}, \quad \text{the map } U_S U_S^\top \text{ projects on singular values indexed by } S \\
\tilde{\phi}_L &= [U_S\ U_Q Z_L] \quad \exists Z_L \in \mathbb{R}^{d_y - r \times d_{L-1} - r} \text{ and } Q = \{1...d_y\} - S \\
\forall h \in \{L-1, ..., 2\},\ \tilde{\phi}_h &= I_r \oplus Z_h, \quad \exists Z_h \in \mathbb{R}^{d_h - r \times d_{h-1} - r} \\
\tilde{\phi}_1 &= [U_S^\top \Sigma_{YX}\Sigma_X^{-1}\ Z_1]^\top \exists Z_1 \in \mathbb{R}^{d_1 - r \times d_0 - r} \\
U_Q Z_L ... Z_2 &= 0
\end{aligned}$$

Let $\tilde{\phi_Z}$ be a representation Q satisfying the conditions of the proposition just above. Define:

$$U^S_{\tilde{\phi_Z}} = \{\phi \in Rep_d | \phi \sim \tilde{\phi_Z}\}$$

In particular, we see that the set of critical point of rank $r = |S|$ associated with the subset of singular values indexed by S satisfies:

$$\mathcal{C}_{L,S} \subset \cup_Z U^S_{\tilde{\phi_Z}} \subset \pi^{-1}(U_S U_S^\top \Sigma_{XY} \Sigma_X^{-1})$$

Proposition 10 (reformulated): Consider ϕ such that ϕ is conjugate to the representation $\tilde{\phi}$ such that:

$$\begin{aligned} \tilde{\phi}_L &= [U_S \ U_Q Z_L] \quad \exists Z_L \in \mathbb{R}^{d_y - r \times d_{L-1} - r} \text{ and } Q \in \{1...d_y\} - S \\ \forall h \in \{L-1, ..., 2\}, \ \tilde{\phi}_h &= I_r \oplus Z_h, \quad \exists Z_h \in \mathbb{R}^{d_h - r \times d_{h-1} - r} \\ \tilde{\phi}_1 &= [U_S^\top \Sigma_{XY} \Sigma_X^{-1} \ Z_1]^\top \exists Z_1 \in \mathbb{R}^{d_1 - r \times d_0 - r} \end{aligned}$$

if $r = r_{max}$ or if there exists $h_1 \neq h_2$ with $Z_{h_1} = Z_{h_2} = 0$ then ϕ is a first order critical point of the loss associated with the subset S.
Let ϕ be a representation Q satisfying the conditions of the proposition just above. Define:

$$U^S_{h_1,h_2} = \{\phi \in Rep_d | \phi \sim \tilde{\phi} \text{ and } \exists h_1 \neq h_2 | Z_{h_1} = Z_{h_2} = 0\}$$

Then we have:

$$U^S_{h_1,h_2} \subset \mathcal{C}_{L,S} \subset \cup_Z U^S_{\tilde{\phi_Z}} \subset \pi^{-1}(U_S U_S^\top \Sigma_{XY} \Sigma_X^{-1})$$

Unfortunately, a full parameterization of the first order critical points is lacking. This is because we could find a critical point that does not fall into $U^S_{h_1,h_2}$ for example or some representation in $U^S_{\tilde{\phi}}$ that is not a critical point.
Recall that a critical point satisfies:

$$\pi_{>l}(\phi)^t E(\phi) \pi_{<l}(\phi)^t = 0$$

Let $\phi^\star := \Sigma_{XY}\Sigma_X^{-1}$ be the target map and let $P_S := U_S U_S^\top$ be a projector on singular values indexed by S. Using proposition 9 we can write this equation as:

$$\pi_{>l}(\phi)^t (I - P_S) \phi^\star \pi_{<l}(\phi)^t = 0$$

Let $P_S^\perp := 1 - P_S$, unfortunately an GL_d group action on the map:

$$\psi_S(\phi) := \pi_{>l}(\phi)^t P_S^\perp \phi^\star \pi_{<l}(\phi)^t$$

allows to define a group of GL_d over the critical points if only if

$$\pi_{>l}(\phi)^t g_L^{-t} P_S^\perp \phi^\star g_0^{-t} \pi_{<l}(\phi)^t = 0$$

Which does not seem to hold in general.
Proposition The gradient is equivariant under the action of G_h and critical points are G_h invariant.
Proof We have already shown that the gradient map satisfies:

$$g * \nabla_l L(\phi) = g_l^t \nabla_l L(\phi) g_{l-1}^t = \nabla_l L(g.\phi)$$

Let ϕ_S be a critical point i.e. we have

$$\nabla L(\phi_S) = 0$$

Acting on ϕ_S via the hidden subgroup G_h yields:

$$\nabla_l L(g\phi_S) = g_l^t \nabla_l L(\phi_S) g_{l-1}^t = 0$$

Since $\nabla_l L(\phi_S) = 0$ for all l.

5 Hessian analysis

The Hessian of the loss function encodes the local curvature of the parameter space and therefore provides a second-order description of the geometry of the optimization landscape. Its eigenvalues quantify curvature along different parameter directions: large eigenvalues correspond to stiff, highly identifiable directions, while small or zero eigenvalues indicate flat or redundant directions. The rank of the Hessian thus measures the effective local dimensionality of the model and reveals structural degeneracies induced by overparameterization or symmetries. In regular models, the Hessian is full rank and the local learning coefficient (LLC) equals half the parameter dimension. In singular models, however, the presence of degenerate Hessian directions reflects non-identifiability and constrains the asymptotic behavior of the marginal likelihood; in particular, the Hessian rank provides an upper bound on twice the LLC and characterizes the quadratic part of the loss that determines the leading-order contribution to the real log canonical threshold.

5.1 Hessian Decomposition: Outer-Product and Functional Terms

Consider a parametric model $f_\theta(x) \in \mathbb{R}^K$ with loss

$$L(\theta) = \mathbb{E}_{(x,y)}\big[\ell(f_\theta(x), y)\big].$$

Let $J_\theta(x) := \nabla_\theta f_\theta(x) \in \mathbb{R}^{K\times p}$ denote the Jacobian of the model output with respect to the parameters, and let $\nabla_f \ell$ and $\nabla_f^2 \ell$ denote the gradient and Hessian of the loss with respect to the model output f. Applying the chain rule twice yields the Gauss–Newton decomposition of the parameter Hessian:

$$\nabla_\theta^2 L(\theta) = H_o(\theta) + H_f(\theta),$$

where

$$H_o(\theta) = \mathbb{E}\big[J_\theta(x)^\top \nabla_f^2 \ell(f_\theta(x), y)\, J_\theta(x)\big]\,,$$

and

$$H_f(\theta) = \mathbb{E}\left[\sum_{k=1}^{K} \frac{\partial \ell}{\partial f_k}(f_\theta(x), y)\, \nabla_\theta^2 f_{\theta,k}(x)\right].$$

Outer-product term. The matrix H_o depends only on first derivatives of the model and second derivatives of the loss. It measures how curvature in output space is pulled back to parameter space through the Jacobian. When $\nabla_f^2 \ell \succeq 0$ (e.g., mean squared error or cross-entropy), H_o is positive semidefinite. For mean squared error,

$$\ell(f, y) = \tfrac{1}{2}\|f - y\|^2, \qquad \nabla_f^2 \ell = I_K,$$

and therefore

$$H_o(\theta) = \mathbb{E}\big[J_\theta(x)^\top J_\theta(x)\big].$$

In this case, H_o coincides with the Gauss–Newton matrix and is closely related to the empirical Fisher and the Neural Tangent Kernel. Its rank is bounded by the rank of the Jacobian and thus reflects the locally identifiable directions supported by the data.

Functional term. The matrix H_f depends on second derivatives of the model itself. It captures curvature induced by the parameterization rather than by the loss geometry. For mean squared error,

$$H_f(\theta) = \mathbb{E}\left[\sum_{k=1}^{K} \big(f_{\theta,k}(x) - y_k\big)\, \nabla_\theta^2 f_{\theta,k}(x)\right].$$

Hence H_f is weighted by residuals and vanishes at exact interpolation. Unlike H_o, it is generally indefinite and encodes structural degeneracies arising from overparameterization or symmetries in the model.

Geometric interpretation and relation to the LLC. The decomposition separates two sources of curvature: data-induced curvature (H_o) and parameterization-induced curvature (H_f). Near well-fitted minima in regression and classification, H_f becomes negligible and the Hessian is dominated by H_o. The nonzero eigenvalues of H_o determine the quadratic directions of the loss landscape and therefore govern the leading-order local geometry. In regular models the Hessian is full rank, whereas in singular models rank deficiency reflects parameter redundancies. Since the local learning coefficient (LLC) is determined by the leading quadratic structure of the loss expansion, the rank of the Hessian provides an upper bound on twice the LLC, while higher-order terms determine whether this bound is tight.

5.2 Exact Rank Formula, Upper Bound, and Rank Deficiency

We consider a depth-L deep linear network with layer widths $d_0, d_1, \ldots, d_L$, where $d_0 = d$ is the input dimension and $d_L = K$ the output dimension. Let $M_\ell = d_\ell$ and let

$$M := \sum_{\ell=1}^{L-1} M_\ell$$

denote the total number of hidden units. Let $\Sigma_{xx} = \mathbb{E}[xx^\top]$ with $r := \mathrm{rk}(\Sigma_{xx})$, and define

$$q := \min\{r, M_1, \ldots, M_{L-1}, K\}.$$

Exact expression for the outer-product term. For the mean squared error loss, the outer-product (Gauss–Newton) term admits the factorization

$$H_o = A_o(I_K \otimes \Sigma_{xx})A_o^\top,$$

where

$$A_o^\top = \big(W_{L:2} \otimes I_d,\, W_{L:\ell+1} \otimes W_{1:\ell-1},\, \ldots,\, I_K \otimes W_{1:L-1}\big).$$

Under the maximal-rank assumption on the weight matrices, the rank of H_o is given exactly by

$$\mathrm{rk}(H_o) = q\,(r + K - q).$$

Upper bound (tight in practice) for the full Hessian. Writing $H_L = H_o + H_f$, one obtains the sharp upper bound

$$\mathrm{rk}(H_L) \leq 2qM - Lq^2 + q(r + K).$$

Empirically (and at initialization), this bound holds with equality:

$$\mathrm{rk}(H_L) = 2qM - Lq^2 + q(r + K).$$

Rank deficiency. Let $p = \sum_{\ell=0}^{L-1} M_\ell M_{\ell+1}$ be the total number of parameters. When $r = d$ (full-rank input covariance), the rank deficiency of the Hessian satisfies

$$\text{rank-deficiency}(H_L) = p - \mathrm{rk}(H_L) = \sum_{\ell=0}^{L-1} (M_\ell - q)(M_{\ell+1} - q).$$

Interpretation. These formulas reveal that the effective dimension of the model, as measured by the Hessian rank, scales *linearly* in the layer widths, whereas the number of parameters scales *quadratically*. The quantity q acts as the effective bottleneck dimension: only q directions per layer participate in curvature. All remaining directions correspond to structurally degenerate parameter combinations.

The rank deficiency formula admits a particularly clean interpretation: it equals the parameter count of a hypothetical network in which each layer width has been reduced by q. In other words, the degeneracy is exactly the redundancy created by overparameterizing beyond the minimum effective dimension.

Consequently, deep networks possess a large nullspace in the Hessian, whose dimension grows quadratically with overparameterization. The Hessian therefore lives on a much lower-dimensional manifold than the parameter space itself. This clarifies that: (i) degeneracy is architectural rather than accidental, (ii) the effective dimension is governed by q and the sum of widths, not by the raw parameter count, and (iii) overparameterization primarily increases flat directions rather than curved ones.

5.3 Link Between H_o, H_f, and Strict vs. Non-Strict Saddles

Consider a deep linear network trained with squared loss

$$L(W) = \frac{1}{2}\|Y - W_{L:1}X\|_F^2.$$

The Hessian admits the Gauss–Newton decomposition

$$H_L = H_o + H_f,$$

where

$$H_o = \mathbb{E}\big[J^\top J\big] \quad \text{and} \quad H_f = \mathbb{E}\left[\sum_{k=1}^{d_L} \frac{\partial \ell}{\partial f_k} \nabla_W^2 f_k\right].$$

For squared loss, H_o is positive semidefinite, since it is a Gram matrix of Jacobians. Hence any negative curvature of the full Hessian must arise from the functional term H_f.

Second-order expansion and correspondence. In the second-order analysis of Achour *et al.*, the loss along a perturbation W' around a first-order critical point is expanded as

$$L(W + tW') = L(W) + t^2 c_2(W, W') + o(t^2),$$

where the quadratic coefficient $c_2(W, W')$ splits into two parts:

$$c_2(W, W') = F^T(W, W') + S^T(W, W').$$

The term F^T is a squared-norm expression (of the form $\| \sum_i T_i \|^2$) and is manifestly nonnegative. This corresponds precisely to the Gauss–Newton curvature $v^\top H_o v$. The term S^T is a residual-weighted interaction term arising from second derivatives of the model; this corresponds to the functional contribution $v^\top H_f v$.

Thus, in quadratic form,

$$v^\top H_L v = v^\top H_o v + v^\top H_f v,$$

which matches

$$c_2(W, W') = F^T + S^T.$$

Strict vs. non-strict saddles. Since $H_o \succeq 0$, negative curvature of H_L can only occur if

$$v^\top H_f v < -v^\top H_o v$$

for some direction v. In other words, the functional term must be sufficiently negative to dominate the positive semidefinite outer-product curvature.

The results of Achour *et al.* show that

- If a critical point is *non-tightened*, then there exists a direction W' such that $c_2(W, W') < 0$. In Hessian language, this means H_f produces a negative eigenvalue of H_L, and the point is a strict saddle.
- If a critical point is *tightened*, the second-order term can be rewritten as a sum of squares, implying $c_2(W, W') \geq 0$ for all directions. Equivalently, H_f cannot overcome the PSD curvature of H_o, and the Hessian has no negative eigenvalues (non-strict saddle).

Interpretation. The Gauss–Newton term H_o captures curvature induced by output sensitivity and is always positive semidefinite. The functional term H_f captures residual-weighted second-order effects of the parameterization and is the sole source of indefiniteness. Non-tightened configurations correspond to structural conditions under which the indefinite component H_f generates a negative curvature direction. Tightened configurations correspond to algebraic rank conditions that prevent H_f from dominating H_o, thereby eliminating strict saddle behavior.
Useful references: Analytic Insights into Structure and Rank of Neural Network Hessian Maps and Depth, Not Data: An Analysis of Hessian Spectral Bifurcation

6 Bounding the critical loci

Using proposition 9, we see that for all the critical points the product map is of the form:

$$\pi\phi = \phi_L \dots \phi_1 = P_S \Sigma_{XY} \Sigma_{XX} + U_Q Z_L \dots Z_1, \text{ with}$$
$$U_Q Z_L \dots Z_1 = 0$$

Let $P_Q = U_Q U_Q^\top$ a projector onto the singular values not in S (i.e. orthogonal to P_S in some sense). We could study the multiplicative map:

$$\pi_Q(Z_L, ..., Z_1) = U_Q Z_L \dots Z_1$$

And focus on $\pi_Q^{-1}(0)$. Indeed, for all critical points the latter equation holds, so perhaps by studying the fibers of this map we can study the geometry of orbits in that fiber that "upper bound" (in the sense of inclusion) the loci of saddle points. We have an action of the group G such that:

$$\pi_Q g.(Z_L, ..., Z_1) = U_Q g_L Z_L \dots Z_1 g_0^{-1} = g_L U_Q g_L^t g_L Z_L \dots Z_1 g_0^{-1} = g.\pi_Q(Z_L, ..., Z_1)$$

This holds if $g_L \in O_L$ commutes with U_Q which implies that P_Q is a fixed point for g_L? Is there an interesting subgroup of O_L that has this property? Another point: in terms of representations of the quiver Q, the Z_i are sub representations of the representation induced by ϕ_i. **Relations between projectors on singular values** One can show that:

- $U_S U_S^\top$ is a projector on r singular values and $U_Q U_Q^\top$ is a projector on $d_y - r$ singular values
- $U_S^\top U_S = 1$ and similar for U_Q
- The SVD of $P_S M$ is $U S_r V^\top$ where S_r is a matrix with singular values $s_1, ..., s_r$ and 0 otherwise
- The SVD of $P_Q M$ is $U S_{d_y - r} V^\top$ where S_{d_y-r} is a matrix with singular values $s_{r+1}, ..., s_{d_y-r}$ and 0 otherwise
- $U_S^\top U_Q = U_Q^\top U_S = 0$

This can be shown by noting that $U = [U_S,\ U_Q]$ up to re-ordering of singular values which does not matters since SVD is invariant under permutations of singular vectors. Using $U^\top U = 1$ we get the orthogonality relations between U_S and U_Q. Since we have $U_Q Z_L ... Z_1 = 0$ then using $U_Q^\top U_Q = 1$ we find this implies $Z_L ... Z_1 = 0$. Hence we have $U_Q Z_L ... Z_1 = 0$ if and only if $Z_L ... Z_1 = 0$. We can therefore study the fiber of the multiplication map

$$\pi_Z(Z_L, ..., Z_1) := Z_L ... Z_1$$

To bound the critical point above (in the sense of inclusion).

Restriction to fibers of the residuals

To study the critical loci of the DLN loss landscape, it would be useful to study the set of points that upper bound and lower bound that loci (in the sense of the inclusion of sets). Recall that from proposition 1 we have for ϕ_c a critical point: $\pi(\phi_c) = U_S U_S^\top M$. The following general fact will be important:

Proposition 6.1. Let U be an orthonormal matrix in O_{d_y}. Let U_S be a matrix defined by taking the first $r < d_y$ columns of U and let U_Q be a matrix defined by taking the remaining columns of U. The matrices U_S and U_Q have orthonormal columns and are mutually orthogonal. In particular, the matrix U_Q satisfies:

$$U_Q^\top U_Q = 1_{d_y - r}$$

.

To prove that proposition, up to re-ordering of the singular values, we can generally write the left singular matrix U as:

$$U = [U_S,\ U_Q]$$

Since U is an orthogonal matrix in O_{d_y} we have

$$\begin{aligned} U^\top U &= 1_{d_y} \\ &= \begin{pmatrix} U_S^\top U_S & U_S^\top U_Q \\ U_Q^\top U_S & U_Q^\top U_Q \end{pmatrix} \end{aligned}$$

Hence by identifying the matrix blocks we obtain:

$$\begin{aligned} U_S^\top U_S &= 1_r \\ U_Q^\top U_Q &= 1_{d_y - r} \\ U_S^\top U_Q &= 0_{r, d_y - r} \\ U_Q^\top U_S &= 0_{d_y - r, r} \end{aligned}$$

Which terminates the proof.
Consider a representation $\phi \in Rep_d$ satisfying proposition 9 or 10 reformulated above. We call a subrepresentation of ϕ defined by the maps $(Z_i)_i$ the residual representation. Recall that we have:

$$\pi_Q(Z_1, ..., Z_L) = U_Q Z_L ... Z_1$$

We have the following proposition:

Proposition 6.2. Let $\pi_Z(Z_1, ..., Z_L) = Z_L ... Z_1$ be the multiplication map of the residuals. We have:

$$\pi_Q(Z_1, ..., Z_L) = 0 \iff \pi_Z(Z_1, ..., Z_L) = 0$$

This proposition enables us to import the results from the fibers of the multiplicative map to π_Z to study the critical loci. Since we work under the conditions of proposition 9 and 10, this proposition is easy to prove by using the orthogonality of the columns of U_Q and $\pi_Q(Z_1, ..., Z_L) = 0$:

$$U_Q^\top \pi_Q(Z_1, ..., Z_L) = U_Q^\top U_Q \pi_Z(Z_1, ..., Z_L) = \pi_Z(Z_1, ..., Z_L) = Z_L ... Z_1 = 0$$

Conversely, if $\pi_Z = 0$ then $\pi_Q = U_Q \pi_Z = 0$. More generally we have for critical points:

$$\begin{aligned} &U_S U_S^\top \phi = U_S U_S^\top M \\ &U_Q U_Q^\top \phi = U_Q U_Q^\top U_Q Z_L .. Z_1 = U_Q Z_L ... Z_1 = 0, \text{ if } \phi \text{ is a critical point} \end{aligned}$$

The set of critical points can therefore be studied via:

$$\{\phi \in Rep_d, Z \in Rep_{d-r} | \pi(\phi) = P_S M + U_Q \pi_Z(Z), \pi_Z(Z) = 0\}$$

7 Saddle points for degenerate data

Consider proposition 1 from the paper "The Loss Landscape of Deep Linear Neural Networks: a Second-order Analysis". The assumption that the matrix Σ has distinct eigenvalues is essential to prove that all critical points are some projection of the global optimum onto a subset of singular values and that they write as a diagonal matrix in the same basis that diagonalizes Σ. Proposition 1 is not true when assuming that the teacher matrix has degenerate eigenvalues. In that case a saddle point still seems to be a projection of the teacher matrix but not necessarily along its singular vector: it projects onto a basis that can be rotated to the singular basis of the teacher matrix.

For a counter example, consider the teacher matrix $M = I_2$, assume that $\Sigma_{XY} = I_2 = \Sigma_X$ i.e. whitened inputs. Consider a two layers DLN with whitened inputs. Consider the projector:

$$P = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

Consider the point:

$$W_1 = W_2 = P$$

We see that the point (P, P) is a critical point by checking that gradient is zero and that we have $P^2 = P$ i.e. P is a projector. Further more the loss at P has value 1 which is not 0 i.e. P is not a global minimum but a saddle point. We have found a counter-example to proposition 1 when relaxing the assumption of data degeneracy.

We propose a conjecture for an extension of the first statement of proposition 1 when the data has some degenerate singular values. We relax the assumption that the matrix

$$\Sigma = \Sigma_{YX}\Sigma_X^{-1}\Sigma_{XY}$$

has distinct eigenvalues. Furthermore, recall that we have:

$$\Sigma = U\Lambda U^\top$$

Where U is the left singular matrix of $\Sigma^{\frac{1}{2}}$ and Λ is a diagonal matrix with eigenvalues λ_i each with multiplicities m_i. We conjecture the following:

Conjecture: Let W be a first order critical point, let I_S be a block diagonal matrix of identities I_{m_i} if $i \in S$ and 0 otherwise, then there exist a subset S indexing the distinct eigenvalues of Σ and a projector P and a orthogonal matrix Q such that:

$$W = P\Sigma_{XY}\Sigma_X^{-1} = UQI_SQ^\top U^\top \Sigma_{XY}\Sigma_X^{-1}$$

The projector P satisfies:

$$P\Sigma = \Sigma P$$

The matrix Λ is diagonal with eigenvalues of multiplicities m_i. The matrix Q is a block diagonal matrix for which each block is an orthogonal matrix in $O(m_i)$.

8 Geometry of the critical points

Define the variety of matrices of rank r:

$$\Sigma_d^r := \{W \in \Omega | rk(W_L ... W_1) = r\}$$

Consider the following group action of $GL_d = \Pi_{l=L}^0 GL_{d_l} = GL_{d_L} \times GL_h \times GL_{d_0}$:

$$(W_1, \dots, W_L) \mapsto (g_1 W_1 g_0^{-1},\ g_2 W_2 g_1^{-1},\ \dots,\ g_L W_L g_{L-1}^{-1})$$

We can also reparametrize Σ_d^r by the following:

$$\Sigma_d^r = \{W \in \Omega, Z_l \in \mathbb{R}^{d_l - r \times d_{l+1} - r} | W_l \sim \begin{pmatrix} I_r & 0 \\ 0. & Z_l \end{pmatrix}, Z_L ... Z_1 = 0\}$$

Note that the latter product over Z's can be defined as the manifold:

$$\Sigma_0^{d-r} = \{Z_l \in \mathbb{R}^{d_l - r \times d_{l+1} - r} | Z_L ... Z_1 = 0\}$$

Define the algebraic variety Γ by the following cyclic conditions:

$$\Gamma = \{Z_l \in \mathbb{R}^{d_l - r \times d_{l+1} - r} \mid Z_{l-1} \cdots Z_1\, \Sigma_{YX} U_Q\, Z_L \cdots Z_{l+1} = 0\}$$

Equivalently, we can view Γ as a space of quiver representations with relations. Let Q_Γ be the cyclic quiver with vertex set $\{0, \dots, L\}$ and arrows

$$a_l : l - 1 \to l \quad (l = 1, \dots, L), \qquad a_0 : L \to 0,$$

together with an additional distinguished arrow

$$a_0 : L \to 0,$$

A representation of Q_Γ is given by vector spaces $V_l \cong \mathbb{R}^{d_l - r}$ and linear maps $\phi_l : V_{s(a_l)} \to V_{t(a_l)}$. We identify

$$\phi_l := Z_l \ (l = 1, \dots, L), \qquad \phi_0 := \Sigma_{XY} U_Q : V_L \to V_0,$$

(i.e. ϕ_0 is specialized/fixed to the linear map $\Sigma_{XY} U_Q$). Then Γ is the subset of such representations satisfying the L cyclic relations

$$\phi_{l-1} \cdots \phi_1\, \phi_0\, \phi_L \cdots \phi_{l+1} = 0, \qquad l = 1, \dots, L,$$

with the convention that an empty product is the identity. If we include the condition that

$$\phi_L ... \phi_1 = 0$$

Then all paths of length L are zero and we denote Γ_d the set of paths representations satisfying this condition. From the 2nd order analysis paper, the set of critical points write:

$$\mathcal{C}_r = \{W \in \Omega, Z_l \in \mathbb{R}^{d_l - r \times d_{l+1} - r} | W_L = [U_S, U_Q Z_L], W_l \sim \begin{pmatrix} I_r & 0 \\ 0. & Z_l \end{pmatrix}, W_1 = [U_S^\top \Sigma_{YX} \Sigma_X^{-1}, Z_1]^\top, Z_l \in \Gamma_d\}$$

We want to show something like:

$$\mathcal{C}_r / G_h \cong \Gamma_d$$

This suggests that studying the codimension of the critical loci amounts to studying the codimensions of Γ_d.

8.1 Link between critical values and cyclic conditions

Consider the following subset of parameters:

$$\Omega_r := \{W \in \Omega | W_l \sim \begin{pmatrix} I_r & 0 \\ 0. & Z_l \end{pmatrix}, Z_l \in \mathbb{R}^{(d_l - r)\times(d_{l+1}-r)}\}$$

8.2 A block-rank stratum and an associated bundle model

Definition 8.1 (Block-rank stratum Ω_r)**.** Fix integers $0 \le r \le \min_{0\le\ell\le L} d_\ell$ and let

$$\Omega \;:=\; \prod_{\ell=1}^{L} \mathrm{Mat}_{d_\ell \times d_{\ell-1}}(\mathbb{R})$$

be the parameter space of depth-L linear networks. Let

$$G \;:=\; \prod_{\ell=0}^{L} GL(d_\ell, \mathbb{R})$$

act on Ω by the standard change-of-basis (gauge) action

$$(g \cdot W)_\ell \;:=\; g_\ell \, W_\ell \, g_{\ell-1}^{-1}, \qquad g = (g_0, \ldots, g_L) \in G,\; W = (W_1, \ldots, W_L) \in \Omega. \tag{1}$$

For each ℓ define the residual matrix space

$$\mathrm{Mat}_\ell^{(r)} \;:=\; \mathrm{Mat}_{(d_\ell - r)\times(d_{\ell-1}-r)}(\mathbb{R}), \qquad \mathcal{Z} \;:=\; \prod_{\ell=1}^{L} \mathrm{Mat}_\ell^{(r)}.$$

Define the embedding $\iota : \mathcal{Z} \to \Omega$ by

$$\iota(Z)_\ell \;:=\; \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} \in \mathrm{Mat}_{d_\ell \times d_{\ell-1}}(\mathbb{R}), \qquad Z = (Z_1, \ldots, Z_L) \in \mathcal{Z}, \tag{2}$$

where the block decomposition uses $\mathbb{R}^{d_{\ell-1}} \cong \mathbb{R}^r \oplus \mathbb{R}^{d_{\ell-1}-r}$ in the domain and $\mathbb{R}^{d_\ell} \cong \mathbb{R}^r \oplus \mathbb{R}^{d_\ell - r}$ in the codomain.

Define the slice

$$S_r \;:=\; \iota(\mathcal{Z}) \;\subset\; \Omega.$$

Finally, define

$$\Omega_r \;:=\; G \cdot S_r \;=\; \{\, g \cdot \iota(Z) \;:\; g \in G,\; Z \in \mathcal{Z} \,\}. \tag{3}$$

Equivalently, $W \in \Omega_r$ iff there exist $g \in G$ and $Z \in \mathcal{Z}$ such that $W_\ell = g_\ell^{-1} \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} g_{\ell-1}$ for all ℓ.

Definition 8.2 (Stabilizer of the slice)**.** Let $H_r \le G$ be the stabilizer subgroup of the slice S_r, i.e.

$$H_r \;:=\; \{\, h \in G : h \cdot S_r = S_r \,\}.$$

Then H_r is the subgroup of block-diagonal elements of the form

$$H_r \;=\; \left\{ h = (h_0, \ldots, h_L) \in G \;:\; h_\ell = \begin{pmatrix} A & 0 \\ 0 & K_\ell \end{pmatrix},\; A \in GL(r, \mathbb{R}),\; K_\ell \in GL(d_\ell - r, \mathbb{R}) \right\}. \tag{4}$$

Proposition 8.3 (Associated bundle model for Ω_r)**.** Let Ω_r be as in Definition 8.1, and let H_r be as in Definition 8.2. Define an action of H_r on $G \times \mathcal{Z}$ by

$$(g, Z) \cdot h := \big(gh,\ h^{-1} \cdot Z\big), \tag{5}$$

where for $h \in H_r$ written as in (4),

$$(h^{-1} \cdot Z)_\ell := K_\ell^{-1}\, Z_\ell\, K_{\ell-1}, \qquad \ell = 1, \dots, L, \tag{6}$$

(and A acts trivially on $\mathcal{Z}$). Let $G \times_{H_r} \mathcal{Z}$ denote the quotient of $G \times \mathcal{Z}$ by this right action.

Define

$$\Phi\colon G \times \mathcal{Z} \to \Omega, \qquad \Phi(g, Z) := g \cdot \iota(Z), \tag{7}$$

with ι as in (2) and the G-action as in (1). Then:

1. $\Phi(G \times \mathcal{Z}) = \Omega_r$.
2. Φ is H_r-invariant for the action (5), hence descends to a well-defined map

   $$\bar{\Phi}\colon G \times_{H_r} \mathcal{Z} \longrightarrow \Omega_r, \qquad \bar{\Phi}([g, Z]) = \Phi(g, Z).$$

   Moreover, $\bar{\Phi}$ is surjective.
3. Define the *principal stabilizer locus*

   $$\mathcal{Z}^\circ := \big\{ Z \in \mathcal{Z} \;\big|\; \mathrm{Stab}_G(\iota(Z)) = H_r \big\},$$

   i.e. the set of Z for which the only elements of G mapping $\iota(Z)$ back into S_r are the block-diagonal elements in H_r. Let $\Omega_r^\circ := \Phi(G \times \mathcal{Z}^\circ)$. Then the restriction

   $$\bar{\Phi}\colon G \times_{H_r} \mathcal{Z}^\circ \xrightarrow{\sim} \Omega_r^\circ$$

   is a bijection. In particular, on the principal stabilizer locus,

   $$\Omega_r^\circ \;\cong\; G \times_{H_r} \mathcal{Z}^\circ. \tag{8}$$

Proof. **Step 1 (Image equals Ω_r).** By definition (3),

$$\Omega_r = \{ g \cdot \iota(Z) : g \in G,\ Z \in \mathcal{Z} \}.$$

But $\Phi(g, Z) = g \cdot \iota(Z)$ by (7), hence $\Phi(G \times \mathcal{Z}) = \Omega_r$.

Step 2 (H_r-invariance). Take $(g, Z) \in G \times \mathcal{Z}$ and $h \in H_r$. Using (5)–(6),

$$\Phi\big((g, Z) \cdot h\big) = \Phi\big(gh,\ h^{-1} \cdot Z\big) = (gh) \cdot \iota(h^{-1} \cdot Z).$$

By the group action property (1),

$$(gh) \cdot \iota(h^{-1} \cdot Z) = g \cdot \big(h \cdot \iota(h^{-1} \cdot Z)\big).$$

Thus it suffices to show

$$h \cdot \iota(h^{-1} \cdot Z) = \iota(Z). \tag{9}$$

Write $h_\ell = \left(\begin{smallmatrix} A & 0 \\ 0 & K_\ell \end{smallmatrix}\right)$ as in (4). Then, using (1) and (2),

$$\big(h \cdot \iota(h^{-1} \cdot Z)\big)_\ell = h_\ell \begin{pmatrix} I_r & 0 \\ 0 & (h^{-1} \cdot Z)_\ell \end{pmatrix} h_{\ell-1}^{-1}.$$

Since $h_{\ell-1}^{-1} = \left(\begin{smallmatrix} A^{-1} & 0 \\ 0 & K_{\ell-1}^{-1} \end{smallmatrix}\right)$ and $(h^{-1} \cdot Z)_\ell = K_\ell^{-1} Z_\ell K_{\ell-1}$ from (6), we compute the block product explicitly:

$$\begin{pmatrix} A & 0 \\ 0 & K_\ell \end{pmatrix} \begin{pmatrix} I_r & 0 \\ 0 & K_\ell^{-1} Z_\ell K_{\ell-1} \end{pmatrix} \begin{pmatrix} A^{-1} & 0 \\ 0 & K_{\ell-1}^{-1} \end{pmatrix}.$$

Multiplying the first two matrices gives

$$\begin{pmatrix} A & 0 \\ 0 & K_\ell \end{pmatrix} \begin{pmatrix} I_r & 0 \\ 0 & K_\ell^{-1} Z_\ell K_{\ell-1} \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & Z_\ell K_{\ell-1} \end{pmatrix},$$

and multiplying by the third matrix yields

$$\begin{pmatrix} A & 0 \\ 0 & Z_\ell K_{\ell-1} \end{pmatrix} \begin{pmatrix} A^{-1} & 0 \\ 0 & K_{\ell-1}^{-1} \end{pmatrix} = \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} = \iota(Z)_\ell.$$

This proves (9), hence $\Phi((g, Z) \cdot h) = \Phi(g, Z)$ for all $h \in H_r$. Therefore Φ is H_r-invariant and descends to a well-defined map $\bar{\Phi} \colon G \times_{H_r} \mathcal{Z} \to \Omega_r$.

Step 3 (Surjectivity of $\bar{\Phi}$). By Step 1, every element of Ω_r has the form $\Phi(g, Z)$ for some (g, Z). Thus every element of Ω_r is in the image of $\bar{\Phi}$.

Step 4 (Injectivity on the principal stabilizer locus). Assume $\Phi([g, Z]) = \Phi([g', Z'])$ with $Z, Z' \in \mathcal{Z}^\circ$. Then $\Phi(g, Z) = \Phi(g', Z')$, i.e.

$$g \cdot \iota(Z) = g' \cdot \iota(Z'). \tag{10}$$

Let $h := g^{-1} g' \in G$. Applying g^{-1} to (10) and using the group action property gives

$$\iota(Z) = h \cdot \iota(Z'). \tag{11}$$

In particular, h maps the point $\iota(Z') \in S_r$ to the point $\iota(Z) \in S_r$.

We now use the assumption that $Z' \in \mathcal{Z}^\circ$, i.e. $\mathrm{Stab}_G(\iota(Z')) = H_r$. We claim that $h \in H_r$. To see this, note that equation (11) states that h maps $\iota(Z')$ into S_r. More precisely, we need to show that any $h \in G$ satisfying $h \cdot \iota(Z') \in S_r$ for some $Z' \in \mathcal{Z}^\circ$ must belong to H_r.

Write $h \cdot \iota(Z') = \iota(Z)$ for some $Z \in \mathcal{Z}$. Consider the one-parameter family $Z'(t) := Z' + t\,\delta Z$ for an arbitrary displacement $\delta Z \in \mathcal{Z}$, and define $W(t) := h \cdot \iota(Z'(t))$. Expanding component-wise,

$$W(t)_\ell = h_\ell \begin{pmatrix} I_r & 0 \\ 0 & Z'_\ell + t\,\delta Z_\ell \end{pmatrix} h_{\ell-1}^{-1}.$$

At $t = 0$, this equals $\iota(Z)_\ell$ by (11). The derivative at $t = 0$ is

$$\frac{d}{dt}\Big|_{t=0} W(t)_\ell = h_\ell \begin{pmatrix} 0 & 0 \\ 0 & \delta Z_\ell \end{pmatrix} h_{\ell-1}^{-1}.$$

Write h_ℓ in blocks compatible with the decomposition $\mathbb{R}^{d_\ell} = \mathbb{R}^r \oplus \mathbb{R}^{d_\ell - r}$:

$$h_\ell = \begin{pmatrix} A_\ell & B_\ell \\ C_\ell & K_\ell \end{pmatrix}, \qquad h_{\ell-1}^{-1} = \begin{pmatrix} \alpha_{\ell-1} & \beta_{\ell-1} \\ \gamma_{\ell-1} & \kappa_{\ell-1} \end{pmatrix}.$$

A direct computation shows

$$h_\ell \begin{pmatrix} 0 & 0 \\ 0 & \delta Z_\ell \end{pmatrix} h_{\ell-1}^{-1} = \begin{pmatrix} B_\ell\, \delta Z_\ell\, \gamma_{\ell-1} & B_\ell\, \delta Z_\ell\, \kappa_{\ell-1} \\ K_\ell\, \delta Z_\ell\, \gamma_{\ell-1} & K_\ell\, \delta Z_\ell\, \kappa_{\ell-1} \end{pmatrix}.$$

Now, $\iota(Z)_\ell$ has the block form $\left(\begin{smallmatrix} I_r & 0 \\ 0 & Z_\ell \end{smallmatrix}\right)$, so for $W(t)$ to remain in S_r (i.e. have the same block structure) to first order for *all* choices of δZ_ℓ, both off-diagonal blocks of the derivative must vanish for all δZ_ℓ. The $(1,2)$-block condition $B_\ell\, \delta Z_\ell\, \kappa_{\ell-1} = 0$ for all δZ_ℓ forces $B_\ell = 0$ or $\kappa_{\ell-1} = 0$; but $h_{\ell-1}$ is invertible, so $\kappa_{\ell-1} \neq 0$, hence $B_\ell = 0$. Similarly, the $(2,1)$-block condition gives $\gamma_{\ell-1} = 0$, which (since $h_{\ell-1}^{-1}$ invertible and $B_{\ell-1} = 0$ by induction) forces $C_{\ell-1} = 0$ as well.

Applying this argument at every layer ℓ shows that each h_ℓ is block-diagonal:

$$h_\ell = \begin{pmatrix} A_\ell & 0 \\ 0 & K_\ell \end{pmatrix},$$

i.e. $h \in H_r$. (This is exactly the content of the principal stabilizer condition: at $Z' \in \mathcal{Z}^\circ$ the slice S_r is "transverse enough" that only block-diagonal elements can map $\iota(Z')$ back into S_r.)

Having established $h \in H_r$, we now compute (11) componentwise using (1):

$$\iota(Z)_\ell = \big(h \cdot \iota(Z')\big)_\ell = h_\ell \begin{pmatrix} I_r & 0 \\ 0 & Z'_\ell \end{pmatrix} h_{\ell-1}^{-1}.$$

Substituting $h_\ell = \left(\begin{smallmatrix} A & 0 \\ 0 & K_\ell \end{smallmatrix}\right)$ and $h_{\ell-1}^{-1} = \left(\begin{smallmatrix} A^{-1} & 0 \\ 0 & K_{\ell-1}^{-1} \end{smallmatrix}\right)$ and multiplying as in Step 2 yields

$$\begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} = \begin{pmatrix} I_r & 0 \\ 0 & K_\ell Z'_\ell K_{\ell-1}^{-1} \end{pmatrix},$$

so

$$Z_\ell = K_\ell\, Z'_\ell\, K_{\ell-1}^{-1} \quad \Longleftrightarrow \quad Z'_\ell = K_\ell^{-1}\, Z_\ell\, K_{\ell-1}. \tag{12}$$

But (12) is exactly the relation $Z' = h^{-1} \cdot Z$ from (6). Therefore $(g', Z') = (gh,\ h^{-1} \cdot Z)$, i.e. $(g', Z') = (g, Z) \cdot h$ in the right action (5). Hence $[g', Z'] = [g, Z]$ in $G \times_{H_r} \mathcal{Z}^\circ$, proving injectivity on the principal stabilizer locus.

Step 5 (Conclusion). Steps 3 and 4 show that $\bar{\Phi}$ restricted to $G \times_{H_r} \mathcal{Z}^\circ$ is bijective onto Ω_r°, hence (8). The global map $\bar{\Phi}\colon G \times_{H_r} \mathcal{Z} \to \Omega_r$ is surjective but may fail to be injective at points where the stabilizer of $\iota(Z')$ in G strictly contains H_r. □

Remark 8.4 (Hidden-layer gauge variant)**.** If only hidden-layer changes of basis are allowed, replace G by $G_h := \prod_{\ell=1}^{L-1} \mathrm{GL}(d_\ell, \mathbb{R})$ (i.e. fix g_0, g_L), and replace H_r by $H_{r,h} := H_r \cap G_h$. All statements and the proof above remain valid verbatim with this replacement, with $\mathcal{Z}^\circ$ redefined using Stab_{G_h} in place of Stab_G.

8.3 Critical points versus the cyclic variety Γ

Setup. Fix a depth $L \geq 2$, widths $(d_0, \ldots, d_L)$, and an integer $0 \leq r \leq \min_\ell d_\ell$. Let

$$\Omega \;:=\; \prod_{\ell=1}^{L} \mathrm{Mat}_{d_\ell \times d_{\ell-1}}(\mathbb{R})$$

and let the base–change group

$$G \;:=\; \prod_{\ell=0}^{L} GL(d_\ell, \mathbb{R})$$

act on Ω by

$$(g \cdot W)_\ell \;:=\; g_\ell \, W_\ell \, g_{\ell-1}^{-1}. \tag{5}$$

We also fix the data-dependent matrices appearing in the normal form of critical points (e.g. from the DLN second-order analysis):

$$U_S \in \mathbb{R}^{d_L \times r}, \qquad U_Q \in \mathbb{R}^{d_L \times (d_L - r)}, \qquad \sigma \;:=\; \Sigma_{YX} U_Q \in \mathrm{Hom}(\mathbb{R}^{d_L - r}, \mathbb{R}^{d_0 - r}).$$

(Here σ is the distinguished "closing arrow" in the cyclic-quiver model.)

Residual parameter space and constraints. Define the residual space

$$\mathcal{Z} \;:=\; \prod_{\ell=1}^{L} \mathrm{Mat}_{(d_\ell - r) \times (d_{\ell-1} - r)}(\mathbb{R}), \qquad Z = (Z_1, \ldots, Z_L).$$

Define the *fiber constraint* (multiplication fiber at 0)

$$\Sigma_0^{d-r} \;:=\; \{\, Z \in \mathcal{Z} : Z_L Z_{L-1} \cdots Z_1 = 0 \,\}. \tag{6}$$

Define the *cyclic constraints with fixed closing arrow* σ by

$$\Gamma_\sigma \;:=\; \Big\{ Z \in \mathcal{Z} \;\Big|\; Z_{\ell-1} \cdots Z_1 \, \sigma \, Z_L \cdots Z_{\ell+1} = 0 \;\text{ for all } \ell = 1, \ldots, L \Big\}. \tag{7}$$

Finally set

$$\mathcal{X}_{r,\sigma} \;:=\; \Sigma_0^{d-r} \cap \Gamma_\sigma \;\subset\; \mathcal{Z}. \tag{8}$$

A gauge-fixed slice of rank-r critical points. Define the map $\iota : \mathcal{Z} \to \Omega$ by putting residual blocks on the diagonal:

$$\iota(Z)_\ell \;:=\; \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} \in \mathrm{Mat}_{d_\ell \times d_{\ell-1}}(\mathbb{R}), \qquad \ell = 2, \ldots, L-1, \tag{9}$$

and set the endpoints to match the standard DLN critical-point normal form:

$$\iota(Z)_L \;:=\; [\, U_S,\, U_Q Z_L \,] \in \mathrm{Mat}_{d_L \times d_{L-1}}(\mathbb{R}), \qquad \iota(Z)_1 \;:=\; \begin{pmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ Z_1 \end{pmatrix} \in \mathrm{Mat}_{d_1 \times d_0}(\mathbb{R}). \tag{10}$$

(Only the block structure matters for what follows; any equivalent fixed endpoint embedding works.)

Define the *gauge-fixed critical slice*

$$\mathcal{S}_{r,\sigma} \;:=\; \iota(\mathcal{X}_{r,\sigma}) \;\subset\; \Omega. \tag{11}$$

Hidden-layer gauge group and its slice stabilizer. Let

$$G_h \;:=\; \prod_{\ell=1}^{L-1} GL(d_\ell, \mathbb{R})$$

act on Ω via (5) with $g_0 = g_L = I$ (endpoints fixed). Define the subgroup $H_{r,\sigma} \leq G_h$ consisting of those hidden reparametrizations that preserve the block decomposition $\mathbb{R}^{d_\ell} \cong \mathbb{R}^r \oplus \mathbb{R}^{d_\ell - r}$ for every hidden layer *and* preserve the endpoint embeddings (10):

$$H_{r,\sigma} \;:=\; \left\{ h = (h_1, \ldots, h_{L-1}) \in G_h \;\middle|\; h_\ell = \begin{pmatrix} A & 0 \\ 0 & K_\ell \end{pmatrix},\; A \in GL(r, \mathbb{R}),\; K_\ell \in GL(d_\ell - r, \mathbb{R}),\; \text{and } h \cdot \mathcal{S}_{r,\sigma} = \mathcal{S}_{r,\sigma} \right\}. \tag{12}$$

(Equivalently: $H_{r,\sigma}$ is the setwise stabilizer of $\mathcal{S}_{r,\sigma}$ inside G_h.)

The induced action of $H_{r,\sigma}$ on $\mathcal{Z}$ is the residual base-change action: for $h_\ell = \left(\begin{smallmatrix} A & 0 \\ 0 & K_\ell \end{smallmatrix}\right)$ we set

$$(h \cdot Z)_\ell \;:=\; K_\ell \, Z_\ell \, K_{\ell-1}^{-1} \qquad (\ell = 2, \ldots, L-1), \tag{13}$$

with the obvious endpoint conventions for $\ell = 1$ and $\ell = L$ determined by the fixed endpoint embeddings (10). By construction, $\mathcal{X}_{r,\sigma} \subset \mathcal{Z}$ is $H_{r,\sigma}$-stable (this is exactly the meaning of $h \cdot \mathcal{S}_{r,\sigma} = \mathcal{S}_{r,\sigma}$).

Theorem 8.5. Let $C_r \subset \Omega$ denote the set of rank-r first-order critical points in the normal form of the DLN second-order characterisation, i.e. the set described by the existence of residual blocks $Z = (Z_1, \ldots, Z_L)$ such that

$$W_L = [U_S,\, U_Q Z_L], \qquad W_\ell \sim \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} \; (\ell = 2, \ldots, L-1), \qquad W_1 = \begin{pmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ Z_1 \end{pmatrix}, \qquad Z \in \Sigma_0^{d-r} \cap \Gamma_\sigma.$$

Then the following hold.

(1) Orbit-saturation description.

$$C_r = G_h \cdot S_{r,\sigma}. \tag{22}$$

(2) Associated bundle model (set-level). Define $\Phi \colon G_h \times \mathcal{X}_{r,\sigma} \to \Omega$ by

$$\Phi(g, Z) := g \cdot \iota(Z), \tag{23}$$

where ι is defined by (17)–(18). Then $\Phi(G_h \times \mathcal{X}_{r,\sigma}) = C_r$ and Φ is invariant under the right action of $H_{r,\sigma}$ on $G_h \times \mathcal{X}_{r,\sigma}$ given by

$$(g, Z) \cdot h := (gh,\; h^{-1} \cdot Z). \tag{24}$$

Hence Φ descends to a surjective map

$$\bar{\Phi} \colon G_h \times_{H_{r,\sigma}} \mathcal{X}_{r,\sigma} \longrightarrow C_r.$$

Define the *transporter into the slice* at a point $Z' \in \mathcal{X}_{r,\sigma}$ by

$$\mathrm{Trans}_{G_h}\big(\iota(Z'),\, S_{r,\sigma}\big) := \big\{ g \in G_h : g \cdot \iota(Z') \in S_{r,\sigma} \big\}.$$

Define the *principal slice locus*

$$\mathcal{X}^{\circ}_{r,\sigma} := \big\{ Z' \in \mathcal{X}_{r,\sigma} \;\big|\; \mathrm{Trans}_{G_h}\big(\iota(Z'),\, S_{r,\sigma}\big) = H_{r,\sigma} \big\},$$

and let $C^{\circ}_r := \Phi(G_h \times \mathcal{X}^{\circ}_{r,\sigma})$. Then the restriction

$$\bar{\Phi}\colon G_h \times_{H_{r,\sigma}} \mathcal{X}^{\circ}_{r,\sigma} \xrightarrow{\sim} C^{\circ}_r$$

is a bijection. In particular, on the principal slice locus one has a canonical identification

$$C^{\circ}_r \;\cong\; G_h \times_{H_{r,\sigma}} \mathcal{X}^{\circ}_{r,\sigma} \qquad \text{(on the principal slice locus).} \tag{25}$$

(3) Quotient of critical points by hidden gauge. On the same principal slice locus,

$$C^{\circ}_r / G_h \;\cong\; \mathcal{X}^{\circ}_{r,\sigma} / H_{r,\sigma}. \tag{26}$$

Proof. We proceed in explicit steps.

Step 1: Proof of (22). By the assumed DLN normal form for rank-r critical points, $W \in C_r$ if and only if there exists $Z \in \mathcal{X}_{r,\sigma}$ and a sequence of hidden invertible changes of basis $g = (g_1, \ldots, g_{L-1}) \in G_h$ such that the transformed weights satisfy the block gauge-fixed identities (17)–(18). Equivalently, $g^{-1} \cdot W = \iota(Z) \in S_{r,\sigma}$. Therefore $W = g \cdot \iota(Z) \in G_h \cdot S_{r,\sigma}$, proving $C_r \subseteq G_h \cdot S_{r,\sigma}$. Conversely, if $W = g \cdot \iota(Z)$ with $Z \in \mathcal{X}_{r,\sigma}$, then W has the required normal form up to the hidden reparametrisation g, and since Z satisfies both (14) and (15), W is critical. Thus $G_h \cdot S_{r,\sigma} \subseteq C_r$. This proves (22).

Step 2: Image of Φ equals C_r. By definition (23), $\Phi(g, Z) = g \cdot \iota(Z) \in G_h \cdot S_{r,\sigma} = C_r$ by Step 1. Hence $\Phi(G_h \times \mathcal{X}_{r,\sigma}) \subseteq C_r$. Conversely, Step 1 already shows every $W \in C_r$ can be written $W = g \cdot \iota(Z)$ with $Z \in \mathcal{X}_{r,\sigma}$. Therefore $\Phi(G_h \times \mathcal{X}_{r,\sigma}) = C_r$. **Step 3: $H_{r,\sigma}$-invariance of Φ.** Fix $(g, Z) \in G_h \times \mathcal{X}_{r,\sigma}$ and $h \in H_{r,\sigma}$. Then

$$\Phi\big((g, Z) \cdot h\big) = \Phi\big(gh,\; h^{-1} \cdot Z\big) = (gh) \cdot \iota(h^{-1} \cdot Z).$$

Using associativity of the group action (13), this equals $g \cdot \big(h \cdot \iota(h^{-1} \cdot Z)\big)$. Thus it suffices to show

$$h \cdot \iota(h^{-1} \cdot Z) = \iota(Z). \tag{$\star$}$$

Write each $h_\ell = \left(\begin{smallmatrix} I_r & 0 \\ 0 & K_\ell \end{smallmatrix}\right)$ (recall that the setwise stabiliser condition and the endpoint structure force $A = I_r$; see the paragraph after Definition 8.2).

Middle layers ($\ell = 2, \ldots, L-1$)*:* Here $(h^{-1} \cdot Z)_\ell = K_\ell^{-1} Z_\ell K_{\ell-1}$, and the computation is identical to Step 2 of Proposition 8.3:

$$h_\ell \begin{pmatrix} I_r & 0 \\ 0 & K_\ell^{-1} Z_\ell K_{\ell-1} \end{pmatrix} h_{\ell-1}^{-1} = \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix} = \iota(Z)_\ell.$$

Last layer ($\ell = L$)*:* Since $g_L = I$ in G_h, the action gives $(h \cdot \iota(h^{-1} \cdot Z))_L = \iota(h^{-1} \cdot Z)_L\, h_{L-1}^{-1}$. Now $(h^{-1} \cdot Z)_L = Z_L K_{L-1}$ (since $g_L = I$ implies the effective residual action at layer L is right multiplication by K_{L-1}), so

$$[U_S,\; U_Q Z_L K_{L-1}] \begin{pmatrix} I_r & 0 \\ 0 & K_{L-1}^{-1} \end{pmatrix} = [U_S,\; U_Q Z_L] = \iota(Z)_L.$$

First layer ($\ell = 1$)*:* Since $g_0 = I$, the action gives $(h \cdot \iota(h^{-1} \cdot Z))_1 = h_1\, \iota(h^{-1} \cdot Z)_1$. Now $(h^{-1} \cdot Z)_1 = K_1^{-1} Z_1$ (since $g_0 = I$), so

$$\begin{pmatrix} I_r & 0 \\ 0 & K_1 \end{pmatrix} \begin{pmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ K_1^{-1} Z_1 \end{pmatrix} = \begin{pmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ Z_1 \end{pmatrix} = \iota(Z)_1.$$

This proves ($\star$), hence Φ is $H_{r,\sigma}$-invariant and descends to $\bar{\Phi}$.

Step 4: Injectivity on the principal slice locus. Assume $\Phi(g, Z) = \Phi(g', Z')$ with $Z, Z' \in X^\circ_{r,\sigma}$. Then $g \cdot \iota(Z) = g' \cdot \iota(Z')$, and setting $h := g^{-1} g' \in G_h$ we obtain

$$\iota(Z) = h \cdot \iota(Z'). \tag{$\star\star$}$$

In particular, h maps the slice point $\iota(Z')$ to another slice point $\iota(Z)$, so $h \in \mathrm{Trans}_{G_h}(\iota(Z'), S_{r,\sigma})$. Since $Z' \in \mathcal{X}^\circ_{r,\sigma}$, we have $\mathrm{Trans}_{G_h}(\iota(Z'), S_{r,\sigma}) = H_{r,\sigma}$ by definition of the principal slice locus. Therefore $h \in H_{r,\sigma}$.

Having established $h \in H_{r,\sigma}$, we compute ($\star\star$) componentwise. By the same block multiplication as in Step 3 (run in reverse), equation ($\star\star$) yields $Z = h \cdot Z'$, i.e. $Z' = h^{-1} \cdot Z$. Therefore $(g', Z') = (gh, h^{-1} \cdot Z) = (g, Z) \cdot h$ in the right action (24), and hence $[g', Z'] = [g, Z]$ in $G_h \times_{H_{r,\sigma}} \mathcal{X}^\circ_{r,\sigma}$. This proves injectivity on the principal slice locus.

Step 5: The principal slice locus is generically non-empty. We show that $\mathcal{X}^\circ_{r,\sigma}$ contains a dense open subset of $X_{r,\sigma}$, so that the bijection (25) holds on a large locus. Take any $h = (h_1, \ldots, h_{L-1}) \in G_h$ satisfying $h \cdot \iota(Z') \in S_{r,\sigma}$ for some $Z' \in X_{r,\sigma}$. We must show $h \in H_{r,\sigma}$ for generic Z'.

Write $h_\ell = \left(\begin{smallmatrix} a_\ell & b_\ell \\ c_\ell & k_\ell \end{smallmatrix}\right)$ in the $r \times (d_\ell - r)$ block decomposition.

Endpoint $\ell = L$*:* The condition $(h \cdot \iota(Z'))_L = \iota(Z)_L$ gives $[U_S, U_Q Z'_L]\, h_{L-1}^{-1} = [U_S, U_Q Z_L]$. Write $h_{L-1}^{-1} = \left(\begin{smallmatrix} \alpha & \beta \\ \gamma & \kappa \end{smallmatrix}\right)$. Comparing the first block column and using $U_S^\top U_Q = 0$ yields $\alpha = I_r$ (from the U_S-component) and $Z'_L \gamma = 0$ (from the U_Q-component). For generic $Z' \in X_{r,\sigma}$, the matrix Z'_L has full row rank $d_L - r$, so $\gamma = 0$. Hence h_{L-1}^{-1} is block upper-triangular; since it is also invertible, its inverse h_{L-1} is block upper-triangular as well.

Endpoint $\ell = 1$*:* The condition $(h \cdot \iota(Z'))_1 = \iota(Z)_1$ gives $h_1 \left(\begin{smallmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ Z'_1 \end{smallmatrix}\right) = \left(\begin{smallmatrix} U_S^\top \Sigma_{YX} \Sigma_X^{-1} \\ Z_1 \end{smallmatrix}\right)$. From the top block: $a_1 U_S^\top \Sigma_{YX} \Sigma_X^{-1} + b_1 Z'_1 = U_S^\top \Sigma_{YX} \Sigma_X^{-1}$. Generically Z'_1 is linearly independent from the rows of $U_S^\top \Sigma_{YX} \Sigma_X^{-1}$, which forces $b_1 = 0$ and $a_1 = I_r$. From the bottom block: $c_1 U_S^\top \Sigma_{YX} \Sigma_X^{-1} + k_1 Z'_1 = Z_1$; for this to admit a solution with Z_1 free, we need $c_1 = 0$ (since otherwise $c_1 U_S^\top \Sigma_{YX} \Sigma_X^{-1}$ would introduce a component in the signal subspace, contradicting the block structure). Hence h_1 is block-diagonal with $a_1 = I_r$.

Middle layers ($\ell = 2, \ldots, L-2$)*:* Having established that h_1 and h_{L-1} are block-diagonal, one propagates inward. At each layer the condition

$$h_\ell \begin{pmatrix} I_r & 0 \\ 0 & Z'_\ell \end{pmatrix} h_{\ell-1}^{-1} = \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix}$$

with $h_{\ell-1}$ already known to be block-diagonal forces $b_\ell Z'_\ell \kappa_{\ell-1} = 0$ and $c_\ell = 0$. Since $\kappa_{\ell-1}$ is invertible and generically Z'_ℓ has full row rank, we get $b_\ell = 0$. Hence h_ℓ is block-diagonal at every layer.

Thus, for Z' in a dense open subset of $X_{r,\sigma}$ (where certain residual blocks have full rank), every element of $\mathrm{Trans}_{G_h}(\iota(Z'), S_{r,\sigma})$ is block-diagonal, i.e. belongs to $H_{r,\sigma}$. Since $H_{r,\sigma} \subseteq \mathrm{Trans}_{G_h}(\iota(Z'), S_{r,\sigma})$

always holds (by Step 3), we conclude $\mathrm{Trans}_{G_h}(\iota(Z'), S_{r,\sigma}) = H_{r,\sigma}$ on this locus, confirming that $X^\circ_{r,\sigma}$ is generically non-empty.

Step 6: Conclusion. Steps 2 and 3 establish that $\bar{\Phi}$ is a well-defined surjection. Step 4 establishes injectivity on $\mathcal{X}^\circ_{r,\sigma}$, giving (25). Taking G_h-quotients gives (26). The global map $\bar{\Phi}\colon G_h \times_{H_{r,\sigma}} \mathcal{X}_{r,\sigma} \to C_r$ is surjective but may fail to be injective at points Z' where certain residual blocks Z'_ℓ are rank-deficient, causing $\mathrm{Trans}_{G_h}(\iota(Z'), S_{r,\sigma})$ to strictly contain $H_{r,\sigma}$. □

Remark 8.6 (How this corrects the conjectural formula)**.** The correct "product/quotient" description involves the hidden gauge group G_h and the intersection $X_{r,\sigma} = \Sigma_0^{d-r} \cap \Gamma_\sigma$ rather than Γ_d alone. A fully global isomorphism $C_r \cong G_h \times_{H_{r,\sigma}} X_{r,\sigma}$ can fail only where the transporter into the slice strictly contains $H_{r,\sigma}$; in that case the correct statement is that C_r is modelled by this associated bundle on each transporter stratum.

Lemma 8.7 (Codimension is determined by the principal slice locus)**.** With notation as in Theorem 8.5, assume the principal slice locus $\mathcal{X}^\circ_{r,\sigma} \subset \mathcal{X}_{r,\sigma}$ is Zariski open and dense in $\mathcal{X}_{r,\sigma}$, and let $C^\circ_r := \Phi(G_h \times \mathcal{X}^\circ_{r,\sigma}) \subset C_r$. Then C°_r is Zariski dense in C_r, and in particular

$$\mathrm{codim}_\Omega(C^\circ_r) \;=\; \mathrm{codim}_\Omega(C_r).$$

Proof. First note that $\mathcal{X}^\circ_{r,\sigma}$ Zariski open and dense in $\mathcal{X}_{r,\sigma}$ implies $G_h \times \mathcal{X}^\circ_{r,\sigma}$ is Zariski open and dense in $G_h \times \mathcal{X}_{r,\sigma}$, since G_h is irreducible and the product of a dense open subset with an irreducible variety is dense open in the product.

Consider the map $\Phi\colon G_h \times \mathcal{X}_{r,\sigma} \to \Omega$ defined in (23). It is a morphism (being the restriction of the algebraic action map $(g, W) \mapsto g \cdot W$ composed with the embedding ι), hence it is continuous in the Zariski topology. By Theorem 8.5(2), $\Phi(G_h \times \mathcal{X}_{r,\sigma}) = C_r$ and by definition $\Phi(G_h \times \mathcal{X}^\circ_{r,\sigma}) = C^\circ_r$.

Since Φ is continuous, for any subset $A \subseteq G_h \times \mathcal{X}_{r,\sigma}$ one has

$$\Phi(\overline{A}) \;\subseteq\; \overline{\Phi(A)},$$

where closures are taken in the Zariski topology. Apply this with $A = G_h \times \mathcal{X}^\circ_{r,\sigma}$. Because $G_h \times \mathcal{X}^\circ_{r,\sigma}$ is dense in $G_h \times \mathcal{X}_{r,\sigma}$, we have

$$\overline{G_h \times \mathcal{X}^\circ_{r,\sigma}} \;=\; G_h \times \mathcal{X}_{r,\sigma}.$$

Therefore,

$$C_r \;=\; \Phi(G_h \times \mathcal{X}_{r,\sigma}) \;=\; \Phi\big(\overline{G_h \times \mathcal{X}^\circ_{r,\sigma}}\big) \;\subseteq\; \overline{\Phi(G_h \times \mathcal{X}^\circ_{r,\sigma})} \;=\; \overline{C^\circ_r}.$$

Since always $C^\circ_r \subseteq C_r$, we also have $\overline{C^\circ_r} \subseteq C_r$, hence

$$\overline{C^\circ_r} \;=\; C_r,$$

i.e. C°_r is Zariski dense in C_r.

Finally, Zariski density implies $\dim C^\circ_r = \dim C_r$ (because C_r is the Zariski closure of C°_r), and hence

$$\mathrm{codim}_\Omega(C^\circ_r) = \dim \Omega - \dim C^\circ_r = \dim \Omega - \dim C_r = \mathrm{codim}_\Omega(C_r).$$

□

Proposition 8.8 (Codimension comparison: C_r versus $\mathcal{X}_{r,\sigma}$)**.** With notation as in Theorem 8.5, assume the hypotheses of Lemma 8.7. Then the codimensions satisfy

$$\operatorname{codim}_\Omega(C_r) \;=\; \operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma}) \;+\; \Big(\dim\Omega - \dim\mathcal{Z} - \dim G_h + \dim H_{r,\sigma}\Big). \tag{14}$$

Equivalently,

$$\operatorname{codim}_\Omega(C_r) \;=\; \operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma}) \;+\; r(d_0 + d_L),$$

where the last equality uses the explicit dimension identities

$$\dim\Omega = \sum_{\ell=1}^{L} d_\ell d_{\ell-1}, \qquad \dim\mathcal{Z} = \sum_{\ell=1}^{L}(d_\ell - r)(d_{\ell-1} - r), \qquad \dim G_h = \sum_{\ell=1}^{L-1} d_\ell^2, \qquad \dim H_{r,\sigma} = r^2 + \sum_{\ell=1}^{L-1}(d_\ell - r)^2.$$

9 Sampling saddle points

We leverage the algebraic parameterization of first-order critical points in deep linear networks (DLNs) given by the rank condition and the cyclic conditions. The objective is to turn these polynomial constraints into a numerical sampler that produces many (typically gauge-inequivalent) saddle points.

9.1 The set of critical points

We can now relate the set of critical points $\mathcal{C}_r$ to the variety Γ_d of elements of $\mathcal{Z}$ satisfying the cyclic conditions. Conjecture: We have:

$$\mathcal{C}_r \cong G_d \otimes \Gamma_d / H_{r,\sigma}$$

With

$$H_{r,\sigma} = \{h \in H_r | h \cdot Z \in \Gamma_d \text{ for all } Z \in \Gamma_{d,\sigma}\}$$

Rank strata and target projection. Let the teacher (or population) matrix be

$$M = U \operatorname{diag}(s_1, \ldots, s_{d_{\text{out}}})\, V^\top,$$

and fix a rank r and a subset S of singular directions with $|S| = r$. Define U_S as the columns of U indexed by S, and the corresponding projector $P_S := U_S U_S^\top$. A rank-r critical stratum is characterized by the end-to-end product

$$W := W_L \cdots W_1 = P_S M, \qquad \operatorname{rank}(W) = r.$$

In practice, we choose $r < r_{\max}$ to avoid global minimizers and bias toward saddle points.

Gauge-fixed normal form and residual variables. We work in a fixed representative of the hidden-layer gauge orbit. Each hidden dimension d_ℓ is split into a signal part of dimension r and a residual part of dimension $d_\ell - r$. In this gauge, the layers are parameterized as

$$W_\ell \;\sim\; \begin{pmatrix} I_r & 0 \\ 0 & Z_\ell \end{pmatrix}, \qquad \ell = 2, \ldots, L-1,$$

with residual blocks $Z_\ell \in \mathbb{R}^{(d_\ell - r)\times(d_{\ell-1}-r)}$. The first and last layers embed the signal path and attach the residuals:

$$W_L = [\,U_S \;\; U_Q Z_L\,], \qquad W_1 = \begin{pmatrix} U_S^\top M \\ Z_1 \end{pmatrix},$$

where U_Q spans the orthogonal complement of U_S. The signal blocks are chosen so that the product along the signal path equals $P_S M$ exactly, ensuring the rank condition by construction.

Cyclic and fiber constraints on the residuals. In this normal form, first-order criticality imposes polynomial constraints on the residual blocks $Z = (Z_1, \ldots, Z_L)$:

- **Fiber (product) constraint:**

$$\mu(Z) := Z_L Z_{L-1} \cdots Z_1 = 0.$$

- **Cyclic constraints:**

$$d\mu_\ell(Z) := Z_{\ell-1} \cdots Z_1 \, \sigma \, Z_L \cdots Z_{\ell+1} = 0, \qquad \ell = 1, \ldots, L,$$

  where σ is the fixed matrix appearing in the theoretical characterization (e.g. $\sigma = \Sigma_{XY} U_Q$ in the population setting, or an empirical analogue).

Energy formulation. To convert these matrix equalities into a scalar objective, we introduce the constraint energy

$$E(Z) \;=\; \frac{1}{2}\|\mu(Z)\|_F^2 \;+\; \frac{1}{2}\sum_{\ell=1}^{L} \|d\mu_\ell(Z)\|_F^2.$$

Then $E(Z) = 0$ if and only if all cyclic and fiber constraints are satisfied exactly.

Ensuring saddle points. A cheap sufficient construction for producing saddle points is to enforce two distinct residual blocks to be zero. Concretely, choose $a \neq b$ and set

$$Z_a = 0, \qquad Z_b = 0,$$

while sampling the remaining Z_ℓ. This automatically enforces the fiber constraint and typically satisfies all cyclic constraints. More expressive (but more expensive) constructions avoid exact zeros and instead certify saddle behavior numerically via negative curvature of the Hessian.

Boltzmann sampling over constrained residuals. To sample many solutions rather than a single feasible point, we sample from a Boltzmann distribution concentrating near the constraint variety:

$$p_\beta(Z) \;\propto\; \exp\big(-\beta\, E(Z)\big)\, \pi(Z),$$

where $\beta > 0$ controls concentration and $\pi(Z)$ is a prior that prevents collapse to trivial solutions. Typical choices include Gaussian or Gaussian with weight decay

$$\log \pi(Z) = -\frac{\alpha}{2}\sum_{\ell=1}^{L} \|Z_\ell\|_F^2,$$

or norm-targeting penalties that keep $\|Z_\ell\|_F$ away from zero.

SGLD updates. Define the negative log-density (up to an additive constant)

$$\mathcal{L}_\beta(Z) := \beta\, E(Z) - \log \pi(Z).$$

Stochastic gradient Langevin dynamics (SGLD) then takes the form

$$Z_{t+1} = Z_t - \frac{\epsilon_t}{2} \nabla \mathcal{L}_\beta(Z_t) + \eta_t, \qquad \eta_t \sim \mathcal{N}(0, \epsilon_t I),$$

with stepsize schedule ϵ_t. When empirical analogues of the constraints are used, gradients can be estimated on minibatches.

Projection to true DLN critical points and Hessian certification. Comment: I'm not sure about the usefulness of doing this: Because the parameterization and the matrix σ may be approximate in finite-sample settings, after obtaining a low-energy configuration Z we assemble the full weights $W_1, \ldots, W_L$ and perform a short refinement step on the true DLN loss to reach $\|\nabla \mathcal{L}\| \approx 0$, while periodically re-balancing layer norms to control gauge drift. **Comment: this is going to be useful:** finally, saddle points are certified by estimating the smallest eigenvalues of the Hessian using Hessian-vector products; the presence of a negative eigenvalue indicates a strict saddle and a non strict saddle otherwise (or perhap§s local minimum in the underparametrize regime).

10 Cyclic quiver representations

A cyclic quiver is a quiver with at list one oriented cycle. In particular we will be interested in a quiver which is exactly one cycle here noted C_n.

Definition 10.1 (Cyclic quiver)**.** A *cyclic quiver* (or *oriented cycle*) on $n \geq 1$ vertices is the quiver C_n with vertex set

$$P = \{0, 1, \ldots, n-1\}$$

and arrow set

$$E = \{a_i : i \to i+1 \mid i = 0, \ldots, n-2\} \cup \{a_{n-1} : n-1 \to 0\},$$

i.e. the vertices are arranged in a directed cycle.

Definition 10.2 (Representation of a cyclic quiver)**.** A *representation* of the cyclic quiver C_n (arrows of C_n are of the opposite orientation) over a field k is the data of vector spaces

$$(V_0, \ldots, V_{n-1})$$

together with linear maps

$$\phi_i : V_{i-1} \to V_i \quad (i = 1, \ldots, n-1), \qquad \phi_n : V_{n-1} \to V_0,$$

i.e. a representation of the opposite quiver C_n.

Proposition 10.3 (The relations defining Γ as a cyclic quiver representation)**.** Fix $L \geq 1$ and consider the cyclic quiver C_{L+1} on vertices

$$\{0, 1, \ldots, L\}.$$

Let r be the rank of the student DLN and:

$$V_l \cong \mathbb{R}^{d_l - r}, \qquad l = 0, \ldots, L,$$

and consider a representation of the opposite quiver C_{L+1} given by linear maps

$$\phi_l : V_{l-1} \to V_l \quad (l = 1, \ldots, L), \qquad \phi_{L+1} : V_L \to V_0,$$

where indices are understood cyclically.

For $\phi = (\phi_1, \ldots, \phi_{L+1}) \in \mathrm{Rep}_d(C_{L+1})$, define for $l = 1, \ldots, L$ the maps[3]:

$$d\mu_l(\phi) := \phi_{l-1} \cdots \phi_1 \, \phi_{L+1} \, \phi_L \cdots \phi_{l+1} \ \in \mathrm{Hom}(V_l, V_{l-1})$$

with the convention $\phi_0 = \mathrm{id}_{V_0}$. The maps $d\mu_l$ correspond to the components of the differential of the product map around the cycle. They correspond to a path of length L.

Define the closed subvariety

$$\Gamma_d := \Big\{ \phi \in \mathrm{Rep}_d(C_{L+1}) \ \Big| \ d\mu_l(\phi) = 0 \text{ for all } l = 1, \ldots, L \Big\}.$$

Identifying $\phi_l = Z_l$ for $l = 1, \ldots, L$, we now *specialize the closing arrow*

$$\phi_{L+1} := \Sigma_{XY} U_Q : V_L \to V_0.$$

For this specialization, the algebraic variety Γ related to the critical loci of a DLN is:

$$\Gamma := \Gamma_d \big|_{\phi_{L+1} = \Sigma_{XY} U_Q}$$

coincides with the space of cyclic representations satisfying the vanishing $d\mu$ relations and with the distinguished arrow ϕ_{L+1} fixed.

Proposition 10.4 (Equivariance of the cyclic composition maps)**.** Let C_{L+1} be the cyclic quiver on vertices $\{0, \ldots, L\}$, let $\mathrm{Rep}_d(C_{L+1})$ be its representation space with $V_l \cong \mathbb{R}^{d_l - r}$, and let

$$\phi = (\phi_1, \ldots, \phi_L, \phi_{L+1}) \in \mathrm{Rep}_d(C_{L+1}).$$

Define the base–change group

$$G_d := \prod_{l=0}^{L} \mathrm{GL}(V_l)$$

acting on $\mathrm{Rep}_d(C_{L+1})$ via

$$(g \cdot \phi)_l := g_l \, \phi_l \, g_{l-1}^{-1}, \quad (l = 0, \ldots, L), \ \mathrm{mod}\ L+1$$

For each $l = 1, \ldots, L$, define

$$d\mu_l(\phi) := \phi_{l-1} \cdots \phi_1 \, \phi_{L+1} \, \phi_L \cdots \phi_{l+1} \in \mathrm{Hom}(V_{l+1}, V_l),$$

with the convention that an empty product is the identity. Then, for all $g \in G_d$ and all $l = 0, \ldots, L$,

$$d\mu_l(g \cdot \phi) = g_l \, d\mu_l(\phi) \, g_{l+1}^{-1}.$$

[3] And such that the linear compositions are compatible with the cyclic quiver



$d\mu_1 : V_2 \to V_1$ (composition along the cycle, omitting ϕ_2)

Figure 1: The cyclic quiver C_n (illustrated here with $n = 8$). The thick arrows indicate the path whose composition gives a typical $d\mu_l$ map along the cycle (example: $d\mu_2 : V_2 \to V_1$).

Proof. Expand $d\mu_l(g \cdot \phi)$ as a product of the maps $(g \cdot \phi)_i$. Each factor contributes a left multiplication by some g_i and a right multiplication by g_{i+1}^{-1}, and all intermediate terms cancel by associativity, leaving only g_l on the left and g_{l+1}^{-1} on the right. □

Proposition 10.5 (Orbit decomposition of Γ_d)**.** Let

$$\Gamma_d := \Big\{ \phi \in \mathrm{Rep}_d(C_{L+1}) \;\Big|\; d\mu_l(\phi) = 0 \text{ for all } l = 1, \dots, L \Big\}.$$

Then Γ_d is G_d-invariant. In particular, Γ_d decomposes as a disjoint union of G_d-orbits:

$$\Gamma_d = \bigsqcup_{\phi \in \Gamma_d} G_d \cdot \phi.$$

Proof. By Proposition 10.4, each map $d\mu_l$ is G_d-equivariant and hence the vanishing set $d\mu_l^{-1}(0)$ is G_d-stable. Intersecting over $l = 1, \dots, L$ shows that Γ_d is G_d-stable. Any group action partitions a set into disjoint orbits, yielding the stated decomposition. □

Note that $\Gamma_d \cong \mathrm{Rep}_d(C_{L+1}) / < d\mu_l >$ is then finite dimensional and we can clasified indecomposable using indecomposable representations of Nakayama algebras.

10.1 Cyclic quiver as a bounded module

Definition 10.6 (Path algebra)**.** Let Q be a quiver. The *path algebra* kQ is the k-vector space with basis given by all directed paths in Q (including the trivial paths e_i of length 0 at each vertex i), with multiplication given by concatenation of composable paths and 0 otherwise, extended by bilinearity.

Definition 10.7 (Bound quiver algebra associated with Γ_d)**.** Let $Q = C_{L+1}$ and for each $l = 0, \dots, L$ let $p_l \in kQ$ be the path

$$p_l := a_{l-1} \cdots a_1 \, a_{L+1} \cdots a_{l+1},$$

interpreting empty products as the corresponding trivial path. Let $I \subset kQ$ be the two-sided ideal generated by $\{p_1, \ldots, p_L\}$. The quotient algebra

$$A := kQ/I$$

is called the *bound quiver algebra* associated with these relations.

Definition 10.8 (Representation of an algebra of dimension d)**.** Let A be a k-algebra and let $d = (d_0, \ldots, d_L)$. A *representation of A of dimension d compatible with the vertices* is a left A-module structure on $V := \bigoplus_{i=0}^{L} k^{d_i}$ such that the idempotents $e_i \in kQ \twoheadrightarrow A$ act as the projections onto the i-th summand. We denote the corresponding representation space by $\mathrm{Rep}_d(A)$.

Proposition 10.9 (From quiver representations with relations to A-modules)**.** There is a well-defined map

$$F : \Gamma_d \to \mathrm{Rep}_d(A)$$

which sends $\phi \in \Gamma_d$ to the unique A-module structure on V for which each arrow a_i acts by the linear map ϕ_i.

Proof. Given $\phi \in \Gamma_d$, the tuple of linear maps $(\phi_i)_i$ defines a representation of the quiver $Q = C_{L+1}^{op}$, hence a left kQ-module structure on V by letting a path act by the corresponding composition of arrow maps. For each l, the defining condition $\mu_l(\phi) = 0$ is exactly the statement that the element $p_l \in kQ$ acts as the zero map on V. Since I is the two-sided ideal generated by the p_l, every element of I also acts by zero. Therefore the kQ-action factors through the quotient $A = kQ/I$, producing a unique A-module structure on V with the required arrow actions. □

Proposition 10.10 (From A-modules to quiver representations with relations)**.** There is a well-defined map

$$G : \mathrm{Rep}_d(A) \to \Gamma_d$$

which sends an A-module structure on V to the induced quiver representation ϕ whose arrow maps are the actions of the classes of the arrows a_i.

Proof. Let V be a representation in $\mathrm{Rep}_d(A)$. Composing the action map $A \to \mathrm{End}_k(V)$ with the quotient map $kQ \twoheadrightarrow A$ gives a kQ-module structure on V. Restricting the action of each arrow basis element a_i to the appropriate vertex summand yields linear maps $\phi_i : V_i \to V_{i+1}$ and hence a point $\phi \in \mathrm{Rep}_d(C_n)$. Because each p_l maps to zero in A, it acts as the zero endomorphism on V. Equivalently, the composed linear map $\mu_l(\phi)$ is zero for every l, so $\phi \in \Gamma_d$. □

Proposition 10.11 (Correspondence between Γ_d and $\mathrm{Rep}_d(A)$)**.** The maps F and G are mutually inverse bijections. In particular,

$$\Gamma_d \cong \mathrm{Rep}_d(A), \qquad A = kQ/I.$$

Proof. Starting from $\phi \in \Gamma_d$, the A-module $F(\phi)$ is by construction generated by the same arrow actions as ϕ, so applying G recovers the same tuple of arrow maps, hence $G(F(\phi)) = \phi$. Conversely, starting from an A-module structure on V, the map G records the arrow actions and the map F reconstructs the unique A-module whose arrow actions agree with those recorded, hence $F(G(V)) = V$. □

Lemma 10.12 (Finite-dimensionality of $A = kQ/I$)**.** Let $Q = C_{L+1}$ be the oriented cycle on $L+1$ vertices, and let $I \subset kQ$ be the two-sided ideal generated by the paths $p_1, \ldots, p_L$, where each p_l is a path of length L (the unique directed path going around the cycle while omitting one arrow). Then every directed path in Q of length $\geq L+1$ lies in I. In particular, $A = kQ/I$ is finite-dimensional.

Proof. Let q be any directed path in Q of length $\geq L+1$. Write $q = b_m b_{m-1} \cdots b_1$ as a concatenation of arrows with $m \geq L+1$. Consider the contiguous subpath $q' := b_L \cdots b_1$ of length L. Since Q is an oriented $L+1$-cycle, there is exactly one directed path of length L from its source to its target, hence q' must coincide with one of the generators p_l. Therefore $q = q''\, p_l$ for some (possibly trivial) path q'', so $q \in I$ because I is a two-sided ideal.

Thus all paths of length $\geq L+1$ vanish in A, so A is spanned by the residue classes of paths of length $< L+1$, a finite set. Hence A is finite-dimensional. □

10.2 Bounded modules are Nakayama algebras

Definition 10.13 (A-module)**.** Let A be a (not necessarily commutative) k-algebra. A (left) *A-module* is a k-vector space M together with a k-bilinear map

$$A \times M \to M, \qquad (a, m) \mapsto a \cdot m,$$

such that for all $a, b \in A$ and $m \in M$ one has

$$(ab) \cdot m = a \cdot (b \cdot m), \qquad 1_A \cdot m = m.$$

A *homomorphism* of A-modules is a k-linear map commuting with the A-action.

Definition 10.14 (Indecomposable module)**.** An A-module M is *indecomposable* if $M \neq 0$ and there do not exist nonzero submodules $M_1, M_2 \subset M$ such that

$$M \cong M_1 \oplus M_2.$$

Equivalently, M is indecomposable if every idempotent endomorphism of M is either 0 or id_M.

Definition 10.15 (Composition series)**.** Let M be a finite-dimensional A-module. A *composition series* of M is a finite chain of submodules

$$0 = M_0 \subset M_1 \subset \cdots \subset M_\ell = M$$

such that each successive quotient M_i/M_{i-1} is *simple*. The integer ℓ is called the *length* of the series.

Definition 10.16 (Projective module)**.** An A-module P is *projective* if for every surjective A-module homomorphism $f : M \twoheadrightarrow N$ and every A-module homomorphism $g : P \to N$, there exists an A-module homomorphism $\tilde{g} : P \to M$ such that $f \circ \tilde{g} = g$. Equivalently, P is projective if the functor $\mathrm{Hom}_A(P, -)$ is exact.

Definition 10.17 (Indecomposable projective module)**.** A *projective indecomposable* A-module is an A-module that is both projective and indecomposable. For a finite-dimensional basic algebra A, every indecomposable projective module is of the form $P_i \cong Ae_i$ for a primitive idempotent $e_i \in A$, and every projective module decomposes uniquely as a direct sum of indecomposable projectives.

Definition 10.18 (Uniserial module)**.** An A-module M is *uniserial* if it admits a unique composition series. Equivalently, the set of submodules of M is totally ordered by inclusion.

Definition 10.19 (**Nakayama algebra**)**.** Let A be a finite-dimensional k-algebra. We say that A is a *Nakayama algebra* if every indecomposable left A-module is *uniserial*, i.e. it admits a unique composition series. Equivalently, A is Nakayama if every indecomposable projective left A-module is uniserial (and hence every indecomposable injective is uniserial).

Definition 10.20 (Admissible ideal)**.** Let Q be a finite quiver and let kQ be its path algebra. Write $(kQ)_{\geq m}$ for the k-span of all paths of length at least m. A two-sided ideal $I \subset kQ$ is called *admissible* if there exists an integer $m \geq 2$ such that

$$(kQ)_{\geq m} \subseteq I \subseteq (kQ)_{\geq 2}.$$

Definition 10.21 (Jacobson radical)**.** Let A be a (not necessarily commutative) ring. The *Jacobson radical* $J(A)$ is the intersection of all maximal left ideals of A. If A is a finite-dimensional k-algebra, then equivalently

$$J(A) = \bigcap_{S \text{ simple}} \ker\big(A \to \mathrm{End}_k(S)\big),$$

i.e. $J(A)$ is the largest two-sided ideal that acts by zero on every simple A-module.

Proposition 10.22 (**Bound quiver criterion for Nakayama algebras**)**.** Let $A \simeq kQ/I$ be a finite-dimensional bound quiver algebra, where Q is a finite quiver and $I \subset kQ$ is an admissible ideal. Assume that for every vertex $i \in Q_0$ there is at most one arrow leaving i and at most one arrow entering i. Then A is a Nakayama algebra.

Moreover, if Q is connected and has exactly one arrow leaving and exactly one arrow entering each vertex, then Q is an oriented cycle and A is a *cyclic Nakayama algebra*. If instead Q is connected and has a unique source and a unique sink, then Q is an oriented line and A is a *linear Nakayama algebra*.

Proof. Let $e_i \in kQ \twoheadrightarrow A$ be the primitive idempotent corresponding to a vertex $i \in Q_0$, and consider the indecomposable projective module $P_i := Ae_i$. As a k-vector space, P_i is spanned by residue classes of paths in Q starting at i (modulo the relations in I). Because there is at most one arrow leaving each vertex, for each $\ell \geq 0$ there is *at most one* directed path of length ℓ starting at i; hence P_i has a basis indexed by (the surviving classes of) these paths.

Let $J \subset A$ denote the Jacobson radical. Since $A \simeq kQ/I$ with I admissible, J is generated (as a two-sided ideal) by the classes of arrows, and one has

$$J^\ell P_i = J^\ell A e_i \;\cong\; (J^\ell) e_i,$$

which is spanned by classes of paths of length $\geq \ell$ starting at i. By the previous paragraph, for each ℓ the quotient $J^\ell P_i / J^{\ell+1} P_i$ is either 0 or a simple module (it is spanned by the class of the unique surviving path of length ℓ, if it exists). Therefore the radical filtration

$$P_i \supset JP_i \supset J^2 P_i \supset \cdots$$

is a chain whose successive quotients are simple (or zero), i.e. P_i is uniserial.

Thus every indecomposable projective P_i is uniserial, and hence A is a Nakayama algebra.

For the final statements: if Q is connected and every vertex has exactly one incoming and one outgoing arrow, then the underlying directed graph is a directed cycle. If Q is connected and has a unique source and sink with all other vertices having one incoming and one outgoing arrow, then the directed graph is a directed line. These correspond to cyclic and linear Nakayama algebras respectively. □

10.3 Cyclic monomial Nakayama algebra with uniform Kupisch series

Throughout this subsection, let $Q = C_{L+1}$ be the oriented $L+1$-cycle, let kQ be its path algebra, and let

$$A := kQ/I$$

be the bound quiver algebra from Definition (Bound quiver algebra associated with Γ_d).

Definition 10.23 (Monomial (path) ideal / monomial algebra)**.** An ideal $I \subset kQ$ is called *monomial* if it is generated (as a two-sided ideal) by paths in Q. A quotient kQ/I with I monomial is called a *monomial algebra*.

Definition 10.24 (Kupisch series)**.** Let A be a finite-dimensional basic algebra with primitive idempotents $e_1, \dots, e_{L+1}$ and indecomposable projectives $P_i := Ae_i$. The *Kupisch series* of A is the $L+1$-tuple

$$(c_1, \dots, c_{L+1}), \qquad c_i := \ell(P_i),$$

where $\ell(P_i)$ denotes the composition length of P_i.

Remark 10.25 (Loewy length)**.** For a finite-dimensional algebra A with Jacobson radical $J = J(A)$, the *Loewy length* is the smallest integer n such that $J^n = 0$.

Proposition 10.26 (Identification and Kupisch series)**.** With $Q = C_{L+1}$ and $I = (p_1, \dots, p_{L+1})$ as above, the algebra $A = kQ/I$ is a *cyclic monomial Nakayama algebra*. Moreover its Kupisch series is uniform:

$$(c_1, \dots, c_{L+1}) = (L, \dots, L).$$

Proof. By construction, the ideal I is generated by the paths $p_1, \dots, p_{L+1}$, hence I is monomial and A is a monomial algebra. By Proposition 10.22, A is Nakayama; since $Q = C_{L+1}$ is an oriented cycle, A is in fact *cyclic* Nakayama.

Fix a vertex i and consider $P_i = Ae_i$. Modulo I, every path in Q of length $\geq L$ starting at i vanishes, while for each $t = 0, 1, \dots, L-1$ there is exactly one directed path of length t starting at i. Therefore P_i has a k-basis given by the residue classes of these L paths (length 0 through $L-1$).

Since $J = J(A)$ is generated by the classes of arrows (as recalled in the proof of Proposition 10.22), the radical filtration

$$P_i \supset JP_i \supset J^2P_i \supset \cdots$$

has successive quotients $J^tP_i/J^{t+1}P_i$ one-dimensional (hence simple) for $t = 0, \dots, L-1$, and zero thereafter. Thus $\ell(P_i) = L$ for every i, which gives the Kupisch series $(L, \dots, L)$. □

Remark 10.27. Equivalently, A can be viewed as a *truncated cycle algebra*: it is the path algebra of an oriented cycle with all paths of length $\geq L$ killed.

10.4 Indecomposable representations of Nakayama algebras

Definition 10.28 (The uniserial modules $M(i,\ell)$)**.** Fix $Q = C_{L+1}$ with vertices $Q_0 = \{0,\dots,L\}$ and arrows $a_i : l-1 \to l$ for $l = 1,\dots,L$ and $a_{L+1} : L \to 0$. For $i \in Q_0$ and $0 \le \ell \le L$, define a representation $M(i,\ell)$ of Q by

$$M(i,\ell)_j := \begin{cases} k, & j \in \{i, i+1, \dots, i+\ell-1\} \pmod{L+1}, \\ 0, & \text{otherwise}, \end{cases}$$

and for each arrow $a_j : j-1 \to j$,

$$M(i,\ell)_{a_j} := \begin{cases} \mathrm{id}_k, & \text{if both source and target are } k, \\ 0, & \text{otherwise}. \end{cases}$$

Lemma 10.29 (The relations vanish on $M(i,\ell)$)**.** Let $Q = C_{L+1}$ and, for each $l \in \{1,\dots,L+1\}$, let

$$p_l := a_{l-1} \cdots a_1\, a_{L+1} \cdots a_{l+1}$$

be the oriented path of length L going once around the cycle and omitting a_l. Set $A := kQ/\langle p_1,\dots,p_{L+1}\rangle$. Then, for every $i \in Q_0$ and $1 \le \ell \le L$, the representation $M(i,\ell)$ factors through A, hence defines a finite-dimensional left A-module.

Proof. Fix i and ℓ. Each path p_l has length L. Since the support of $M(i,\ell)$ contains only $\ell \le L$ consecutive vertices, the corresponding composition along p_l necessarily passes through a vertex where $M(i,\ell)$ is zero. Therefore p_l acts by the zero map on $M(i,\ell)$ for every l, so all generators of the ideal $\langle p_1,\dots,p_L\rangle$ act trivially. Hence the kQ-action factors through A. □

Lemma 10.30 (Uniseriality of $M(i,\ell)$)**.** For every $i \in Q_0$ and $1 \le \ell \le L+1$, the A-module $M(i,\ell)$ is uniserial, and in particular indecomposable.

Proof. Any nonzero submodule $N \subseteq M(i,\ell)$ must contain the one-dimensional space at some vertex in the support. Because all arrow maps inside the support are identities, containing a vertex forces N to contain all vertices reachable along the directed arrows within the support. Consequently, the nonzero submodules are exactly the modules $M(i+t,\ell-t)$ for $t = 0,\dots,\ell-1$ (indices taken modulo n), and they form a chain under inclusion. Thus submodules are totally ordered and $M(i,\ell)$ is uniserial. □

Lemma 10.31 (Endomorphisms of $M(i,\ell)$)**.** For every $i \in Q_0$ and $1 \le \ell \le L$, one has $\mathrm{End}_A\big(M(i,\ell)\big) \cong k$.

Proof. Let $f \in \mathrm{End}_A(M(i,\ell))$. On each vertex j in the support, f is multiplication by some scalar $\lambda_j \in k$. For each arrow inside the support, the arrow map is id_k, so commutativity forces $\lambda_{j+1} = \lambda_j$. Hence all λ_j coincide with a single scalar λ, and $f = \lambda\,\mathrm{id}$. □

Lemma 10.32 (Projective truncations realize $M(i,\ell)$)**.** Let $P(i) := Ae_i$ be the indecomposable projective at vertex i and let $J := J(A)$. Then, for every $i \in Q_0$ and $1 \le \ell \le L$, there is an isomorphism of A-modules

$$P(i)/J^\ell P(i) \;\cong\; M(i,\ell).$$

Proof. By Proposition 10.22 and the previous subsection, A is a cyclic Nakayama algebra with uniform Kupisch series $(L, \dots, L)$. In particular, $P(i)$ is uniserial of length L, and $J^t P(i)$ is spanned by residue classes of paths of length $\geq t$ starting at i. Thus $P(i)/J^\ell P(i)$ has a basis given by the residue classes of the unique paths of lengths $0, 1, \dots, \ell - 1$ starting at i. Under the induced quiver description, this quotient has one-dimensional spaces on the vertices $i, i+1, \dots, i+\ell-1 \pmod{L+1}$ and identity maps along arrows inside the support, which is exactly $M(i, \ell)$. □

Proposition 10.33 (Indecomposable representations of the cyclic truncated Nakayama algebra)**.** Let $Q = C_{L+1}$ be the oriented cyclic quiver with arrows $a_i : i \to i+1$ for $i = 1, \dots, L$ and $a_{L+1} : L \to 0$. For each $l \in \{1, \dots, L+1\}$, let p_l be the path of length L going once around the cycle and omitting a_l, and set

$$A := kQ/\langle p_1, \dots, p_{L+1} \rangle.$$

Then every finite-dimensional indecomposable left A-module is isomorphic to a unique module $M(i, \ell)$ with $i \in Q_0$ and $1 \leq \ell \leq L$. Equivalently, every indecomposable A-module is of the form $P(i)/J^\ell P(i)$ for a unique pair (i, ℓ). In particular, A has exactly $(L+1)L$ indecomposable modules up to isomorphism.

Proof. By Lemma 10.29, each $M(i, \ell)$ is a well-defined finite-dimensional A-module, and by Lemma 10.30 it is indecomposable.

Conversely, since A is Nakayama (Proposition 10.22), every indecomposable A-module is uniserial and hence a quotient of a unique indecomposable projective $P(i) = Ae_i$. Its composition length is some ℓ with $1 \leq \ell \leq \ell(P(i)) = L$, and uniseriality forces the quotient to be $P(i)/J^\ell P(i)$. By Lemma 10.32, this module is isomorphic to $M(i, \ell)$.

Uniqueness follows because the top simple module determines i and the length determines ℓ. Counting pairs (i, ℓ) with $i \in Q_0$ and $1 \leq \ell \leq L$ gives $(L+1)L$. □

More details about this can be found in the following lecture notes: Noncommutative Algebra 2: Representations of finite-dimensional algebras in particular theorem and corroloray section 1.6. We have a Krull schmidt theorem for these Nakayama algebras.

Proposition 10.34 (Krull–Schmidt for the cyclic truncated Nakayama algebra)**.** Let $A = kQ/I$ be the cyclic truncated Nakayama algebra, where $Q = C_{L+1}$ is the oriented $L+1$-cycle and I is the ideal generated by the length-(L) paths $p_1, \dots, p_{L+1}$ (equivalently, all paths of length $\geq L+1$ vanish in A). Let mod-A denote the category of finite-dimensional left A-modules. Then every $M \in$ mod-A admits a decomposition as a finite direct sum of indecomposable A-modules. Moreover, this decomposition is unique up to reordering and isomorphism of the indecomposable summands. Equivalently, using the explicit indecomposables $M(i, \ell)$ from Proposition 10.33, there exist uniquely determined multiplicities $m_{i,\ell} \in \mathbb{Z}_{\geq 0}$ such that

$$M \cong \bigoplus_{i=0}^{L} \bigoplus_{\ell=1}^{L} M(i, \ell)^{\oplus m_{i,\ell}}.$$

Proof sketch. Since A is finite-dimensional over k, it is left artinian (and left noetherian), hence every object of mod-A has finite length. In particular, mod-A is a Krull–Schmidt category: every object decomposes as a finite direct sum of indecomposables, and such a decomposition is unique up to permutation and isomorphism.

Concretely, existence follows by choosing a nonzero module M and repeatedly splitting off a nontrivial direct summand until all summands are indecomposable; the process terminates because M has finite length. Uniqueness follows because each indecomposable has a local endomorphism ring (for example, by Lemma 10.31 in the explicit classification), and the classical Krull–Schmidt–Azumaya argument applies.

Finally, Proposition 10.33 classifies the indecomposables as the modules $M(i,\ell)$, so any Krull–Schmidt decomposition can be rewritten uniquely in the stated multiplicity form. □

Note: This enables to parameterize the orbits of Γ with the indecomposables representations.

Orbits, isomorphism classes, and multiplicity patterns. Fix a finite-dimensional k-algebra A and a dimension vector $d = (d_0, \dots, d_L)$. Recall that

$$\mathrm{Rep}_d(A) = \Big\{\text{left } A\text{-module structures on } V = \bigoplus_{i=0}^{L} k^{d_i} \text{ such that each idempotent } e_i \text{ acts as the projection onto } k^{d_i}\Big\}.$$

Let

$$G_d := \prod_{i=0}^{L} \mathrm{GL}(k^{d_i})$$

act on $\mathrm{Rep}_d(A)$ by *change of bases at the vertices* (equivalently, by transport of structure): for $g = (g_0, \dots, g_L) \in G_d$ and a module structure $\rho : A \to \mathrm{End}_k(V)$, define a new module structure $g \cdot \rho$ by

$$(g \cdot \rho)(a) \;:=\; g\,\rho(a)\,g^{-1}, \qquad g := \bigoplus_{i=1}^{n} g_i \in \mathrm{GL}(V).$$

When $A \simeq kQ/I$ is presented as a bound quiver algebra, this action is equivalently given on arrow maps by $(g \cdot \phi)_a = g_{t(a)}\,\phi_a\,g_{s(a)}^{-1}$. Then two points $\rho, \rho' \in \mathrm{Rep}_d(A)$ lie in the same G_d-orbit if and only if the corresponding A-modules (V, ρ) and (V, ρ') are isomorphic in mod-A. Consequently, there is a canonical bijection

$$\big(G_d \backslash \mathrm{Rep}_d(A)\big) \;\cong\; \Big\{\text{isomorphism classes of } A\text{-modules } M \text{ with } \dim(e_i M) = d_i \text{ for all } i\Big\}.$$

Now specialize to the cyclic truncated Nakayama algebra $A = kQ/I$ considered above. Its indecomposable modules are the uniserials $M(i,\ell)$ with $i \in \{0, \dots, L\}$ (indices taken modulo $L+1$) and $1 \le \ell \le L$ (Proposition 10.33). By the Krull–Schmidt theorem (Proposition 10.34), every finite-dimensional A-module M admits a decomposition

$$M \;\simeq\; \bigoplus_{i=0}^{L} \bigoplus_{\ell=1}^{L} M(i,\ell)^{\oplus m_{i,\ell}}, \qquad m_{i,\ell} \in \mathbb{Z}_{\ge 0},$$

and this decomposition is unique up to permutation of summands. The multiplicities $(m_{i,\ell})$ determine (and are constrained by) the dimension vector d via the *covering equations*

$$d_k \;=\; \sum_{i=0}^{L} \sum_{\ell=1}^{L} m_{i,\ell}\, \mathbf{1}\{k \in \{i, i+1, \dots, i+\ell-1\} \pmod{L+1}\}, \qquad k = 0, \dots, L, \tag{15}$$

since $M(i,\ell)$ contributes a one-dimensional space at vertex k exactly when k lies in its cyclic support interval of length ℓ starting at i. Define the set of *multiplicity patterns*

$$M_d^+(A) := \left\{ (m_{i,\ell}) \in \mathbb{Z}_{\geq 0}^{L(L+1)} \;\middle|\; (15) \text{ holds for all } k \right\}.$$

Then the above discussion yields a concrete parametrization of orbits: every G_d-orbit in $\mathrm{Rep}_d(A)$ corresponds to a unique isomorphism class in mod-A, and hence to a unique multiplicity pattern $(m_{i,\ell}) \in M_d^+(A)$ via its Krull–Schmidt decomposition. In other words, we obtain canonical bijections

$$G_d \backslash \mathrm{Rep}_d(A) \;\cong\; \left\{ \text{isomorphism classes in mod-}A \text{ with dimension vector } d \right\} \;\cong\; M_d^+(A),$$

where the last map sends an A-module to the multiplicities of the indecomposables $M(i,\ell)$ in its Krull–Schmidt decomposition.

Proposition 10.35 (Orbit stratification of Γ_d by indecomposable Nakayama modules)**.** Fix $L \geq 1$ and let $Q = C_{L+1}$ be the oriented cyclic quiver on vertices $\{0, \ldots, L\}$. Let $A := kQ/I$ be the bound quiver algebra, where I is the ideal generated by the length-L paths $p_1, \ldots, p_{L+1}$ obtained by going once around the cycle and omitting one arrow. Let $d = (d_0, \ldots, d_L)$ be a dimension vector.

Define

$$\Gamma_d := \{ \phi \in \mathrm{Rep}_d(Q) \mid d\mu_\ell(\phi) = 0 \text{ for all } \ell = 1, \ldots, L \},$$

where

$$d\mu_\ell(\phi) := \phi_{\ell-1} \cdots \phi_1 \, \phi_{L+1} \, \phi_L \cdots \phi_{\ell+1}.$$

Let

$$G_d := \prod_{i=0}^{L} \mathrm{GL}(k^{d_i})$$

act on Γ_d by base change at the vertices.

Then the following hold.

1. There is a canonical identification

$$\Gamma_d \;\cong\; \mathrm{Rep}_d(A),$$

under which the G_d-action coincides with transport of structure on A-modules.

2. The algebra A is a finite-dimensional cyclic Nakayama algebra. Its indecomposable finite-dimensional left modules are, up to isomorphism, the uniserial modules

$$M(i,\ell), \qquad i \in \{0, \ldots, L\}, \quad 1 \leq \ell \leq L,$$

supported on ℓ consecutive vertices starting at i (indices taken modulo $L+1$).

3. Every A-module M of dimension vector d admits a unique (up to permutation) Krull–Schmidt decomposition

$$M \;\cong\; \bigoplus_{i=0}^{L} \bigoplus_{\ell=1}^{L} M(i,\ell)^{\oplus m_{i,\ell}},$$

for uniquely determined multiplicities $m_{i,\ell} \in \mathbb{Z}_{\geq 0}$ satisfying the dimension constraints

$$d_k \;=\; \sum_{i=0}^{L} \sum_{\ell=1}^{L} m_{i,\ell}\, \mathbf{1}\{k \in \{i, i+1, \ldots, i+\ell-1\} \pmod{L+1}\}\,, \quad k = 0, \ldots, L.$$

Denote by $\mathcal{M}_d^+(A)$ the finite set of all such multiplicity patterns $m = (m_{i,\ell})$.

4. For each $m \in \mathcal{M}_d^+(A)$, define the stratum

$$\Gamma_d^{(m)} := \left\{ \phi \in \Gamma_d \;\middle|\; F(\phi) \cong \bigoplus_{i,\ell} M(i,\ell)^{\oplus m_{i,\ell}} \right\},$$

where $F(\phi)$ denotes the associated A-module.

Then $\Gamma_d^{(m)}$ is a single G_d-orbit.

5. The variety Γ_d decomposes as a disjoint union of these orbits:

$$\Gamma_d \;=\; \bigsqcup_{m \in \mathcal{M}_d^+(A)} \Gamma_d^{(m)} \;=\; \bigsqcup_{m \in \mathcal{M}_d^+(A)} G_d \cdot \phi_m,$$

where ϕ_m is any representative whose associated A-module has multiplicity pattern m.

In particular, the G_d-orbits in Γ_d are canonically parameterized by the multiplicity data of indecomposable A-modules, i.e. by the set $\mathcal{M}_d^+(A)$.

11 Stratification of Γ_{XY}

While the indecomposable representation of Γ are parameterized by the $M(i,\ell)$ from the previous modules over Nakayama algebras we are actually interested in decomposing:

$$\Gamma_\sigma := \{(\phi_1, \ldots, \phi_{L+1}) \in \mathrm{Rep}_d(C_{L+1}) \mid \phi_{l-1} \cdots \phi_1\, \phi_{L+1}\, \phi_L \cdots \phi_{l+1} = 0,\ \phi_{L+1} = \sigma\}\,.$$

where the map $\sigma \in \mathrm{Hom}(V_L, V_0)$ is fixed (in DLN it is $\Sigma_{XY} U_Q$). The map

$$d\mu_{l,\sigma} := \phi_{l-1} \cdots \phi_1\, \sigma\, \phi_L \cdots \phi_{l+1}$$

is no longer G_d-equivariant.

Definition 11.1 (Stabilizer group of the fixed arrow)**.** Let $\sigma \in \mathrm{Hom}(V_L, V_0)$ be fixed. Define

$$\mathrm{Stab}(\sigma) := \{(g_0, g_L) \in \mathrm{GL}(V_0) \times \mathrm{GL}(V_L) \mid g_0\, \sigma\, g_L^{-1} = \sigma\},$$

and the corresponding subgroup of the base–change group

$$G_\sigma := \{g = (g_0, \ldots, g_L) \in \prod_{l=0}^{L} \mathrm{GL}(V_l) \mid (g_0, g_L) \in \mathrm{Stab}(\sigma)\}.$$

Proposition 11.2 (G_σ-equivariance of the cyclic composition maps)**.** Fix $\sigma \in \mathrm{Hom}(V_L, V_0)$ and consider the representation space with $\phi_{L+1} = \sigma$. For $\phi = (\phi_1, \dots, \phi_L, \sigma)$ and each $l = 1, \dots, L$, define

$$d\mu_{l,\sigma}(\phi) := \phi_{l-1} \cdots \phi_1 \, \sigma \, \phi_L \cdots \phi_{l+1} \in \mathrm{Hom}(V_{l+1}, V_l),$$

with the convention that an empty product is the identity. Then, for all $g \in G_\sigma$ and all l,

$$d\mu_{l,\sigma}(g \cdot \phi) = g_l \, \mu_{l,\sigma}(\phi) \, g_{l+1}^{-1}.$$

In particular, the subset $\Gamma_\sigma := \bigcap_{l=1}^{L} d\mu_{l,\sigma}^{-1}(0)$ is G_σ-invariant.

Proof. Expand $d\mu_{l,\sigma}(g \cdot \phi)$ as a product of the maps $(g \cdot \phi)_i$. All intermediate base-change factors cancel. The only remaining factors are g_l on the left and g_{l+1}^{-1} on the right. The middle arrow is $(g \cdot \phi)_{L+1} = g_1 \sigma g_L^{-1}$, which equals σ because $g \in G_\sigma$. □

For all modules in Γ_σ Krull-Schmidt applies (as a category of A-modules where A is finite dimensional). We want to parameterize the orbits of Γ_σ under the action of G_σ and study their geometry (e.g. codimension).

Lemma 11.3 (Intersection of a G_d–orbit with a fixed–arrow fiber)**.** Let $L \geq 2$ and let $V_0, \dots, V_L$ be finite–dimensional vector spaces over an algebraically closed field $\Bbbk$. Set

$$G_d \;:=\; \prod_{i=0}^{L} GL(V_i).$$

Let Γ_d be an affine variety equipped with an algebraic action of G_d. Assume there exists a morphism

$$F : \Gamma_d \longrightarrow \mathrm{Hom}(V_L, V_0)$$

satisfying the equivariance property

$$F(g \cdot x) \;=\; g_0 \, F(x) \, g_L^{-1} \qquad \text{for all } g = (g_0, \dots, g_L) \in G_d, \; x \in \Gamma_d. \tag{16}$$

Fix $\sigma \in \mathrm{Hom}(V_L, V_0)$ and define the fiber

$$\Gamma_\sigma \;:=\; F^{-1}(\sigma).$$

Define the stabilizer subgroup

$$G_\sigma \;:=\; \{g = (g_0, \dots, g_L) \in G_d \mid g_0 \, \sigma \, g_L^{-1} = \sigma\}.$$

Let $\mathcal{O} \subset \Gamma_d$ be a single G_d–orbit. Then either $\mathcal{O} \cap \Gamma_\sigma = \varnothing$, or else for every $x \in \mathcal{O} \cap \Gamma_\sigma$ one has

$$\mathcal{O} \cap \Gamma_\sigma \;=\; G_\sigma \cdot x.$$

In particular, whenever $\mathcal{O} \cap \Gamma_\sigma \neq \varnothing$, this intersection consists of a single G_σ–orbit.

Proof. We proceed in several steps.

Step 1: Γ_σ is stable under the action of G_σ.

Let $x \in \Gamma_\sigma$, so by definition $F(x) = \sigma$. Let $g \in G_\sigma$. Using the equivariance property (16),

$$F(g \cdot x) \;=\; g_1\, F(x)\, g_L^{-1} \;=\; g_1\, \sigma\, g_L^{-1}.$$

Since $g \in G_\sigma$, we have $g_1\, \sigma\, g_L^{-1} = \sigma$, and hence $F(g \cdot x) = \sigma$. Therefore $g \cdot x \in \Gamma_\sigma$. This shows that Γ_σ is invariant under the action of G_σ and, in particular,

$$G_\sigma \cdot x \subseteq \Gamma_\sigma \qquad \text{for all } x \in \Gamma_\sigma. \tag{$*$}$$

Step 2: If $x \in \mathcal{O} \cap \Gamma_\sigma$, then $G_\sigma \cdot x \subseteq \mathcal{O} \cap \Gamma_\sigma$.

Let $x \in \mathcal{O} \cap \Gamma_\sigma$. Since $\mathcal{O}$ is a G_d–orbit, it is stable under the action of G_d and hence under the subgroup $G_\sigma \subseteq G_d$. Thus $G_\sigma \cdot x \subseteq \mathcal{O}$. By $(*)$, $G_\sigma \cdot x \subseteq \Gamma_\sigma$. Combining these two inclusions gives

$$G_\sigma \cdot x \subseteq \mathcal{O} \cap \Gamma_\sigma. \tag{$**$}$$

Step 3: Any two points of $\mathcal{O} \cap \Gamma_\sigma$ differ by the action of G_σ.

Assume $x, y \in \mathcal{O} \cap \Gamma_\sigma$. Because x and y lie in the same G_d–orbit $\mathcal{O}$, there exists $g = (g_1, \ldots, g_L) \in G_d$ such that

$$y = g \cdot x.$$

Applying F and using (16), we obtain

$$F(y) \;=\; F(g \cdot x) \;=\; g_1\, F(x)\, g_L^{-1}.$$

Since $x, y \in \Gamma_\sigma$, we have $F(x) = F(y) = \sigma$. Therefore

$$\sigma = g_1\, \sigma\, g_L^{-1}.$$

By definition of G_σ, this implies $g \in G_\sigma$. Hence $y = g \cdot x \in G_\sigma \cdot x$, and we conclude that

$$\mathcal{O} \cap \Gamma_\sigma \subseteq G_\sigma \cdot x. \tag{$***$}$$

Step 4: Conclusion.

If $\mathcal{O} \cap \Gamma_\sigma = \varnothing$, there is nothing to prove. Otherwise choose any $x \in \mathcal{O} \cap \Gamma_\sigma$. Inclusion $(**)$ gives $G_\sigma \cdot x \subseteq \mathcal{O} \cap \Gamma_\sigma$, while inclusion $(***)$ gives the reverse inclusion. Therefore

$$\mathcal{O} \cap \Gamma_\sigma = G_\sigma \cdot x.$$

This proves that whenever the intersection is not empty, it consists of a single G_σ–orbit. The lemma follows. □

Proposition 11.4 (Rank label for Nakayama orbits and the fiber non-emptiness criterion over $\mathbb{R}$)**.** Fix $L \geq 2$ and finite-dimensional real vector spaces $V_1, \ldots, V_L$. Let

$$G_d \;:=\; \prod_{i=0}^{L} GL(V_i) \qquad \text{and} \qquad W \;:=\; \mathrm{Hom}_{\mathbb{R}}(V_L, V_0).$$

Let Γ_d be a real algebraic variety equipped with an algebraic action of G_d. Assume there is a morphism of real varieties

$$F : \Gamma_d \longrightarrow W$$

satisfying the equivariance identity

$$F\big((g_0,\dots,g_L)\cdot x\big) = g_0\,F(x)\,g_L^{-1} \qquad \text{for all } (g_0,\dots,g_L)\in G_d,\ x\in\Gamma_d. \tag{17}$$

Definition (Orbit label and rank label). Assume Γ_d admits a decomposition into G_d–orbits indexed by a parameter set Λ,

$$\Gamma_d \;=\; \bigsqcup_{\lambda\in\Lambda} \mathcal{O}_\lambda,$$

where $\mathcal{O}_\lambda$ denotes the G_d–orbit corresponding to λ (in the applications of this paper, λ encodes the Krull–Schmidt multiplicities of the Nakayama indecomposables). For each $\lambda\in\Lambda$, define the integer

$$r(\lambda) \;:=\; \operatorname{rank}\big(F(x)\big) \quad \text{for any } x\in\mathcal{O}_\lambda. \tag{18}$$

Then the following statements hold.

(i) The integer $r(\lambda)$ in (18) is well-defined, i.e. it does not depend on the choice of $x\in\mathcal{O}_\lambda$.

(ii) Fix $\sigma\in W$ and define the fiber $\Gamma_\sigma := F^{-1}(\sigma)$. Then for every $\lambda\in\Lambda$,

$$\mathcal{O}_\lambda\cap\Gamma_\sigma\neq\varnothing \qquad\Longleftrightarrow\qquad r(\lambda)=\operatorname{rank}(\sigma). \tag{19}$$

Proof. We prove (i) and (ii) in separate steps.

Step 1: A basic invariance identity for ranks under left–right change of bases. Let $A\in GL(V_0)$, $B\in GL(V_L)$, and $T\in W=\operatorname{Hom}_{\mathbb{R}}(V_L,V_0)$. Consider the map $T' := A\,T\,B^{-1}\in W$. We have $\operatorname{rank}(T')=\operatorname{rank}(T)$.

Step 2: Proof of (i) (well-definedness of $r(\lambda)$). Fix $\lambda\in\Lambda$ and choose two points $x,y\in\mathcal{O}_\lambda$. By definition of orbit, there exists $g=(g_0,\dots,g_L)\in G_d$ such that

$$y = g\cdot x.$$

Apply F to both sides and use the equivariance identity (17):

$$F(y)=F(g\cdot x)=g_0\,F(x)\,g_L^{-1}.$$

Now apply Step 1 with $A=g_0$, $B=g_L$, and $T=F(x)$. This yields

$$\operatorname{rank}(F(y))=\operatorname{rank}(g_0\,F(x)\,g_L^{-1})=\operatorname{rank}(F(x)).$$

Thus the rank of $F(\,\cdot\,)$ is constant along $\mathcal{O}_\lambda$, so the integer $r(\lambda)$ defined by (18) is independent of the choice of $x\in\mathcal{O}_\lambda$. This proves (i).

Step 3: A real classification fact for linear maps by rank. Let $S,T\in W=\operatorname{Hom}_{\mathbb{R}}(V_L,V_1)$. We use the following statement:

Claim. If $\operatorname{rank}(S)=\operatorname{rank}(T)$, then there exist $A\in GL(V_1)$ and $B\in GL(V_L)$ such that

$$S = A\,T\,B^{-1}. \tag{20}$$

Step 4: Proof of (ii), forward implication. Assume $\mathcal{O}_\lambda \cap \Gamma_\sigma \neq \varnothing$. Choose $x \in \mathcal{O}_\lambda \cap \Gamma_\sigma$. By definition of $\Gamma_\sigma = F^{-1}(\sigma)$ we have $F(x) = \sigma$, hence

$$\operatorname{rank}(\sigma) = \operatorname{rank}(F(x)).$$

By definition of $r(\lambda)$ (and Step 2), $\operatorname{rank}(F(x)) = r(\lambda)$ for any $x \in \mathcal{O}_\lambda$. Therefore $\operatorname{rank}(\sigma) = r(\lambda)$, proving the forward direction of (19).

Step 5: Proof of (ii), reverse implication. Assume $r(\lambda) = \operatorname{rank}(\sigma)$. Choose any $x_0 \in \mathcal{O}_\lambda$. Then, by definition of $r(\lambda)$,

$$\operatorname{rank}(F(x_0)) = r(\lambda) = \operatorname{rank}(\sigma).$$

By the claim in Step 3 (applied to $T = F(x_0)$ and $S = \sigma$), there exist $A \in GL(V_0)$ and $B \in GL(V_L)$ such that

$$\sigma = A\, F(x_0)\, B^{-1}.$$

Define $g = (g_0, \ldots, g_L) \in G_d$ by setting

$$g_0 := A, \qquad g_L := B, \qquad g_i := \mathrm{Id}_{V_i} \ \text{ for } i = 1, \ldots, L-1.$$

Then $g \in G_d$ and, using equivariance (17),

$$F(g \cdot x_0) = g_0\, F(x_0)\, g_L^{-1} = A\, F(x_0)\, B^{-1} = \sigma.$$

Hence $g \cdot x_0 \in \Gamma_\sigma$. Also, since $x_0 \in \mathcal{O}_\lambda$ and $\mathcal{O}_\lambda$ is a G_d–orbit, we have $g \cdot x_0 \in \mathcal{O}_\lambda$. Therefore $g \cdot x_0 \in \mathcal{O}_\lambda \cap \Gamma_\sigma$, so $\mathcal{O}_\lambda \cap \Gamma_\sigma \neq \varnothing$.

Combining Steps 4 and 5 yields (19). This completes the proof. □

Proposition 11.5 (Rank stratification of Γ_{XY} by Nakayama types)**.** Let $d = (d_1, \ldots, d_L)$ be a dimension vector and let

$$\Gamma_d \subset \mathrm{Rep}_d(C_{L+1})$$

be the variety defined in Section §9. Recall that Γ_d admits a decomposition into G_d–orbits indexed by

$$\mathcal{M}_d^+ \;=\; \left\{ (m_{i,\ell}) \;\middle|\; \sum_{i,\ell} m_{i,\ell}\, \underline{\dim} M(i,\ell) = d \right\},$$

where $M(i,\ell)$ are the indecomposable modules of the cyclic Nakayama algebra and $(m_{i,\ell})$ records their multiplicities. For $\mathbf{m} = (m_{i,\ell}) \in \mathcal{M}_d^+$, denote by $\mathcal{O}_{\mathbf{m}} \subset \Gamma_d$ the corresponding G_d–orbit.

Let

$$F : \Gamma_d \longrightarrow \mathrm{Hom}(V_L, V_1), \qquad F(\phi) = \phi_{L+1},$$

and fix a linear map $\sigma \in \mathrm{Hom}(V_L, V_1)$. Define

$$\Gamma_{XY} := \Gamma_\sigma := F^{-1}(\sigma), \qquad G_\sigma := \{ g \in G_d : \ g_1 \sigma g_L^{-1} = \sigma \}.$$

For $\mathbf{m} \in \mathcal{M}_d^+$, define the integer

$$r(\mathbf{m}) \;:=\; \operatorname{rank}\big(F(\phi)\big) \quad \text{for any } \phi \in \mathcal{O}_{\mathbf{m}}.$$

Then:

(i) The integer $r(\mathbf{m})$ is well-defined (independent of the choice of $\phi \in \mathcal{O}_{\mathbf{m}}$).

(ii) For every $\mathbf{m} \in \mathcal{M}_d^+$,

$$\mathcal{O}_{\mathbf{m}} \cap \Gamma_{XY} \neq \varnothing \quad \Longleftrightarrow \quad r(\mathbf{m}) = \operatorname{rank}(\sigma).$$

(iii) If $\mathcal{O}_{\mathbf{m}} \cap \Gamma_{XY} \neq \varnothing$, then

$$\mathcal{O}_{\mathbf{m}} \cap \Gamma_{XY}$$

is a single G_σ–orbit.

(iv) Consequently, Γ_{XY} admits the disjoint decomposition

$$\Gamma_{XY} \;=\; \bigsqcup_{\mathbf{m}\in\mathcal{M}_d^+\,:\,r(\mathbf{m})=\operatorname{rank}(\sigma)} \big(\mathcal{O}_{\mathbf{m}} \cap \Gamma_{XY}\big),$$

and each stratum in this decomposition is a single G_σ–orbit.

Proposition 11.6 (Rank formula for the closing arrow in terms of Nakayama multiplicities)**.** Work over $\mathbb{R}$. Fix $L \geq 2$ and a dimension vector $d = (d_0, \ldots, d_L)$. Let $\Gamma_d \subset \operatorname{Rep}_d(C_{L+1})$ be the variety from the document, and recall that Γ_d decomposes into G_d–orbits indexed by $\mathcal{M}_d^+$, i.e. for each

$$\mathbf{m} = (m_{i,\ell}) \in \mathcal{M}_d^+$$

there is a corresponding G_d–orbit $\mathcal{O}_{\mathbf{m}} \subset \Gamma_d$ such that every $\phi \in \mathcal{O}_{\mathbf{m}}$ has associated Nakayama module

$$M(\phi) \;\cong\; \bigoplus_{i,\ell} M(i,\ell)^{\oplus m_{i,\ell}}$$

(up to isomorphism), where $M(i,\ell)$ denote the indecomposable modules of the cyclic Nakayama algebra.

Define the morphism

$$F : \Gamma_d \to \operatorname{Hom}_{\mathbb{R}}(V_L, V_1), \qquad F(\phi) = \phi_{L+1}.$$

For $\mathbf{m} \in \mathcal{M}_d^+$ define

$$r(\mathbf{m}) \;:=\; \operatorname{rank}(F(\phi)) \qquad \text{for any } \phi \in \mathcal{O}_{\mathbf{m}},$$

which is well-defined by Proposition 11.4.

Definition (support interval and activity indicator). For an indecomposable $M(i,\ell)$, define its *support interval* to be the cyclic set of vertices

$$\operatorname{Supp}(i,\ell) \;:=\; \{\, i,\ i+1,\ \ldots,\ i+\ell-1 \,\} \subset \mathbb{Z}/(L+1)\mathbb{Z},$$

where addition is taken modulo L and we identify vertices with their residue classes. Say that $M(i,\ell)$ is $(L \to 0)$–*active* if the arrow $L \to 0$ acts nontrivially on $M(i,\ell)$, i.e. if the structure map

$$M(i,\ell)(L) \longrightarrow M(i,\ell)(0)$$

induced by the quiver arrow $L \to 0$ is nonzero. Define the indicator

$$\epsilon(i,\ell) \;:=\; \begin{cases} 1, & \text{if } M(i,\ell) \text{ is } (L \to 0)\text{–active}, \\ 0, & \text{otherwise.} \end{cases}$$

Then for every $\mathbf{m} = (m_{i,\ell}) \in \mathcal{M}_d^+$ one has the formula

$$r(\mathbf{m}) \;=\; \sum_{i=0}^{L} \sum_{\ell=1}^{L} \epsilon(i,\ell)\, m_{i,\ell}. \tag{21}$$

Proof. Fix $\mathbf{m} = (m_{i,\ell}) \in \mathcal{M}_d^+$. Choose any $\phi \in \mathcal{O}_{\mathbf{m}}$ and write $M := M(\phi)$ for the corresponding Nakayama module. By definition of $\mathcal{O}_{\mathbf{m}}$, there exists an isomorphism of modules

$$M \;\cong\; \bigoplus_{i,\ell} M(i,\ell)^{\oplus m_{i,\ell}}. \tag{22}$$

We will compute $\operatorname{rank}(\phi_{L+1}) = \operatorname{rank}(F(\phi))$ from this decomposition.

Step 1: Describe the vector spaces at vertices 0 and L using (22).

For each vertex $j \in \{0, L\}$ the representation space is $V_j = M(j)$. Evaluating (22) at the vertex j gives an isomorphism of real vector spaces

$$V_j \;=\; M(j) \;\cong\; \bigoplus_{i,\ell} M(i,\ell)(j)^{\oplus m_{i,\ell}}. \tag{23}$$

For cyclic Nakayama indecomposables, $M(i,\ell)(j)$ is either 0 or a 1–dimensional real vector space (depending on whether $j \in \operatorname{Supp}(i,\ell)$). Therefore each summand in (23) is either 0 or $\mathbb{R}^{m_{i,\ell}}$.

Step 2: The closing arrow respects the direct-sum decomposition.

The linear map $\phi_{L+1} : V_L \to V_0$ is the structure map on the quiver arrow $L \to 0$. Because M is a direct sum of submodules as in (22), the action of every quiver arrow, and in particular the arrow $L \to 0$, decomposes as the direct sum of its actions on each direct summand. Concretely, after choosing identifications as in (23) for $j = L$ and $j = 0$, the map ϕ_{L+1} can be written as a block-diagonal direct sum

$$\phi_{L+1} \;\cong\; \bigoplus_{i,\ell} \big(\phi_{L+1}\big)_{i,\ell}, \tag{24}$$

where

$$\big(\phi_{L+1}\big)_{i,\ell} :\; M(i,\ell)(L)^{\oplus m_{i,\ell}} \longrightarrow M(i,\ell)(1)^{\oplus m_{i,\ell}}$$

is the map induced by $L \to 1$ on the summand $M(i,\ell)^{\oplus m_{i,\ell}}$.

Step 3: Compute each block $\big(\phi_{L+1}\big)_{i,\ell}$.

Fix a pair (i,ℓ). There are two cases.

Case A: $\epsilon(i,\ell) = 0$. By definition, $\epsilon(i,\ell) = 0$ means the arrow $L \to 1$ acts trivially on $M(i,\ell)$, i.e. the map

$$M(i,\ell)(L) \to M(i,\ell)(1)$$

is the zero map. Taking $m_{i,\ell}$ copies, the induced map

$$\big(\phi_{L+1}\big)_{i,\ell} :\; M(i,\ell)(L)^{\oplus m_{i,\ell}} \to M(i,\ell)(1)^{\oplus m_{i,\ell}}$$

is also the zero map. Hence

$$\operatorname{rank}\big(\big(\phi_{L+1}\big)_{i,\ell}\big) = 0. \tag{25}$$

Case B: $\epsilon(i,\ell) = 1$. By definition, $\epsilon(i,\ell) = 1$ means the arrow $L \to 1$ acts nontrivially on $M(i,\ell)$. For Nakayama indecomposables, both spaces $M(i,\ell)(L)$ and $M(i,\ell)(1)$ are then 1–dimensional and the map between them is a nonzero linear map $\mathbb{R} \to \mathbb{R}$. Choose bases of these 1–dimensional spaces; in these bases the map is multiplication by a nonzero scalar $a_{i,\ell} \in \mathbb{R}^\times$. On $m_{i,\ell}$ copies, the induced map is multiplication by the same scalar on each copy; in coordinates it is

$$a_{i,\ell} \cdot I_{m_{i,\ell}} : \ \mathbb{R}^{m_{i,\ell}} \to \mathbb{R}^{m_{i,\ell}},$$

which is invertible because $a_{i,\ell} \neq 0$. Therefore

$$\operatorname{rank}\big(\big(\phi_{L+1}\big)_{i,\ell}\big) = m_{i,\ell}. \tag{26}$$

Step 4: Sum the ranks of the blocks.

By (24), after choosing bases compatible with the decomposition, ϕ_{L+1} is block diagonal with blocks $\big(\phi_{L+1}\big)_{i,\ell}$. The rank of a block-diagonal linear map is the sum of the ranks of its diagonal blocks (because its image is the direct sum of the images of the blocks). Hence

$$\operatorname{rank}(\phi_{L+1}) \ = \ \sum_{i,\ell} \operatorname{rank}\big(\big(\phi_{L+1}\big)_{i,\ell}\big).$$

Substituting (25) in Case A and (26) in Case B yields

$$\operatorname{rank}(\phi_{L+1}) \ = \ \sum_{i,\ell} \epsilon(i,\ell)\, m_{i,\ell}.$$

By definition $r(\mathbf{m}) = \operatorname{rank}(\phi_{L+1})$ for $\phi \in \mathcal{O}_{\mathbf{m}}$, so this proves (21). □

Corollary 11.7 (Orbit decomposition of Γ_{XY} indexed by $\mathcal{M}_d^+$)**.** Work over $\mathbb{R}$ and keep the notation of Proposition 11.5 and Proposition 11.6. In particular, $\Gamma_{XY} = \Gamma_\sigma := F^{-1}(\sigma)$, the G_d–orbits in Γ_d are denoted $\mathcal{O}_{\mathbf{m}}$ for $\mathbf{m} \in \mathcal{M}_d^+$, and $G_\sigma := \{g \in G_d : \ g_1 \sigma g_L^{-1} = \sigma\}$.

Then Γ_{XY} decomposes as a disjoint union of G_σ–orbits indexed by those $\mathbf{m} \in \mathcal{M}_d^+$ satisfying the rank constraint:

$$\Gamma_{XY} \ = \ \bigsqcup_{\mathbf{m} \in \mathcal{M}_d^+ \,:\, r(\mathbf{m}) = \operatorname{rank}(\sigma)} \big(\mathcal{O}_{\mathbf{m}} \cap \Gamma_{XY}\big),$$

and for each such $\mathbf{m}$ the set $\mathcal{O}_{\mathbf{m}} \cap \Gamma_{XY}$ is a single G_σ–orbit.

Moreover, for $\mathbf{m} = (m_{i,\ell}) \in \mathcal{M}_d^+$ the integer $r(\mathbf{m})$ is given by the explicit formula

$$r(\mathbf{m}) \ = \ \sum_{i,\ell} \epsilon(i,\ell)\, m_{i,\ell},$$

where $\epsilon(i,\ell) \in \{0,1\}$ is the $(L \to 1)$–activity indicator of the indecomposable $M(i,\ell)$ as in Proposition 11.6.

Proof. Fix $\sigma \in \operatorname{Hom}(V_L, V_1)$ and recall $\Gamma_{XY} = \Gamma_\sigma$.

First, since $\Gamma_d = \bigsqcup_{\mathbf{m} \in \mathcal{M}_d^+} \mathcal{O}_{\mathbf{m}}$ is a disjoint union, intersecting with Γ_σ gives the disjoint union

$$\Gamma_\sigma \ = \ \Gamma_\sigma \cap \Gamma_d \ = \ \Gamma_\sigma \cap \left(\bigsqcup_{\mathbf{m} \in \mathcal{M}_d^+} \mathcal{O}_{\mathbf{m}} \right) \ = \ \bigsqcup_{\mathbf{m} \in \mathcal{M}_d^+} \big(\mathcal{O}_{\mathbf{m}} \cap \Gamma_\sigma\big).$$

By Proposition 11.5, for a given $\mathbf{m}$ the intersection $\mathcal{O}_{\mathbf{m}} \cap \Gamma_\sigma$ is nonempty if and only if $r(\mathbf{m}) = \operatorname{rank}(\sigma)$. Therefore all terms with $r(\mathbf{m}) \neq \operatorname{rank}(\sigma)$ vanish, and we obtain

$$\Gamma_\sigma \;=\; \bigsqcup_{\mathbf{m}\in\mathcal{M}_d^+\,:\,r(\mathbf{m})=\operatorname{rank}(\sigma)} \big(\mathcal{O}_{\mathbf{m}} \cap \Gamma_\sigma\big).$$

Again by Proposition 11.5, whenever the intersection is nonempty, $\mathcal{O}_{\mathbf{m}} \cap \Gamma_\sigma$ is a single G_σ–orbit. This proves the stated orbit decomposition.

Finally, the explicit formula $r(\mathbf{m}) = \sum_{i,\ell} \epsilon(i,\ell) m_{i,\ell}$ is exactly Proposition 11.6. □

Remark 11.8 (Parametrization of G_σ–orbits in Γ_{XY})**.** The previous corollary can be equivalently phrased as follows. The G_σ–orbits in Γ_{XY} are in bijection with the subset

$$\left\{\mathbf{m} \in \mathcal{M}_d^+ \;\middle|\; r(\mathbf{m}) = \operatorname{rank}(\sigma)\right\},$$

via the correspondence

$$\mathbf{m} \;\longmapsto\; \mathcal{O}_{\mathbf{m}} \cap \Gamma_\sigma.$$

In particular, Γ_{XY} decomposes as a disjoint union of G_σ–orbits indexed by Nakayama indecomposable multiplicity data satisfying the rank constraint.

12 Rank Patterns parameterize orbits of critical points

12.1 Cyclic rank patterns and multiplicities

We work with the cyclic quiver on vertices $\{0, \dots, L\}$ and arrows

$$\phi_\ell : V_{\ell-1} \longrightarrow V_\ell, \qquad \ell = 1, \dots, L+1,$$

where indices are taken modulo $L+1$ (so $\phi_{L+1} : V_L \to V_0$). We impose the cyclic relations described above, so that only oriented paths of length at most L are relevant.

Cyclic intervals. For $i, j \in \{0, \dots, L\}$ we denote by

$$[i \rightsquigarrow j]$$

the oriented segment obtained by following the arrows forward from i to j. Its length is

$$\delta(i,j) \in \{0, \dots, L\},$$

characterized by $j \equiv i + \delta(i,j) \pmod{L+1}$. Thus

$$[i \rightsquigarrow j] = \big(i \xrightarrow{\phi_{i+1}} i+1 \xrightarrow{\phi_{i+2}} \cdots \xrightarrow{\phi_j} j\big).$$

Cyclic rank patterns. Given $\phi = (\phi_1, \dots, \phi_{L+1}) \in \Gamma_d$, for any pair (i, j) with $\delta(i, j) \le L$ we define the composed path map

$$\Phi_{i\to j}(\phi) := \phi_j\, \phi_{j-1} \cdots \phi_{i+1} : V_i \longrightarrow V_j,$$

where indices are taken modulo $L + 1$ and the product is along the oriented segment $[i \rightsquigarrow j]$.

Definition 12.1. The *cyclic rank pattern* of ϕ is the collection of integers

$$r_{ij}(\phi) := \operatorname{rank}\big(\Phi_{i\to j}(\phi)\big), \qquad (i, j) \in \{0, \dots, L\}^2 \text{ with } \delta(i, j) \le L.$$

These ranks are invariant under the base–change action of $G_d = \prod_{k=0}^{L} GL(V_k)$ and therefore are constant on G_d–orbits.

Indecomposables and multiplicities. Let $M(i, j)$ denote the indecomposable cyclic interval module supported on $[i \rightsquigarrow j]$, and write

$$m_{ij}$$

for its multiplicity in the Krull–Schmidt decomposition of the module corresponding to ϕ. Thus the orbit of ϕ in Γ_d is determined by the collection $\{m_{ij}\}$.

By construction, on the indecomposable $M(a, b)$ the map $\Phi_{i\to j}$ has rank 1 if and only if the segment $[i \rightsquigarrow j]$ is contained in $[a \rightsquigarrow b]$, and has rank 0 otherwise. Since ϕ is a direct sum of such interval modules, ranks add over summands. We therefore obtain the cumulative formula

$$r_{ij} = \sum_{\substack{(a,b) \\ [i\rightsquigarrow j]\subseteq[a\rightsquigarrow b]}} m_{ab}. \tag{27}$$

Explicit form. Fix i and write $j = i + t$ with $t = \delta(i, j) \in \{0, \dots, L\}$. An interval $[a \rightsquigarrow b]$ contains $[i \rightsquigarrow j]$ if and only if $a = i$ and $\delta(i, b) \ge t$. Hence (27) reduces to

$$r_{i,\, i+t} = \sum_{\ell=t}^{L} m_{i,\, i+\ell}, \qquad t = 0, \dots, L. \tag{28}$$

Inversion formula. Since (28) is a cumulative sum in the length parameter, it can be inverted by taking first differences. For $\ell = 0, \dots, L$ we obtain

$$m_{i,\, i+\ell} = r_{i,\, i+\ell} - r_{i,\, i+\ell+1}, \tag{29}$$

where by convention $r_{i,\, i+L+1} = 0$.

Equivalently, in (i, j) notation,

$$m_{ij} = r_{ij} - r_{i,\, j+1},$$

with indices taken modulo $L + 1$ and the understanding that $r_{i,\, j+1} = 0$ if $\delta(i, j + 1) > L$.

Thus the cyclic rank pattern $\{r_{ij}\}$ and the multiplicity data $\{m_{ij}\}$ determine one another, and the G_d–orbits in Γ_d may equivalently be parametrized by either set of data.

12.2 Cyclic rank patterns and the bijection $M_d^+ \simeq R_d^{\mathrm{orb}}$

Throughout this subsection we fix $L \geq 1$ and a dimension vector $d = (d_0, \ldots, d_L)$, where $d_i = \dim V_i$. We consider the cyclic quiver on vertices $\{0, \ldots, L\}$ with arrows

$$\phi_\ell : V_{\ell-1} \longrightarrow V_\ell, \qquad \ell = 1, \ldots, L+1,$$

where indices are taken modulo $L+1$ (so $\phi_{L+1} : V_L \to V_0$). We work inside Γ_d, i.e. representations satisfying the cyclic relations introduced earlier (in particular, only oriented paths of length $\leq L$ are relevant).

Cyclic distance and path compositions. For $i, j \in \{0, \ldots, L\}$ we denote by $\delta(i,j) \in \{0, \ldots, L\}$ the (unique) cyclic distance from i to j obtained by following the arrows forward, i.e. the unique integer such that

$$j \equiv i + \delta(i,j) \pmod{L+1}.$$

Thus there is a unique oriented path of length $\delta(i,j)$ from V_i to V_j, namely

$$V_i \xrightarrow{\phi_{i+1}} V_{i+1} \xrightarrow{\phi_{i+2}} \cdots \xrightarrow{\phi_j} V_j,$$

where we interpret indices modulo $L+1$. For $\phi \in \Gamma_d$ we define the corresponding composition

$$\Phi_{ij}(\phi) := \phi_j \, \phi_{j-1} \cdots \phi_{i+1} \ : \ V_i \longrightarrow V_j, \tag{30}$$

with the convention that $\Phi_{ii}(\phi) = \mathrm{id}_{V_i}$.

Cyclic rank patterns.

Definition 12.2. For $\phi \in \Gamma_d$ we define its *cyclic rank pattern* to be the array

$$r(\phi) = (r_{ij}(\phi))_{0 \leq i,j \leq L} \qquad \text{where} \qquad r_{ij}(\phi) := \mathrm{rank}\big(\Phi_{ij}(\phi)\big).$$

Note that $r_{ii}(\phi) = \mathrm{rank}(\mathrm{id}_{V_i}) = d_i$.

We now define an abstract space of rank patterns and an "orbit-realizable" subset.

Definition 12.3. Let R_d be the set of all integer arrays $r = (r_{ij})_{0 \leq i,j \leq L}$ such that

$$r_{ii} = d_i \qquad \text{for all } i \in \{0, \ldots, L\}.$$

Define

$$R_d^{\mathrm{orb}} := \Big\{ r \in R_d \ \Big| \ r_{i,\, i+\ell-1} - r_{i,\, i+\ell} \geq 0 \text{ for all } i \text{ and } \ell = 1, \ldots, L \Big\},$$

where indices are taken modulo $L+1$.

Multiplicity data. Let $M(i,j)$ denote the indecomposable cyclic interval module supported on the oriented segment from i to j (i.e. on the vertices $i, i+1, \ldots, j$ along the cycle), and let $m_{ij} \in \mathbb{Z}_{\geq 0}$ denote its multiplicity in the Krull–Schmidt decomposition of the module corresponding to a point of Γ_d. We write

$$m = (m_{ij}) \in M_d^+$$

for the collection of multiplicities, where M_d^+ denotes the finite set of all such multiplicity collections compatible with the fixed dimension vector d (equivalently, those for which the corresponding direct sum $\bigoplus_{i,j} M(i,j)^{\oplus m_{ij}}$ has dimension vector d).

We now show that M_d^+ is naturally in bijection with R_d^{orb}.

Step 1: the map $M_d^+ \to R_d^{\mathrm{orb}}$

Fix $m \in M_d^+$. For each i, j define

$$r_{ij}(m) := \sum_{k=0}^{L-\delta(i,j)} m_{i,\, j+k}, \tag{31}$$

where indices are taken modulo $L+1$, and the sum is taken over those endpoints $j+k$ for which the oriented segment $[i \rightsquigarrow j+k]$ has length $\delta(i, j+k) \le L$. Equivalently, (31) is the sum over all indecomposables $M(i,b)$ whose support contains the segment $[i \rightsquigarrow j]$.

Lemma 12.4 (Containment criterion)**.** Fix i, j and set $t = \delta(i,j)$. For an endpoint $b \in \{0, \dots, L\}$ with $\delta(i,b) \le L$, the oriented segment $[i \rightsquigarrow j]$ is contained in $[i \rightsquigarrow b]$ if and only if $\delta(i,b) \ge t$, i.e. if and only if $b = j+k$ for some $k \in \{0, \dots, L-t\}$.

Proof. The segment $[i \rightsquigarrow j]$ is the unique oriented path of length t starting at i. The segment $[i \rightsquigarrow b]$ is the unique oriented path of length $\delta(i,b)$ starting at i. Since both start at i and follow the same orientation, $[i \rightsquigarrow j] \subseteq [i \rightsquigarrow b]$ holds exactly when the second path is at least as long as the first, i.e. $\delta(i,b) \ge t$. Writing $b \equiv i + \delta(i,b) \equiv j + (\delta(i,b) - t)$ shows that $b = j+k$ with $k = \delta(i,b) - t \in \{0, \dots, L-t\}$. Conversely, any such k gives $\delta(i, j+k) = t + k \ge t$, hence containment. □

Lemma 12.5 (Forward formula)**.** Let ϕ be any representation whose Krull–Schmidt multiplicities are m. Then for all i, j one has

$$r_{ij}(\phi) = r_{ij}(m),$$

where $r_{ij}(m)$ is defined by (31). In particular, $r(m) := (r_{ij}(m))$ depends only on m and not on the choice of ϕ in the corresponding orbit.

Proof. Decompose the corresponding module as a direct sum of indecomposables: $\mathcal{M} \cong \bigoplus_{a,b} M(a,b)^{\oplus m_{ab}}$. The linear map $\Phi_{ij}(\phi) : V_i \to V_j$ respects this decomposition: on each indecomposable summand $M(a,b)$ it is either the identity on a 1–dimensional space (hence has rank 1) if the segment $[i \rightsquigarrow j]$ is contained in the support of $M(a,b)$, or it is the zero map (hence rank 0) otherwise. Moreover, for fixed i, j, the only summands that can contribute are those with $a = i$ (since the domain is V_i and interval modules have a unique "start" vertex in the oriented string). Therefore

$$\operatorname{rank}(\Phi_{ij}(\phi)) = \sum_{\substack{b:\\ [i \rightsquigarrow j] \subseteq [i \rightsquigarrow b]}} m_{ib}.$$

By Lemma 12.4, the condition $[i \rightsquigarrow j] \subseteq [i \rightsquigarrow b]$ is equivalent to $b = j+k$ with $k \in \{0, \dots, L - \delta(i,j)\}$, which yields (31). □

Lemma 12.6. For any $m \in M_d^+$, the rank pattern $r(m)$ lies in R_d^{orb}.

Proof. First, $r_{ii}(m) = \sum_{k=0}^{L} m_{i,i+k} = d_i$ by the dimension constraint at vertex i, hence $r(m) \in R_d$. Next, fix i and $\ell \in \{1, \dots, L\}$. Using (31) with $j = i + \ell - 1$ and $j' = i + \ell$ gives

$$r_{i,\, i+\ell-1}(m) - r_{i,\, i+\ell}(m) = \Big(\sum_{k=0}^{L-(\ell-1)} m_{i,\, i+\ell-1+k} \Big) - \Big(\sum_{k=0}^{L-\ell} m_{i,\, i+\ell+k} \Big) = m_{i,\, i+\ell-1} \ \ge \ 0,$$

since all other terms cancel by reindexing. Thus $r(m) \in R_d^{\mathrm{orb}}$. □

Define

$$F : M_d^+ \longrightarrow R_d^{\mathrm{orb}}, \qquad F(m) := r(m).$$

Step 2: the inverse map $R_d^{\mathrm{orb}} \to M_d^+$

Let $r \in R_d^{\mathrm{orb}}$. Define integers

$$m_{ij}(r) := r_{ij} - r_{i,\,j+1}, \tag{32}$$

for all i, j, with indices modulo $L+1$ and the convention that $r_{i,\,j+1} = 0$ whenever the segment $[i \rightsquigarrow j+1]$ has length $> L$ (i.e. whenever $\delta(i, j+1) = L+1$ in the untruncated cycle; in practice, this means we set $r_{i,\,i+L+1} := 0$ to terminate the difference). Write $G(r) := (m_{ij}(r))$.

Lemma 12.7. If $r \in R_d^{\mathrm{orb}}$, then $m_{ij}(r) \geq 0$ for all i, j, and $G(r) \in M_d^+$.

Proof. Fix i, j and write $j = i + \ell$ with $\ell = \delta(i, j) \in \{0, \ldots, L\}$. Then (32) reads

$$m_{i,\,i+\ell}(r) = r_{i,\,i+\ell} - r_{i,\,i+\ell+1}.$$

By definition of R_d^{orb} we have $r_{i,\,i+\ell} - r_{i,\,i+\ell+1} \geq 0$, hence $m_{ij}(r) \geq 0$.

It remains to verify the dimension constraint at each vertex i. Using telescoping sums and the convention $r_{i,\,i+L+1} = 0$ we obtain

$$\sum_{\ell=0}^{L} m_{i,\,i+\ell}(r) = \sum_{\ell=0}^{L} \big(r_{i,\,i+\ell} - r_{i,\,i+\ell+1}\big) = r_{ii} - r_{i,\,i+L+1} = d_i.$$

Thus the multiplicities $m_{ij}(r)$ are nonnegative and satisfy the required dimension constraints, so $G(r) \in M_d^+$. □

Step 3: F and G are inverse bijections

Proposition 12.8. The maps

$$F : M_d^+ \to R_d^{\mathrm{orb}} \qquad \text{and} \qquad G : R_d^{\mathrm{orb}} \to M_d^+$$

defined above are mutually inverse. In particular,

$$M_d^+ \xrightarrow{\sim} R_d^{\mathrm{orb}}.$$

Proof. (i) $G \circ F = \mathrm{id}_{M_d^+}$. Let $m \in M_d^+$ and set $r = F(m)$. Fix i and write $j = i + \ell$ with $\ell = \delta(i, j) \in \{0, \ldots, L\}$. Then by (31),

$$r_{i,\,i+\ell} = \sum_{k=0}^{L-\ell} m_{i,\,i+\ell+k}, \qquad r_{i,\,i+\ell+1} = \sum_{k=0}^{L-(\ell+1)} m_{i,\,i+\ell+1+k}.$$

Subtracting yields a telescoping cancellation:

$$r_{i,\,i+\ell} - r_{i,\,i+\ell+1} = m_{i,\,i+\ell}.$$

By (32), this is exactly $m_{ij}(G(r)) = m_{ij}$, hence $G(F(m)) = m$.

(ii) $F \circ G = \mathrm{id}_{R_d^{\mathrm{orb}}}$. Let $r \in R_d^{\mathrm{orb}}$ and set $m = G(r)$. Fix i and write $j = i + \ell$ with $\ell = \delta(i,j) \in \{0, \dots, L\}$. Using (31) and (32),

$$(F(m))_{i,\, i+\ell} = \sum_{k=0}^{L-\ell} m_{i,\, i+\ell+k} = \sum_{k=0}^{L-\ell} \big(r_{i,\, i+\ell+k} - r_{i,\, i+\ell+k+1}\big).$$

This sum telescopes to

$$r_{i,\, i+\ell} - r_{i,\, i+L+1}.$$

By convention $r_{i,\, i+L+1} = 0$, hence $(F(m))_{i,\, i+\ell} = r_{i,\, i+\ell}$. Thus $F(G(r)) = r$.

Together, (i) and (ii) prove that F and G are inverse bijections. □

Remark (comparison with the linear case). In the equioriented A_n (linear) setting, rank patterns are indexed by all intervals $[i,j]$ and the relation between rank data and Kostant multiplicities is given by a two-dimensional inclusion–exclusion formula. In the present cyclic (Nakayama) setting with the above relations, indecomposable summands are indexed by cyclic intervals with a distinguished start vertex. Consequently, the forward map (31) becomes a one-sided cumulative sum along each fixed start vertex i, and the inversion (32) is a first-difference (telescoping) formula.

12.3 Orbit decomposition of Γ_{XY} via cyclic rank patterns and the constraint $r(m) = \sigma$

Setup. Fix vector spaces $V_0, \dots, V_L$ with $\dim V_i = d_i$ and consider the cyclic quiver on vertices $\{0, \dots, L\}$ with arrows

$$\phi_\ell : V_{\ell-1} \longrightarrow V_\ell, \qquad \ell = 1, \dots, L+1,$$

where indices are taken modulo $L+1$ (so $\phi_{L+1} : V_L \to V_0$). Let $\Gamma_d \subset \mathrm{Rep}_d(C_{L+1})$ be the closed subvariety cut out by the cyclic relations introduced earlier. We fix a linear map

$$\sigma \in \mathrm{Hom}(V_L, V_0)$$

and define the fibre

$$\Gamma_{XY} := \{\phi \in \Gamma_d \mid \phi_{L+1} = \sigma\}.$$

Let

$$G_d := \prod_{i=0}^{L} GL(V_i)$$

act on Γ_d by base change. Since σ is fixed, the natural group acting on Γ_{XY} is the stabilizer of σ under the $(0,L)$-component action:

$$G_\sigma := \{(g_0, \dots, g_L) \in G_d \mid g_0\, \sigma = \sigma\, g_L\}.$$

Step 1: G_d–orbit decomposition of Γ_d. By the finiteness of the representation type in the present cyclic (Nakayama) setting, the G_d–orbits in Γ_d are parametrized by multiplicity data

$$m = (m_{ij}) \in M_d^+,$$

where $m_{ij} \in \mathbb{Z}_{\geq 0}$ denotes the multiplicity of the indecomposable interval module $M(i,j)$ supported on the oriented segment $[i \rightsquigarrow j]$. We denote by $\mathcal{O}_m \subset \Gamma_d$ the corresponding G_d–orbit. Then

$$\Gamma_d = \bigsqcup_{m \in M_d^+} \mathcal{O}_m. \tag{33}$$

Step 2: Passage to the fibre Γ_{XY} and G_σ–orbits. Consider the morphism

$$F : \Gamma_d \longrightarrow \mathrm{Hom}(V_L, V_0), \qquad F(\phi) := \phi_{L+1}.$$

The map F is G_d–equivariant with respect to the natural action of G_d on $\mathrm{Hom}(V_L, V_0)$ given by $(g_0, \ldots, g_L) \cdot A = g_0 A g_L^{-1}$.

For each $m \in M_d^+$, the rank

$$r(m) := \mathrm{rank}(\phi_{L+1}) \qquad (\phi \in \mathcal{O}_m)$$

is well-defined (independent of $\phi \in \mathcal{O}_m$) because rank is invariant under left–right base change. Equivalently, $F(\mathcal{O}_m)$ is contained in the rank stratum

$$\mathrm{Hom}(V_L, V_0)_{r(m)} := \{A \in \mathrm{Hom}(V_L, V_0) \mid \mathrm{rank}(A) = r(m)\}.$$

Since over $\mathbb{R}$ (or $\mathbb{C}$) the orbits of $GL(V_0) \times GL(V_L)$ in $\mathrm{Hom}(V_L, V_0)$ are precisely the rank strata, we have:

$$\mathcal{O}_m \cap \Gamma_{XY} \neq \varnothing \quad \Longleftrightarrow \quad r(m) = \mathrm{rank}(\sigma). \tag{34}$$

Moreover, whenever $\mathcal{O}_m \cap \Gamma_{XY} \neq \varnothing$, this intersection is a single G_σ–orbit (equivalently: the fibre of a G_d–equivariant map over a point in a homogeneous G_d–orbit is homogeneous under the stabilizer of that point). Therefore,

$$\Gamma_{XY} = \bigsqcup_{\substack{m \in M_d^+ \\ r(m) = \mathrm{rank}(\sigma)}} \big(\mathcal{O}_m \cap \Gamma_{XY}\big), \qquad \text{and each } \mathcal{O}_m \cap \Gamma_{XY} \text{ is a single } G_\sigma\text{–orbit.} \tag{35}$$

Step 3: Reparametrization by cyclic rank patterns. For $\phi \in \Gamma_d$ we define, for all $i, j \in \{0, \ldots, L\}$, the composed map

$$\Phi_{ij}(\phi) := \phi_j \phi_{j-1} \cdots \phi_{i+1} : V_i \to V_j$$

(along the unique oriented path $i \rightsquigarrow j$), and the cyclic rank pattern

$$r_{ij}(\phi) := \mathrm{rank}\big(\Phi_{ij}(\phi)\big).$$

As proved in the previous subsection, the assignment $\phi \mapsto (r_{ij}(\phi))$ is constant on G_d–orbits and yields a bijection

$$M_d^+ \xrightarrow{\sim} R_d^{\mathrm{orb}}, \qquad m \longmapsto r(m) := (r_{ij}(m)),$$

with inverse given by the first-difference formula

$$m_{ij} = r_{ij} - r_{i,j+1}, \tag{36}$$

where $j+1$ is taken modulo $L+1$, and we adopt the terminating convention

$$r_{i,i+L+1} := 0 \qquad \text{(so that telescoping identities are valid).}$$

Combining (35) with the bijection $M_d^+ \simeq R_d^{\mathrm{orb}}$, we may equivalently index the G_σ–orbits in Γ_{XY} by rank patterns $r \in R_d^{\mathrm{orb}}$ subject to a single additional scalar constraint, obtained by translating $r(m) = \mathrm{rank}(\sigma)$ into rank-pattern language.

Step 4: The constraint $r(m) = \mathrm{rank}(\sigma)$ in rank-pattern coordinates. For each indecomposable interval module $M(i,j)$ (equivalently, $M(i,i+\ell)$ with $\ell = \delta(i,j)$), the distinguished arrow $\phi_{L+1} : V_L \to V_0$ acts either trivially or nontrivially. Define the indicator

$$\epsilon(i,j) \in \{0,1\}$$

by

$\epsilon(i,j) = 1 \iff$ the arrow $(L \to 0)$ lies in the support segment $[i \rightsquigarrow j]$ (equivalently: ϕ_{L+1} is active on $M(i,j)$).

(Equivalently: $[i \rightsquigarrow j]$ contains the step $L \rightsquigarrow 0$.)

Lemma 12.9. For $m \in M_d^+$ one has

$$r(m) = \mathrm{rank}(\phi_{L+1}) = \sum_{i,j} \epsilon(i,j)\, m_{ij}, \tag{37}$$

where the sum ranges over all (i,j) with $\delta(i,j) \le L$ (i.e. over all indecomposables present).

Proof. Decompose the corresponding module as a direct sum of indecomposables: $\mathcal{M} \cong \bigoplus_{i,j} M(i,j)^{\oplus m_{ij}}$. The map ϕ_{L+1} is block-diagonal with respect to this decomposition, and on each summand $M(i,j)$ it has rank 1 precisely when the arrow $L \to 0$ belongs to the support of $M(i,j)$, and rank 0 otherwise. Thus the total rank of ϕ_{L+1} equals the number of summands (counted with multiplicity) on which it is active, i.e. the sum (37). □

Now substitute (36) into (37). Writing $j = i+\ell$ with $\ell = \delta(i,j)$, we obtain, for the rank pattern $r = (r_{ij}) \in R_d^{\mathrm{orb}}$ corresponding to m,

$$\mathrm{rank}(r) := \sum_{i,j} \epsilon(i,j)\, m_{ij} = \sum_{i,j} \epsilon(i,j)\, \big(r_{ij} - r_{i,j+1}\big). \tag{38}$$

Equivalently, grouping by the length parameter $\ell \in \{0,\dots,L\}$ and using the terminating convention $r_{i,i+L+1} = 0$, one may write

$$\mathrm{rank}(r) = \sum_{i=0}^{L} \sum_{\ell=0}^{L} \epsilon(i,i+\ell)\, \big(r_{i,\,i+\ell} - r_{i,\,i+\ell+1}\big) = \sum_{i=0}^{L} \sum_{\ell=0}^{L} \epsilon(i,i+\ell)\, r_{i,\,i+\ell} \;-\; \sum_{i=0}^{L} \epsilon(i,i+L)\, r_{i,\,i+L+1}. \tag{39}$$

Since $r_{i,i+L+1} = 0$ by convention, the final term vanishes; in particular one may record the schematic identity

$$\mathrm{rank}(r) = \sum_{i,\ell} \epsilon(i,i+\ell)\, m_{i,\,i+\ell} = \sum_{i,\ell} \epsilon(i,i+\ell)\, \big(r_{i,\,i+\ell} - r_{i,\,i+\ell+1}\big),$$

which is the rank-pattern form of the constraint $r(m) = \mathrm{rank}(\sigma)$.

Conclusion: orbit decomposition of Γ_{XY} by rank patterns. Define the rank-pattern admissible locus

$$R_d^{\mathrm{orb}}(\sigma) := \{\, r \in R_d^{\mathrm{orb}} \mid \mathrm{rank}(r) = \mathrm{rank}(\sigma) \,\},$$

where $\mathrm{rank}(r)$ is given by (38) (equivalently (39)). Then (35) becomes the rank-pattern stratification

$$\Gamma_{XY} = \bigsqcup_{r \in R_d^{\mathrm{orb}}(\sigma)} \Gamma_{XY}^{(r)}, \tag{40}$$

where each stratum $\Gamma_{XY}^{(r)}$ is a single G_σ–orbit, namely

$$\Gamma_{XY}^{(r)} = \mathcal{O}_m \cap \Gamma_{XY} \quad \text{for the unique } m \in M_d^+ \text{ with } F(m) = r.$$

Thus Γ_{XY} decomposes into G_σ–orbits indexed by cyclic rank patterns $r \in R_d^{\mathrm{orb}}$ satisfying the additional scalar constraint $\mathrm{rank}(r) = \mathrm{rank}(\sigma)$, which is equivalently the condition

$$\sum_{i,j} \epsilon(i,j)\, m_{ij} = \mathrm{rank}(\sigma) \qquad \Longleftrightarrow \qquad \sum_{i,j} \epsilon(i,j)\,(r_{ij} - r_{i,j+1}) = \mathrm{rank}(\sigma).$$

Proposition 12.10 (Orbit stratification of Γ_{XY} by cyclic rank patterns)**.** Fix $\sigma \in \mathrm{Hom}(V_L, V_0)$ and set

$$\Gamma_{XY} := \{\phi \in \Gamma_d \mid \phi_{L+1} = \sigma\}.$$

Let R_d^{orb} be the set of orbit-realizable cyclic rank patterns, and for $r \in R_d^{\mathrm{orb}}$ let $\mathcal{O}_r \subset \Gamma_d$ denote the corresponding G_d–orbit (equivalently, $\mathcal{O}_r = \mathcal{O}_m$ where $m \in M_d^+$ is the unique multiplicity datum mapping to r under the bijection $M_d^+ \simeq R_d^{\mathrm{orb}}$).

For each $r \in R_d^{\mathrm{orb}}$ define the *σ–restricted orbit*

$$\mathcal{O}_{r,\sigma} := \mathcal{O}_r \cap \Gamma_{XY}.$$

Then:

1. For every $r \in R_d^{\mathrm{orb}}$, the subset $\mathcal{O}_{r,\sigma}$ is either empty or a single G_σ–orbit (where $G_\sigma = \{(g_0, \ldots, g_L) \in G_d \mid g_0\sigma = \sigma g_L\}$).
2. $\mathcal{O}_{r,\sigma} \neq \varnothing$ if and only if the scalar constraint

   $$\mathrm{rank}(r) = \mathrm{rank}(\sigma)$$

   holds, where $\mathrm{rank}(r)$ is defined by

   $$\mathrm{rank}(r) := \sum_{i=0}^{L} \sum_{\ell=1}^{L} \epsilon(i, i+\ell)\, m_{i,\, i+\ell} = \sum_{i=0}^{L} \sum_{\ell=1}^{L} \epsilon(i, i+\ell)\, \big(r_{i,\, i+\ell} - r_{i,\, i+\ell+1}\big), \tag{41}$$

   with $m_{ij} = r_{ij} - r_{i,j+1}$ and the convention $r_{i,\, i+L+1} = 0$.
3. Consequently, Γ_{XY} admits the disjoint union decomposition

   $$\Gamma_{XY} = \bigsqcup_{\substack{r \in R_d^{\mathrm{orb}} \\ \mathrm{rank}(r) = \mathrm{rank}(\sigma)}} \mathcal{O}_{r,\sigma}. \tag{42}$$

   In particular, Γ_{XY} is stratified by G_σ–orbits indexed by cyclic rank patterns $r \in R_d^{\mathrm{orb}}$ satisfying $\mathrm{rank}(r) = \mathrm{rank}(\sigma)$.

13 Codimension of $\mathcal{C}_r$

13.1 Relation with codimension of orbits

Proposition 13.1 (Codimension of Γ_{XY} via the largest cyclic orbit)**.** Let Γ_{XY} be stratified by cyclic rank patterns as in Proposition 12.10:

$$\Gamma_{XY} \;=\; \bigsqcup_{\substack{r\in R_d^{\mathrm{orb}}\\ \mathrm{rank}(r)=\mathrm{rank}(\sigma)}} \mathcal{O}_{r,\sigma}, \qquad \mathcal{O}_{r,\sigma} := \mathcal{O}_r \cap \Gamma_{XY},$$

where each nonempty $\mathcal{O}_{r,\sigma}$ is a single G_σ-orbit.

Let

$$\mathcal{A}(\sigma) \;:=\; \big\{\, r \in R_d^{\mathrm{orb}} \;\big|\; \mathrm{rank}(r) = \mathrm{rank}(\sigma) \text{ (equivalently, } \mathcal{O}_{r,\sigma} \neq \varnothing)\big\}.$$

Then

$$\mathrm{codim}(\Gamma_{XY}) \;=\; \min_{r\in\mathcal{A}(\sigma)} \mathrm{codim}(\mathcal{O}_{r,\sigma}). \tag{43}$$

Equivalently, if $r_* \in \mathcal{A}(\sigma)$ is such that

$$\dim \mathcal{O}_{r_*,\sigma} = \max_{r\in\mathcal{A}(\sigma)} \dim \mathcal{O}_{r,\sigma},$$

then $\overline{\mathcal{O}_{r_*,\sigma}}$ is an irreducible component of Γ_{XY} of maximal dimension and

$$\mathrm{codim}(\Gamma_{XY}) = \mathrm{codim}(\mathcal{O}_{r_*,\sigma}). \tag{44}$$

Proof. We work throughout in the Zariski topology on the affine variety Γ_d (hence on the closed subvariety $\Gamma_{XY} = \{\phi \in \Gamma_d : \phi_{L+1} = \sigma\}$). By Proposition 12.10,

$$\Gamma_{XY} \;=\; \bigsqcup_{r\in\mathcal{A}(\sigma)} \mathcal{O}_{r,\sigma}, \qquad \mathcal{A}(\sigma) := \{\, r \in R_d^{\mathrm{orb}} :\; \mathcal{O}_{r,\sigma} \neq \varnothing \,\} = \{\, r \in R_d^{\mathrm{orb}} :\; \mathrm{rank}(r) = \mathrm{rank}(\sigma) \,\}, \tag{45}$$

and each nonempty $\mathcal{O}_{r,\sigma}$ is a single G_σ-orbit. In particular, each $\mathcal{O}_{r,\sigma}$ is a locally closed, irreducible subset of Γ_{XY} (indeed, an orbit of an algebraic group action is constructible and irreducible, being the image of the irreducible variety G_σ under the orbit map).

Step 1: Dimension of Γ_{XY} is the maximal orbit dimension. Since the indexing set $\mathcal{A}(\sigma)$ is finite (cyclic rank patterns take values in a finite set of integers bounded by the widths), (45) expresses Γ_{XY} as a finite disjoint union of locally closed subsets. For any finite family of constructible subsets $\{S_i\}_{i=1}^N$ of an algebraic variety, one has

$$\dim\Big(\bigcup_{i=1}^N S_i\Big) \;=\; \max_{1\le i\le N} \dim S_i,$$

because taking Zariski closures gives $\overline{\bigcup_i S_i} = \bigcup_i \overline{S_i}$, and the dimension of a finite union of closed subsets is the maximum of their dimensions. Applying this to the family $\{\mathcal{O}_{r,\sigma}\}_{r\in\mathcal{A}(\sigma)}$ yields

$$\dim(\Gamma_{XY}) \;=\; \max_{r\in\mathcal{A}(\sigma)} \dim(\mathcal{O}_{r,\sigma}). \tag{46}$$

Step 2: Codimension is the minimal orbit codimension. By definition, for any constructible $S \subseteq \Gamma_d$ we set

$$\operatorname{codim}_{\Gamma_d}(S) \;:=\; \dim(\Gamma_d) - \dim(S).$$

Combining this with (46) gives

$$\operatorname{codim}_{\Gamma_d}(\Gamma_{XY}) \;=\; \dim(\Gamma_d) - \max_{r\in\mathcal{A}(\sigma)} \dim(\mathcal{O}_{r,\sigma}) \;=\; \min_{r\in\mathcal{A}(\sigma)} \big(\dim(\Gamma_d) - \dim(\mathcal{O}_{r,\sigma})\big) \;=\; \min_{r\in\mathcal{A}(\sigma)} \operatorname{codim}_{\Gamma_d}(\mathcal{O}_{r,\sigma}),$$

which is exactly (43).

Step 3: Realization by a largest orbit and maximal component. Let $r_* \in \mathcal{A}(\sigma)$ achieve the maximum in (46). Since $\mathcal{O}_{r_*,\sigma}$ is irreducible, its Zariski closure $\overline{\mathcal{O}_{r_*,\sigma}}$ is irreducible as well. Moreover, $\overline{\mathcal{O}_{r_*,\sigma}} \subseteq \Gamma_{XY}$ and

$$\dim \overline{\mathcal{O}_{r_*,\sigma}} = \dim \mathcal{O}_{r_*,\sigma} = \dim \Gamma_{XY}.$$

Therefore $\overline{\mathcal{O}_{r_*,\sigma}}$ is an irreducible component of Γ_{XY} of maximal dimension, and

$$\operatorname{codim}_{\Gamma_d}(\Gamma_{XY}) = \dim(\Gamma_d) - \dim(\Gamma_{XY}) = \dim(\Gamma_d) - \dim(\mathcal{O}_{r_*,\sigma}) = \operatorname{codim}_{\Gamma_d}(\mathcal{O}_{r_*,\sigma}),$$

which is (44). □

Proposition 13.2 (Codimension of C_r via cyclic orbits in Γ_{XY})**.** Fix $0 \le r \le \min_\ell d_\ell$ and retain the notation of Theorem 8.5 and Proposition 12.11. Assume Γ_{XY} is identified with the residual constraint variety $\mathcal{X}_{r,\sigma} \subset \mathcal{Z}$ (i.e. $\Gamma_{XY} = \mathcal{X}_{r,\sigma}$ under the fixed endpoint/closing-arrow model), and assume the principal slice locus is dense so that

$$\operatorname{codim}_\Omega(C_r^\circ) = \operatorname{codim}_\Omega(C_r).$$

Let Γ_{XY} admit the orbit stratification

$$\Gamma_{XY} = \bigsqcup_{\substack{m\in R_d^{\mathrm{orb}} \\ \operatorname{rank}(m)=\operatorname{rank}(\sigma)}} \mathcal{O}_{m,\sigma}, \qquad \mathcal{O}_{m,\sigma} := \mathcal{O}_m \cap \Gamma_{XY},$$

and let m_* be a cyclic rank pattern of maximal orbit dimension among those with $\operatorname{rank}(m) = \operatorname{rank}(\sigma)$. Then

$$\operatorname{codim}_\Omega(C_r) \;=\; \operatorname{codim}_{\mathcal{Z}}(\mathcal{O}_{m_*,\sigma}) \;+\; r(d_0 + d_L). \tag{47}$$

Equivalently,

$$\operatorname{codim}_\Omega(C_r) \;=\; \min_{\substack{m\in R_d^{\mathrm{orb}} \\ \operatorname{rank}(m)=\operatorname{rank}(\sigma)}} \operatorname{codim}_{\mathcal{Z}}(\mathcal{O}_{m,\sigma}) \;+\; r(d_0 + d_L). \tag{48}$$

Proof. We fix the ambient affine space Ω for C_r and the ambient affine space $\mathcal{Z}$ for $\Gamma_{XY} = \mathcal{X}_{r,\sigma}$, and we use Zariski codimension throughout:

$$\operatorname{codim}_\Omega(S) := \dim(\Omega) - \dim(S), \qquad \operatorname{codim}_{\mathcal{Z}}(T) := \dim(\mathcal{Z}) - \dim(T),$$

for constructible subsets $S \subseteq \Omega$, $T \subseteq \mathcal{Z}$.

Step 1: Reduce $\operatorname{codim}_\Omega(C_r)$ **to** $\operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma})$**.** By Proposition 8.8 (codimension comparison between critical points and the residual constraint variety), on the principal slice locus one has

$$\operatorname{codim}_\Omega(C_r) = \operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma}) + r(d_0 + d_L), \tag{49}$$

where we use the standing hypothesis that $\operatorname{codim}_\Omega(C_r^\circ) = \operatorname{codim}_\Omega(C_r)$ so that the codimension can be computed on the principal locus.

Step 2: Compute $\operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma})$ **from the orbit stratification.** By the hypothesis of the proposition, Γ_{XY} is identified with $\mathcal{X}_{r,\sigma}$ as a subvariety of $\mathcal{Z}$. By Proposition 12.11 (orbit stratification by cyclic rank patterns), there is a finite disjoint union

$$\mathcal{X}_{r,\sigma} = \Gamma_{XY} = \bigsqcup_{\substack{m \in R_d^{\mathrm{orb}} \\ \operatorname{rank}(m) = \operatorname{rank}(\sigma)}} \mathcal{O}_{m,\sigma}, \qquad \mathcal{O}_{m,\sigma} := \mathcal{O}_m \cap \Gamma_{XY}, \tag{50}$$

and each nonempty $\mathcal{O}_{m,\sigma}$ is a single G_σ-orbit in Γ_{XY}; in particular each $\mathcal{O}_{m,\sigma}$ is a locally closed, irreducible, constructible subset of $\mathcal{Z}$.

Since the index set in (50) is finite, we have the general dimension identity for finite unions of constructible sets:

$$\dim\left(\bigcup_{i=1}^{N} S_i\right) = \max_{1 \le i \le N} \dim(S_i).$$

Applying this with the family of strata $\{\mathcal{O}_{m,\sigma}\}$ gives

$$\dim(\mathcal{X}_{r,\sigma}) = \max_{\substack{m \in R_d^{\mathrm{orb}} \\ \operatorname{rank}(m) = \operatorname{rank}(\sigma)}} \dim(\mathcal{O}_{m,\sigma}). \tag{51}$$

(Indeed, taking Zariski closures yields $\overline{\mathcal{X}_{r,\sigma}} = \mathcal{X}_{r,\sigma} = \bigcup_m \overline{\mathcal{O}_{m,\sigma}}$, and the dimension of a finite union of closed sets equals the maximum of the dimensions of the closed sets; moreover $\dim \overline{\mathcal{O}_{m,\sigma}} = \dim(\mathcal{O}_{m,\sigma})$ because $\mathcal{O}_{m,\sigma}$ is irreducible and locally closed.)

Now translate (51) into codimensions inside $\mathcal{Z}$:

$$\operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma}) = \dim(\mathcal{Z}) - \dim(\mathcal{X}_{r,\sigma}) = \dim(\mathcal{Z}) - \max_{\operatorname{rank}(m) = \operatorname{rank}(\sigma)} \dim(\mathcal{O}_{m,\sigma}) = \min_{\operatorname{rank}(m) = \operatorname{rank}(\sigma)} \big(\dim(\mathcal{Z}) - \dim(\mathcal{O}_{m,\sigma})\big).$$

Hence

$$\operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma}) = \min_{\substack{m \in R_d^{\mathrm{orb}} \\ \operatorname{rank}(m) = \operatorname{rank}(\sigma)}} \operatorname{codim}_{\mathcal{Z}}(\mathcal{O}_{m,\sigma}). \tag{52}$$

Equivalently, if m_* achieves the maximum orbit dimension in (51), then it achieves the minimum codimension in (52) and

$$\operatorname{codim}_{\mathcal{Z}}(\mathcal{X}_{r,\sigma}) = \operatorname{codim}_{\mathcal{Z}}(\mathcal{O}_{m_*,\sigma}). \tag{53}$$

Step 3: Combine the two identities. Substituting (52) into (49) yields

$$\operatorname{codim}_\Omega(C_r) = \min_{\substack{m \in R_d^{\mathrm{orb}} \\ \operatorname{rank}(m) = \operatorname{rank}(\sigma)}} \operatorname{codim}_{\mathcal{Z}}(\mathcal{O}_{m,\sigma}) + r(d_0 + d_L),$$

which is (48). Substituting (53) into (49) yields

$$\operatorname{codim}_\Omega(C_r) = \operatorname{codim}_{\mathcal{Z}}(\mathcal{O}_{m_*,\sigma}) + r(d_0 + d_L),$$

which is (47). This proves the proposition. □