

Dantzig-Wolfe decomposition

Guillaume Dalle

MPRO – Decomposition methods for integer programming

All of the content is taken from [1] or [2].

Contents

I.	Motivation: complicating constraints	1
II.	Dantzig-Wolfe relaxation	3
II.1.	Definition	3
II.2.	Extended formulation	4
II.3.	Examples	5
III.	Column generation	6
III.1.	Main problem and pricing	6
III.2.	Column generation	8
III.3.	Examples	8
IV.	Branch and price	9
IV.1.	Branching techniques	9
IV.2.	Implementation tricks	9
	Bibliography	9

I. Motivation: complicating constraints

For each of the following problems:

1. Give a mixed integer programming formulation.
2. Describe the structure of the constraint matrix.

Generalized assignment – modeling

Exercise 1

There are a number of tasks $i \in [m]$ to be performed by agents $j \in [n]$. Assigning task i to agent j generates profit c_{ij} and takes time t_{ij} . Each agent has a time budget T_j . Agents do not collaborate on tasks. The goal is to maximize the profit subject to time constraints.

Let x_{ij} be a binary variable equal to 1 if task i is assigned to agent j . The GAP can be formulated as:

$$\max_x \sum_{i,j} c_{ij} x_{ij} \quad \text{s.t.} \quad \begin{cases} \sum_j x_{ij} \leq 1 & \forall i \\ \sum_i t_{ij} x_{ij} \leq T_j & \forall j \\ x_{ij} \in \{0, 1\} & \forall (i, j) \end{cases} \quad (1)$$

The constraint matrix has a block diagonal structure with one block per agent j and a coupling block on top (at most one agent per task).

Cutting stock – modeling

Exercise 2

At a paper mill, we have large rolls of paper of width W , to be cut into smaller rolls. The client's order is to produce b_i rolls of width w_i for each $i \in [m]$. The goal is to minimize the number of large rolls necessary.

We assume that at most p large rolls will be necessary. Let y_j be a binary variable equal to 1 if roll $j \in [p]$ is used, and z_{ij} be an integer variable denoting the number of small rolls of width w_i cut out from large roll j . The CSP can be formulated as:

$$\min_{y,z} \sum_j y_j \quad \text{s.t.} \quad \begin{cases} \sum_i w_i z_{ij} \leq W y_j & \forall j \\ \sum_j z_{ij} \geq b_i & \forall i \\ y_j \in \{0, 1\} & \forall j \\ z_{i,j} \in \mathbb{N} & \forall (i, j) \end{cases} \quad (2)$$

The constraint matrix has a block diagonal structure with one block per large roll j and a coupling block on top (demand satisfaction).

Facility location – modeling

Exercise 3

A company would like to set up facilities to serve its customers. Each customer $i \in [m]$ has annual demand d_i . For each possible facility location $j \in [n]$, the fixed annual operating cost is f_j and the cost of serving client i is c_{ij} . The goal is to select which facilities to open in order to minimize total annual costs.

Let x_j be a binary variable equal to 1 if facility j is open, and y_{ij} be the fraction of demand d_i transported from j to i . The FLP can be formulated as:

$$\min_{x,y} \sum_{i,j} c_{ij} y_{ij} d_i \quad \text{s.t.} \quad \begin{cases} \sum_j y_{ij} = 1 & \forall i \\ \sum_i y_{ij} \leq x_j & \forall (i, j) \\ x_j \in \{0, 1\} & \forall j \\ y_{i,j} \geq 0 & \forall (i, j) \end{cases} \quad (3)$$

The constraint matrix has a block diagonal structure with one block per facility j and a coupling block on top (serving clients).

Network design – modeling

Exercise 4

Let $G = (N, A)$ be a directed graph of potential arcs to be constructed. We need to select a subset of arcs from A to route commodities $k \in [K]$. Each commodity k goes from source s_k to target t_k with volume v_k . Each arc $a \in A$ has capacity c_a and construction cost f_a . The goal is to design a network at minimum cost that is able to route all the flow.

Let x_a be a binary variable equal to 1 if arc a is constructed. Let y_a^k denote the amount of commodity k flowing through a . The NDP can be formulated as:

$$\min_{x,y} \sum_a f_a x_a \quad \text{s.t.} \quad \begin{cases} \sum_{a \in \delta^+(i)} y_a^k - \sum_{a \in \delta^-(i)} y_a^k = (1_{i=s_k} - 1_{i=t_k}) v_k & \forall i \\ \sum_k y_a^k \leq c_a x_a & \forall a \\ x_a \in \{0,1\} & \forall j \\ y_a^k \geq 0 & \forall (a,k) \end{cases} \quad (4)$$

There are two ways to look at this constraint matrix:

- Block structure in the commodities
- Block structure in the arcs

Parallel subproblems

Idea 5

In each of these problems, removing the coupling constraint would allow parallel solution of simpler subproblems. Can we use this insight to obtain a better relaxation for branch & bound?

II. Dantzig-Wolfe relaxation

II.1. Definition

Setting

Definition 6

Consider the following integer program with rational coefficients:

$$z_{\text{ILP}} = \max_x c^\top x \quad \text{s.t.} \quad \begin{cases} A_1 x \leq b_1 & \text{(hard constraint)} \\ A_2 x \leq b_2 & \text{(easy constraint)} \\ x \in \mathbb{Z}^n & \text{(integrality)} \end{cases} \quad (5)$$

and its continuous relaxation

$$z_{\text{LP}} = \max_x c^\top x \quad \text{s.t.} \quad \begin{cases} A_1 x \leq b_1 \\ A_2 x \leq b_2 \end{cases} \quad (6)$$

We define $S = \{x \in \mathbb{Z}^n : A_1 x \leq b_1, A_2 x \leq b_2\}$ the integer feasible set and $Q = \{x \in \mathbb{Z}^n : A_2 x \leq b_2\}$ the integer feasible set with the hard constraint relaxed. Furthermore, we assume that problems of the form $\max\{\tilde{c}^\top x : x \in Q\}$ are easy to solve.

Dantzig-Wolfe relaxation

Definition 7

The Dantzig-Wolfe relaxation of the ILP in Equation 5 is defined as

$$z_{\text{DW}} = \max_x c^\top x \quad \text{s.t.} \quad \begin{cases} A_1 x \leq b_1 \\ x \in \text{conv}(Q) \end{cases} \quad (7)$$

Often better than the continuous relaxation

Proposition 8

We have $z_{\text{ILP}} \leq z_{\text{DW}} \leq z_{\text{LP}}$.

The feasible sets of these three problems satisfy

$$S = \{A_1x \leq b_1\} \cap Q \subseteq \{A_1x \leq b_1\} \cap \text{conv}(Q) \leq \{A_1x \leq b_1\} \cap \{A_2x \leq b_2\} \quad (8)$$

Not always

Exercise 9

In which cases is the Dantzig-Wolfe relaxation equal to the linear programming relaxation?

When $\text{conv}(Q) = \{A_2x \leq b_2\}$, the Dantzig-Wolfe relaxation does not give a better bound than the linear programming relaxation (but it may still be easier to compute, see below).

II.2. Extended formulation

Extended formulation

Proposition 10

There exist v_k and r_h such that the Dantzig-Wolfe relaxation can be formulated as

$$\begin{aligned} \max_{\lambda, \mu} \quad & \sum_{k \in K} \lambda_k c^\top v_k + \sum_{h \in H} \mu_h c^\top r_h \\ \text{s.t.} \quad & \sum_{k \in K} \lambda_k A_1 v_k + \sum_{h \in H} \mu_h A_1 r_h \leq b_1 \\ & \sum_{k \in K} \lambda_k = 1 \\ & \lambda \in \mathbb{R}_+^K, \mu \in \mathbb{R}_+^H, \end{aligned} \quad (9)$$

By Meyer's theorem, $\text{conv}(Q)$ is a rational polyhedron. By the Minkowski-Weyl theorem, it can be written as the sum of a polytope and a finitely-generated cone.

Reformulation, general case

Remark 11

To obtain a reformulation of the original ILP, just add the following constraint to the DW relaxation:

$$\sum_{k \in K} \lambda_k A_1 v_k + \sum_{h \in H} \mu_h A_1 r_h \in \mathbb{Z}^n \quad (10)$$

Reformulation, simple case

Remark 12

When Q is bounded, we can also pick v_k to be all points in Q (not just vertices), and then a solution to Equation 7 is integral if $\lambda \in \{0, 1\}^K$.

Many practical problems look like this (considering the pure binary case for simplicity):

$$\begin{aligned}
& \max_{x \in \{0,1\}^{n \times p}} && c_1^\top x_1 + c_2^\top x_2 + \dots + c_p^\top x_p \\
& \text{s.t.} && B_1 x_1 \leq b_1 \\
& && B_2 x_2 \leq b_2 \\
& && B_p x_p \leq b_p \\
& && D_1 x_1 + D_2 x_2 + \dots + D_p x_p \leq d
\end{aligned} \tag{11}$$

Let $Q_j = \{x_j \in \{0,1\}^n : B_j x \leq b_j\}$, the Dantzig-Wolfe relaxation is given by

$$\begin{aligned}
& \max_{\lambda} && \sum_j \sum_{v \in Q_j} \lambda_v^j c_j^\top v \\
& \text{s.t.} && \sum_j \sum_{v \in Q_j} \lambda_v^j D_j v \leq d \\
& && \sum_{v \in Q_j} \lambda_v^j = 0 \quad \forall j \\
& && \lambda \in \mathbb{R}_+^{n \times p}
\end{aligned} \tag{12}$$

II.3. Examples

Generalized assignment – DW

Exercise 13

Write the Dantzig-Wolfe reformulation of Equation 1.

Cutting stock – DW

Exercise 14

Write the Dantzig-Wolfe reformulation of Equation 2.

Facility location – DW

Exercise 15

Write the Dantzig-Wolfe reformulation of Equation 3.

Network design – DW

Exercise 16

Write the Dantzig-Wolfe reformulation of Equation 4.

III. Column generation

Exponential size

Remark 17

The Dantzig-Wolfe relaxation in Equation 9 has an exponential number of variables (size of the Minkowski-Weyl representation of $\text{conv}(Q)$). It cannot be solved “as is” by existing solvers.

We describe an approach to tackle this exponential size by considering a few generators only.

III.1. Main problem and pricing

Restricted Dantzig-Wolfe relaxation

Definition 18

The restricted Dantzig-Wolfe relaxation is defined by using a subset of vertices $K' \subset K$ and a subset of rays $H' \subset H$ in Equation 9:

$$\begin{aligned}
 \max_{\lambda, \mu} \quad & \sum_{k \in K'} \lambda_k c^\top v_k + \sum_{h \in H'} \mu_h c^\top r_h \\
 \text{s.t.} \quad & \sum_{k \in K'} \lambda_k A_1 v_k + \sum_{h \in H'} \mu_h A_1 r_h \leq b_1 \\
 & \sum_{k \in K'} \lambda_k = 1 \\
 & \lambda \in \mathbb{R}_+^{K'}, \mu \in \mathbb{R}_+^{H'},
 \end{aligned} \tag{13}$$

If it is unbounded, then Equation 9 is also unbounded. Otherwise, let $(\bar{\lambda}, \bar{\mu})$ be an optimal solution of Equation 13.

If $(\bar{\lambda}, \bar{\mu})$ is not optimal for Equation 9, the question is how to select new generator candidates to insert into K' and H' .

Dual of main problem

Proposition 19

The dual of Equation 9 is given by

$$\begin{aligned}
 \min_{\sigma, \pi} \quad & \sigma + \pi^\top b_1 \\
 \text{s.t.} \quad & (c - A_1^\top \pi)^\top v_k - \sigma \leq 0 \quad \forall k \in K \\
 & (c - A_1^\top \pi)^\top r_h \leq 0 \quad \forall h \in H \\
 & \pi \geq 0
 \end{aligned} \tag{14}$$

In the unrestricted Dantzig-Wolfe relaxation, let $\sigma \in \mathbb{R}$ be the Lagrange multiplier associated with constraint $\sum_{k \in K} \lambda_k = 1$, and $\pi \geq 0$ the vector of Lagrange multipliers associated with $\sum_{k \in K} \lambda_k A_1 v_k + \sum_{h \in H} \mu_h A_1 r_h \leq b_1$. The Lagrangian writes

$$\begin{aligned}
\mathcal{L}(\lambda, \mu; \sigma, \pi) &= \sum_k \lambda_k c^\top v_k + \sum_h \mu_h c^\top r_h \\
&\quad + \sigma \left(1 - \sum_k \lambda_k \right) \\
&\quad + \pi^\top \left(b_1 - \sum_k \lambda_k A_1 v_k - \sum_h \mu_h A_1 r_h \right) \\
\mathcal{L}(\lambda, \mu; \sigma, \pi) &= \sum_k \lambda_k (c^\top v_k - \pi^\top A_1 v_k - \sigma) \\
&\quad + \sum_h \mu_h (c^\top r_h - \pi^\top A_1 r_h) \\
&\quad + \sigma + \pi^\top b_1
\end{aligned} \tag{15}$$

And thus the dual is $\min_{\sigma, \pi} \max_{\lambda, \mu} \mathcal{L}(\lambda, \mu; \sigma, \pi)$, which boils down to

$$\begin{aligned}
\min_{\sigma, \pi} \quad & \sigma + \pi^\top b_1 \\
\text{s.t.} \quad & c^\top v_k - \pi^\top A_1 v_k - \sigma \leq 0 \quad \forall k \\
& c^\top r_h - \pi^\top A_1 r_h \leq 0 \quad \forall h \\
& \pi \geq 0
\end{aligned} \tag{16}$$

Dual variables are shadow prices

Idea 20

Let $(\bar{\sigma}, \bar{\pi})$ be a dual optimal solution associated with $(\bar{\lambda}, \bar{\mu})$ for Equation 13. The dual variables tell us whether it is worth adding new generators to K' and H' .

Finding new generators

Proposition 21

For $k \in K$, let $\bar{c}_k = (c - A_1^\top \bar{\pi})^\top v_k - \bar{\sigma}$ be the reduced cost associated with vertex v_k . For $h \in H$, let $\bar{c}_h = (c - A_1^\top \bar{\pi})^\top r_h$ be the reduced cost associated with ray r_h . If all $\bar{c}_k$ are zero and all $\bar{c}_h$ are zero, then $(\bar{\lambda}, \bar{\mu})$ is an optimal solution to the full relaxation Equation 13. Otherwise, generators with positive reduced costs can be added to the main problem to improve the solution.

Indeed, if $((\bar{\lambda}, \bar{\mu}), (\bar{\sigma}, \bar{\pi}))$ is a primal-dual optimal pair for Equation 13, then they satisfy complementary slackness. Furthermore, $(\bar{\lambda}, \bar{\mu})$ is feasible for the full problem Equation 9. Thus, $(\bar{\lambda}, \bar{\mu})$ is optimal for Equation 9 as well if and only if $(\bar{\sigma}, \bar{\pi})$ is feasible for its dual Equation 14. If it is not feasible, there must be a violated constraint, and we can add it to the dual, which means adding the corresponding variable to the primal.

Pricing subproblem

Definition 22

The pricing subproblem is the optimization problem

$$\zeta = -\bar{\sigma} + \max_{x \in Q} (c - A_1^\top \bar{\pi})^\top x \quad (17)$$

There can be three situations:

- The problem Equation 17 is unbounded if and only if there is an extreme ray r_h such that $(c - A_1^\top \bar{\pi})^\top r_h > 0$, i.e. with positive reduced cost.
- If it is bounded and $\eta > 0$, then there is a vertex v_k with positive reduced cost.
- If it is bounded and $\eta \leq 0$, no variable has a positive reduced cost.

III.2. Column generation

Column generation for large MILPs

Algorithm 23

1. Start with variable subsets K' and H' .
2. Solve the restricted main problem (Equation 13), obtain $(\bar{\lambda}, \bar{\mu})$ and $(\bar{\sigma}, \bar{\pi})$.
3. Solve the pricing subproblem (Equation 17), and depending on its result, either terminate or add variables to K' / H' and loop.

Bounding the optimal objective

Proposition 24

Identical subproblems

Remark 25

III.3. Examples

Generalized assignment – pricing

Exercise 26

Identify the pricing subproblem of Equation 12. How would you solve it?

Cutting stock – pricing

Exercise 27

Identify the pricing subproblem of Equation 12. How would you solve it?

Facility location – pricing

Exercise 28

Identify the pricing subproblem of Equation 12. How would you solve it?

Network design – pricing

Exercise 29

Identify the pricing subproblem of Equation 12. How would you solve it?

IV. Branch and price

Branch & price

Algorithm 30

The branch & price algorithm is a branch & bound where at each node, the relaxation is given by Dantzig-Wolfe.

IV.1. Branching techniques

Adding branching inequalities

Remark 31

It is better to add branching inequalities to the main problem than the pricing problem.

Otherwise one may disturb the specific structure that makes pricing easy.

Variable to branch on

Remark 32

It is better to branch on the original variables x as opposed to the reformulation variables λ .

Two reasons:

- Fixing $\lambda_k = 1$ means separating just one vertex from the rest of the polytope, not a very useful split.
- Fixing $\lambda_k = 0$ doesn't mean we won't try to include v_k further down in the pricing step, which means we'd have to look for the second-best solution of pricing.

IV.2. Implementation tricks

Suboptimal pricing

Remark 33

Still works as long as the columns have positive reduced cost.

Several columns at once

Remark 34

Bibliography

- [1] M. Conforti, G. Cornuéjols, and G. Zambelli, *Integer Programming*, 1st ed. 2014., no. 271. in Graduate Texts in Mathematics. Cham: Springer International Publishing : Imprint: Springer, 2014. doi: 10.1007/978-3-319-11008-0.
- [2] E. Uchoa, A. Pessoa, and L. Moreno, "Optimizing with Column Generation: Advanced Branch-Cut-and-Price Algorithms (Part I)," technical report L-2024-3, Aug. 2024. [Online]. Available: <https://optimizingwithcolumngeneration.github.io/>