

Assignment-based Subjective Questions

1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?

Ans.1. Season: Fall has the highest demand for rental bikes followed by summer. Spring is the season where people least rent bikes.


2. yr: The demand for rental bikes has significantly increased in 2019.

3. mnth: First the demand for bikes kept increasing month after month. Sept has the most count for renting bikes. After that there is a decrease in demand.

4. holiday: When there is a holiday, the demand has increased.

5. weekday: Sunday has more demand for rental bikes as compared to other even though the weekday is not giving a clear picture for demand.

6. weathersit: The clear weather has highest demand.




2. Why is it important to use `drop_first=True` during dummy variable creation?

Ans. The `drop_first = True` is used while creating dummy variables to drop the base/reference category. The reason for this is to avoid the multi-collinearity getting added into the model if all dummy variables are included. The reference category can be easily deduced where 0 is present in a single row for all the other dummy variables of a particular category.

3. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?

Ans.




“temp” and “atemp” are the variables which have the highest correlation with target variable(cnt).

4. How did you validate the assumptions of Linear Regression after building the model on the training set?


Ans.

1. **Linear relationship between independent and dependent variables** – The linearity is validated by looking at the points distributed symmetrically around the diagonal line of the actual vs predicted plot as shown in the below figure.




2. **Error terms are independent of each other** – We can see there is no specific Pattern observed in the Error Terms with respect to Prediction, hence we can say Error terms are independent of each other.

3. **Error terms are normally distributed:** Histogram and distribution plot helps to understand the normal distribution of error terms along with the mean of 0. The figure below clearly depicts the same.



4. Error terms have constant variance (homoscedasticity):

We can see Error Terms have approximately a Constant Variance, hence it follows the Assumption of Homoscedasticity.



5. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?

Ans. The top 3 variables are:

i. **Temp :**

Temperature is the Most Significant Feature which affects the Business positively, Whereas the other Environmental condition such as Raining, Humidity, Windspeed and Cloudy affects the Business negatively.

ii. **'Yr':**

The growth year on year seems organic from 2018 to 2019.

iii. **'Season':**

Fall has the highest demand for rental bikes followed by summer.

General Subjective Questions

1. Explain the linear regression algorithm in detail.

Ans. i. Linear regression is the method of finding the best linear relationship within the independent variables and dependent variables. The algorithm uses the best fitting line to map the association between independent variables with dependent variable.

ii. There are 2 types of linear regression algorithms

- Simple Linear Regression – Single independent variable is used.
 - $Y = \beta_0 + \beta_1 X$ is the line equation used for SLR.
- Multiple Linear Regression – Multiple independent variables are used.
 - $Y = \beta_0 + \beta_1 X_1 + \dots + \beta_p X_p + \epsilon$ is the line equation for MLR.
- $\beta_0 = \text{value of the } Y \text{ when } X = 0 \text{ (Y intercept)}$
- $\beta_1, \beta_2, \dots, \beta_p = \text{Slope or the gradient.}$

iii. Cost functions – The cost functions help to identify the best possible values for the $\beta_0, \beta_1, \beta_2, \dots, \beta_p$ which helps to predict the probability of the target variable. The minimization approach is used to reduce the cost functions to get the best fitting line to predict the dependent variable. There are 2 types of cost function minimization approaches – **Unconstrained and constrained.**

- Sum of squared function is used as a cost function to identify the best fit line. The cost functions are usually represented as
 - The straight-line equation is $Y = \beta_0 + \beta_1 X$
 - The prediction line equation would be $Y_{pred} = \beta_0 + \beta_1 x_i$ and the actual Y is as Y_i .
 - Now the cost function will be $J(\beta_1, \beta_0) = \sum (y_i - \beta_1 x_i - \beta_0)^2$
- The unconstrained minimization are solved using 2 methods
 - Closed form
 - Gradient descent
- a. While finding the best fit line we encounter that there are errors while mapping the actual values to the line. These errors are nothing but the residuals. To minimize the error squares OLS (Ordinary least square) is used.
 - i. $e_i = y_i - y_{pred}$ provides the error for each of the data point.
 - ii. OLS is used to minimize the total e^2 which is called as Residual sum of squares.
 - iii. $RSS = \sum_{i=1}^n (y_i - y_{pred})^2$
- b. Ordinary Least Squares method is used to minimize Residual Sum of Squares and estimate beta coefficients.

2. Explain the Anscombe's quartet in detail.

Ans. Statistics like variance and standard deviation are usually considered good enough parameters to understand the variation of some data without actually looking at every data point. The statistics are great for describing the general trends and aspects of the data. Francis Anscombe realized in 1973 that only statistical measures are not good enough to depict the data sets. He created several data sets all with several identical statistical properties to illustrate the fact.

- Illustrations


- One of the data set is as follows:

	x1	x2	x3	x4	y1	y2	y3	y4
0	10	10	10	8	8.040000	9.140000	7.460000	6.580000
1	8	8	8	8	6.950000	8.140000	6.770000	5.760000
2	13	13	13	8	7.580000	8.740000	12.740000	7.710000
3	9	9	9	8	8.810000	8.770000	7.110000	8.840000
4	11	11	11	8	8.330000	9.260000	7.810000	8.470000
5	14	14	14	8	9.960000	8.100000	8.840000	7.040000
6	6	6	6	8	7.240000	6.130000	6.080000	5.250000
7	4	4	4	19	4.260000	3.100000	5.390000	12.500000
8	12	12	12	8	10.840000	9.130000	8.150000	5.560000
9	7	7	7	8	4.820000	7.260000	6.420000	7.910000
10	5	5	5	8	5.680000	4.740000	5.730000	6.890000

- If the descriptive statistics are checked for above data set then all looks same:

	x1	x2	x3	x4	y1	y2	y3	y4
count	11.000000	11.000000	11.000000	11.000000	11.000000	11.000000	11.000000	11.000000
mean	9.000000	9.000000	9.000000	9.000000	7.500909	7.500909	7.500000	7.500909
std	3.316625	3.316625	3.316625	3.316625	2.031568	2.031657	2.030424	2.030579
min	4.000000	4.000000	4.000000	8.000000	4.260000	3.100000	5.390000	5.250000
25%	6.500000	6.500000	6.500000	8.000000	6.315000	6.695000	6.250000	6.170000
50%	9.000000	9.000000	9.000000	8.000000	7.580000	8.140000	7.110000	7.040000
75%	11.500000	11.500000	11.500000	8.000000	8.570000	8.950000	7.980000	8.190000
max	14.000000	14.000000	14.000000	19.000000	10.840000	9.260000	12.740000	12.500000

- However, when plotted these points, the relation looks completely different as depicted below.



- Anscombe's Quartet signifies that multiple data sets with many similar statistical properties could still be different from one another when plotted.
- The dangers of outliers in data sets are warned by the quartet. Check the bottom 2 graphs. If those outliers would have not been there the descriptive stats would have been completely different in that case.

- Important points
 - Plotting the data is very important and a good practice before analysing the data.
 - Outliers should be removed while analysing the data.
 - Descriptive statistics do not fully depict the data set in its entirety.

3. What is Pearson's R?

Ans. The Pearson's R (also known as Pearson's correlation coefficients) measures the strength between the different variables and the relation with each other. The Pearson's R returns values between -1 and 1. The interpretation of the coefficients are:

- *-1 coefficient indicates strong inversely proportional relationship.*
- *0 coefficient indicates no relationship.*
- *1 coefficient indicates strong proportional relationship.*

$$r = \frac{n(\sum x * y) - (\sum x) * (\sum y)}{\sqrt{[n\sum x^2 - (\sum x)^2] * [n\sum y^2 - (\sum y)^2]}}$$

Where:

N = the number of pairs of scores

$\sum xy$ = the sum of the products of paired scores

$\sum x$ = the sum of x scores

$\sum y$ = the sum of y scores

$\sum x^2$ = the sum of squared x scores

$\sum y^2$ = the sum of squared y scores

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

Ans.

- Definition: The scaling is the data preparation step for regression model. The scaling normalizes these varied datatypes to a particular data range.
- Most of the times the feature data is collected at public domains where the interpretation of variables and units of those variables are kept open collect as much as possible. This results in to the high variance in units and ranges of data. If scaling is not done on these data sets, then the chances of processing the data without the appropriate unit conversion are high. Also the higher the range then higher the possibility that the coefficients are impaired to compare the dependent variable variance.
The scaling only affects the coefficients. The prediction and precision of prediction stays unaffected after scaling.
- Normalization/Min-Max scaling – The Min max scaling normalizes the data within the range of 0 and 1. The Min max scaling helps to normalize the outliers as well.

$$\text{MinMaxScaling: } x = \frac{x - \min(x)}{\max(x) - \min(x)}$$

- Standardization converges all the data points into a standard normal distribution where mean is 0 and standard deviation is 1.

$$\text{Standardization: } x = \frac{x - \text{mean}(x)}{\text{sd}(x)}$$

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

$$VIF = \frac{1}{1 - R^2}$$

Ans. The VIF formula clearly signifies when the VIF will be infinite. If the R^2 is 1 then the VIF is infinite. The reason for R^2 to be 1 is that there is a perfect correlation between 2 independent variables.

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression.

Ans. Q-Q plots are the quantile-quantile plots. It is a graphical tool to assess the 2 data sets are from common distribution. The theoretical distributions could be of type normal, exponential or uniform. The Q-Q plots are useful in the linear regression to identify the train data set and test data set are from the populations with same distributions. This is another method to check the normal distribution of the data sets in a straight line with patterns explained below

- Interpretations
 - Similar distribution: If all the data points of quantile are lying around the straight line at an angle of 45 degree from x-axis.
 - Y values < X values: If y-values quantiles are lower than x-values quantiles.
 - X values < Y values: If x-values quantiles are lower than y-values quantiles.
 - Different distributions – If all the data points are lying away from the straight line.
- Advantages
 - Distribution aspects like loc, scale shifts, symmetry changes and the outliers all can be identified from the single plot.
 - The plot has a provision to mention the sample size as well.