

Vasculogenesis in Hemangioblastomas associated with von Hippel-Lindau Disease

Sven Gläsker^{1,2}, Jonathan Smith^{2,3}, Mark Raffeld⁴, Russell R. Lonser², Alexander O. Vortmeyer^{5,2}

¹Department of Neurosurgery, Freiburg University Medical Center, Germany
²Surgical Neurology Branch, NINDS, National Institutes of Health, Bethesda, MD, USA
³Department of Neurology, Mayo Clinic, Rochester, MN, USA
⁴Laboratory of Pathology, National Cancer Institute, Bethesda, MD, USA
⁵Department of Pathology, Yale University School of Medicine, New Haven, CT, USA

Albert-Ludwigs-Universität Freiburg


UNI FREIBURG

Introduction

Arvid Lindau *Proceedings of the Royal Society of Medicine* (1931)
 hemangioblastoma tissue may be derived from a "congenital anlage"

The histological picture reveals an "... embryological type of the tumor cells"

Stein et al. *J Neurosurg* (1960)
 angiomeseenchymal origin of hemangioblastoma based on developmental biologic observations




early blood island formation from angiogenic mesenchyme incipient vasoformation

Albert-Ludwigs-Universität Freiburg


UNI FREIBURG

Introduction

Choi et al. *Development* (1998)
 Identification of the hemangioblast:
 A common precursor for hematopoietic and endothelial cells.



Huber et al. *Nature* (2004)
 Hemangioblast is a subpopulation of mesoderm that co-express SCL, Brachyury and FLK-1



Albert-Ludwigs-Universität Freiburg

UNI FREIBURG

Methods

Material

- 60 VHL disease-associated hemangioblastomas
- Investigate vascular differentiation

Methods


- Histomorphology
- Immunohistochemistry (CD31, NSE)
- Microdissection-based genetic analysis
- HIF

Albert-Ludwigs-Universität Freiburg

UNI FREIBURG

Results

- In 47/60 tumors no evidence of vasculogenesis or hematopoiesis
- Tumor cells scattered between reactive angiogenetic vessels or arranged in solid clusters




CD31

Albert-Ludwigs-Universität Freiburg

UNI FREIBURG







Results

- 13/60 tumors epithelial vasoformation
- lined by cuboidal or flat cells
- frequently contained red blood cell precursors or mature red blood cells



Albert-Ludwigs-Universität Freiburg

UNI FREIBURG



Conclusion

Albert-Ludwigs-Universität Freiburg

UNI
FREIBURG

- Hemangioblastomas show two types of vascularization:
 1. intense mature capillary network as result of VEGF-driven reactive angiogenesis
 2. vasculogenesis by the neoplastic "stromal" cells
- Both processes next to each other without transition or connection
- Hemangioblastomas resemble the process of vasculogenesis accompanied by extensive angiogenesis

A

B

C

Legend:
● = stromal cell
○ = endothelial cell
○ = capillary
● = pericyte
○ = angioblastic endothelium