


Pomegranates, Polyphenols, and Kidney Cancer




Allan J. Pantuck, MD, MS, FACS
Associate Professor of Urology
David Geffen School of Medicine at UCLA
June 25, 2009

Agents with Cancer Chemopreventive Potential

Anti-androgens [SAR-12, AR] (Enasteride, Dutasteride, DIM, SARMs)	Anti-estrogens [ER-alpha/beta, Aromatase, TGFalpha] (Toremifene, Raloxifene, Arimidex, I3C/DIM)	Anti-angiogenesis Modulators (Thalidomide analogs, Avastin, Celecoxib)
Antioxidant Modulators [ROS, GSH/GST-P1, iNOS] (Selenium, Vitamin E, Lycopene, Polyphenols, Resveratrol, Sulforaphanes)	Differentiation Agents [VDR, RXR, HDAC, DNMT] (Vit D Analogs, Targretin, SAHA, valproic acid)	PPAR (gamma/delta) Modulators (Glitazones, DHEA Analogs, Retinoids, NSAIDs, 15d-PGJ2)
Arachidonic Acid Modulators [NFKB, COX (PGE2), LOX (HETE), PPARs] (ASA/NSAIDs, Celecoxib, Rofecoxib, R-flurbiprofen, NO-ASA, Vit E, Zileuton)	Anti-proliferation/Cell Cycle Agents [DDC, RXR/RAR, p21, p27] (DMFO, Panretin, Vitamin D)	IGF-1/IGFBP-3 Modulators (Soy, Retinoids, Tamoxifen, Lycopene, Vit D)
Phytoestrogens [ER-beta, SAR, AR, RTK] (Genistein, Daidzein/Equol, Resveratrol)	Signal Transduction Modulators (Kinases) [EGFR (Tarceva), PDGFR (STI-571), VEGFR/POGFR (SU-11248), PI3K/AKT/mTOR (Celecoxib, Rapamycin), NFkB (polyphenols)]	Novel Growth Factor Modulators [ET-1, IL-6, NGF] (Atrasentan, Celecoxib, Bowman-Birk Inhibitor)
	Ras/Farnesyl/Geranyl Transferase Modulators (Statins, SCH 66336, L-778,123, Perillyl Alcohol)	Telomerase Modulators Antiandrogens, SERMs, Polyphenols, Vit D, NGF
		Gene-based vaccines (PS3, Rb, E1A/PSA, PSMA, PSA/B7/GMCSF)

Phenolics in Pomegranate Juice



*Phenolics include **phenols**, which have one phenol unit in their structure, and **polyphenols**, which have >1 phenol unit in their structure. For simplicity, we call all the compounds Polyphenols.

Pomegranates and Prostate Cancer Pre-Clinical Data Summary: 2001

In vitro studies show:

- 59-75% growth inhibition of PC3
- Delayed progression into S phase
- Induction of apoptosis

In vivo studies in SCID mice show:

- 52% growth inhibition of LAPC-9 tumors
- 70% reduction in PSA
- Prolonged survival


Phase II Study for Men with Rising PSA After RRP or XRT: 2002

- Men with Recurrent Prostate Cancer
- Rising PSA after RRP or XRT
- Low risk: PSA < 5, Gleason < 8
- No evidence of metastatic disease
- No previous hormone therapy
- Baseline PSA DT

Pantuck et al
Clinical Cancer Research, 2006

Treatment

8 oz. Pomegranate juice daily
providing 1.5 mmol polyphenols



Updated Results: PSA Doubling Time Year 5

	PSADT Before Baseline	PSADT After Baseline	Change
Mean \pm SD	15.0 months	58 months	43.0 months
		Non-Active 51 months Active 69 months	P=0.0001


ASCO, 2008

2009 Pomegranate Research Portfolio

- POM Phase II Continuation Study
- POM Phase III Multi-Center Study
- Neoadjuvant POM Pill Study
- POM and NF κ B Pathway
- POM and IGF Axis
- POM and Kidney Cancer


2009 Pomegranate Research Portfolio

- POM Phase II Continuation Study
- POM Phase III Multi-Center Study
- Neoadjuvant POM Pill Study
- POM and NF κ B Pathway
- POM and IGF Axis
- POM and Kidney Cancer




POMX inhibits constitutive and TNF α -induced NF- κ B activity

POMX inhibits constitutive and TNF α -induced NF- κ B CL1




Cell growth inhibition is attributable to stimulation of apoptosis (Annexin V)



Dilution	Apoptosis (%)
Veh	11.1
1:4000	20.6
1:2000	31.4
1:1000	60.7
1:500	67.4

Pom is Inhibiting NFkB at the activation of the IKK kinase



Something New for 2009

Pom and Kidney Cancer


Kidney Cancer: A Family Of Tumors

RCC


Type	Clear cell	Papillary type 1	Papillary type 2	Chromophobe	Oncocytoma
Incidence (%)	75%	5%	10%	5%	5%
Associated mutations	VHL	c-Met	FH	BHD	BHD

BHD=Birt-Hogg-Dubé; FH=fumarate hydratase; VHL= von Hippel-Lindau.
Modified from Linehan WM et al. J Urol. 2003;170:2163-2172.

Kidney Cancer Molecular Pathways




VHL, HIF and NFkB




Pom Inhibits:




Both VHL+ and VHL- RCC Cells Sensitive to Killing Effects of POM-X




Isogenic VHL + and VHL - cell lines




High NF-κB Activity in RCC cells lacking VHL




VHL, NFkB and EMT




VHL Status and Epithelial and Mesenchymal Phenotype of Kidney Cancer Cell Lines



Inhibiting NFkB with a repressor in kidney cancer cells decreases N-Cadherin levels




....So does chronic low dose administration of Pom-X



....and it increases levels of E-Cadherin



POM inhibits invasion of kidney cancer cells



Plans for 2009-2010

Complete current clinical trials

Further evaluation of POM
for kidney cancer

?Clinical trials in RCC?

Acknowledgements

UCLA Urology
William Aronson, MD
Arie Beldegrun, MD
Allan Pantuck, MD
Nazy Zomorodian, MS

Technion Institute
Michael Aviram, PhD



Sense Foundation

UCLA Medicine
Pinchas Cohen, MD
David Heber, MD
Louis Ignarro, PhD
Harley Liker, MD
Matthew Rettig, MD
Navindra Seeram, PhD

UCLA Biomathematics
Robert Elashoff, PhD
Jeffrey Gornbein, PhD
Hejing Wang, MD