

Use of Molecular Pathology to Predict Outcomes in RCC

Federico A. Monzon M.D.

CGL
Genetic Diagnostic Laboratory

Variability in Clinical Behavior

- As many as 20-30% of surgically treated patients (T1/T2) with clear cell renal cell carcinoma (ccRCC) develop local or distant recurrence, and nearly 50% will eventually develop stage IV (metastatic) disease.
 - Can we predict which tumors are going to recur?
- One of the key dilemmas in the management of metastatic ccRCC is the manifestation of innate or acquired resistance to anti-angiogenic therapy.
 - Can we predict which tumors are going to respond?

CGL
Genetic Diagnostic Laboratory

Variability in Clinical Behavior

- As many as 20-30% of surgically treated patients (T1/T2) with clear cell renal cell carcinoma (ccRCC) develop local or distant recurrence, and nearly 50% will eventually develop stage IV (metastatic) disease.
 - Can we predict which tumors are going to recur?
- One of the key dilemmas in the management of metastatic ccRCC is the manifestation of innate or acquired resistance to anti-angiogenic therapy.
 - Can we predict which tumors are going to respond?

CGL
Genetic Diagnostic Laboratory

Summary Profiles of Renal Tumors

Clear Cell RCC

Papillary RCC

Chromophobe RCC

Oncocytoma

CGL
Genetic Diagnostic Laboratory

Molecular Subtypes of ccRCC

Genetic Clustering of Clear Cell Renal Cell Carcinoma based on array-CGH: Its Association with DNA Methylation Alteration and Patient Outcome. Clin Cancer Res 2008;14(17): Sep 1, 2008

Brannon, et al. Molecular Stratification of Clear Cell Renal Cell Carcinoma by Consensus Clustering Reveals Distinct Subtypes and Survival Patterns. Genes Cancer. Genes Cancer. 2010 February 1(2): 151-163.

CGL
Genetic Diagnostic Laboratory

Recurrence Prognosis – Additional Evidence

Other studies have confirmed the role of chromosomal imbalances as prognostic markers in clear cell and papillary RCC.

- ccRCC, 9p loss – bad prognostic factor (Brunelli M, et al. Pathol. 2008, 21(1):1-6).
- ccRCC, 9p loss – adverse predictor (Klatte T, et al. J Clin Oncol. 2009, 27(5):746-53).
- pRCC, 1q gain – adverse predictor (Szponar A, et al. Int J Cancer. 2009, 124(9):2071-6).
- ccRCC, Genomic profiles predict recurrence (Arai E, et al. Clin Cancer Res 2008;14(17) Sep 1, 2008)

A

Recurrence-free survival rate (%)







Days after resection

B

Overall survival rate (%)

Days after resection

CGL
Genetic Diagnostic Laboratory




Studies on Response to Therapy

- MDACC Sorafenib Study**
 - Sorafenib alone vs. sorafenib-IFN (post nephrectomy)
 - Tumor response, progression-free survival (PFS) and overall survival (OS) were analyzed with regard to marker levels in 26 tumor samples.
- MDACC Bevacizumab Study**
 - 50 patient study pretreated newly diagnosed mRCC patients with four cycles of bevacizumab 10mg/kg IV every two weeks prior to nephrectomy, and continued therapy in patients who demonstrated stable disease or response.


Sorafenib Cohort

- Multivariate Cox Proportional Hazards Regression for Progression Free Survival (PFS)
 - Longer PFS
 - 5q gain - HR = 0.25, 95% CI 0.08 to 0.81, P = 0.021
 - Shorter PFS
 - 1q gain - HR = 0.067 CI 0.006-0.76, P = 0.0292 (univariate model for sorafenib alone arm)
 - 9p loss - HR = 0.29, 95% CI 0.08 to 1.09, P = 0.0673




Sorafenib Cohort

- Multivariate model to predict time to progression
 - 14q loss - HR = 20.478, CI 1.300- 322.601, P = 0.0318 (Interaction with treatment arm);
 - 14q loss - HR = 5.146, CI 0.971 - 27.267, P = 0.0541 (Sorafenib only arm)
 - 18p loss - HR (for normal 18p) = 0.045, CI 0.002- 0.826, P = 0.0368




Bevacizumab Cohort

- Significant association with response, Fisher's exact test
 - 14q loss, CRPR/SDPD, P = 0.0473
 - 5p/q gain, CRPRSD / PD, P = 0.02
 - 8q gain, CRPRSD / PD, P = 0.04




Bevacizumab Cohort

- Significant association with overall survival (OS), log rank
 - 5q gain, HR = 0.59, 95% CI (0.36 , 0.96), P = 0.002
- Significant association with recurrence free survival (RFS), multivariate Cox
 - 5q gain, HR = 0.06, 95% CI (0.01, 0.31), P = 0.00083
 - 15q loss, HR = 0.23, 95% CI (0.07, 0.82), P = 0.024



Gene Expression and Chromosomal Profiles

- Patient cohort treated with Bevacizumab (neoadjuvant)
- Tumors with cca gene expression profile show a good prognosis chromosomal profile
 - 5q gain
 - Normal 8q, 9p and 14q



What's next

- Confirm association of chromosomal loss with therapeutic response to antiangiogenic agents.
 - Bevacizumab, Erlotinib, Sunitinib
- Insights into biology and/or marker discovery
 - Loss of chromosomal regions associated with outcome could reflect:
 - Loss of additional tumor suppressor genes
 - Selection of mutated (constitutively active) oncogenes
 - Study correlations of VHL-HIF-VEGF proteins with outcome/genomic patterns
 - Search for TSGs in areas of association or genes that interact with treatment pathways
- Role of endothelial cells
 - Phenotype and Genotype

BCM
Baylor College of Medicine

CGL

Conclusions

- SNP arrays are a powerful tool for whole-genome analysis of chromosomal lesions (virtual karyotyping) in renal tumors.
- Chromosomal imbalances are associated with recurrence and outcome measures and could be used as prognostic markers
 - 5q gain is associated with better PFS in both Sorafenib and Bevacizumab cohorts and with better response in Bevacizumab treated patients .
 - 14q loss is associated with recurrence, shorter time to progression in Sorafenib cohort and associated with worse response in Bevacizumab treated patients .
 - Unclear if effect is related to prognosis (aggressive tumors) or therapeutic response prediction (tumors less sensitive to therapy), or both.
- Need to identify candidate genes to establish molecular mechanisms for recurrence and/or resistance to therapy.

BCM
Baylor College of Medicine

CGL

Acknowledgements

Methodist The Methodist Hospital



Martin Powers MD, Karla Alvarez BSc, Benita Mathai, Hyun-Jung Kim MD



Eric Jonasch MD



Bob Amato DO

More Collaborators:

- Jill Hagenkord MD, Zoran Gatalica MD & Shera Kash, PhD – Creighton
- Pheroze Tamboli MD, Kanishka Sircar MD – MD Anderson Cancer Center
- Clinical Genomics Facility, UPMC – Maureen Lyons-Weiler MSc and Christin Sciulli BSc
- Julia Bridge MD - U of Nebraska,

Clinical Genomics






Tuan Phuong M.D., Alberto Ayala, M.D. Steven Sherr M.D., Ph.D. & Rae Jo M.D., M.S., Ph.D.

BCM
Baylor College of Medicine

CGL

Questions?

BCM
Baylor College of Medicine

CGL