

よくある訴えと慢性疾患

内科大全外来編 文献

下書き 2026-10-07 23:03

2026年9月

文献

本文の右肩の番号 (1)2) …) は、章ごとの文献番号である。章の番号と題は本文と同じにしてある。

第1部 外来診療の基本

1-1 外来診療の進め方

- 1) Singh H, Meyer AN, Thomas EJ. The frequency of diagnostic errors in outpatient care: estimations from three large observational studies involving US adult populations. *BMJ Qual Saf* 2014;23:727-31.
- 2) Marvel MK, Epstein RM, Flowers K, Beckman HB. Soliciting the patient's agenda: have we improved? *JAMA* 1999;281:283-7.
- 3) Singh H, Giardina TD, Meyer AN, Forjuoh SN, Reis MD, Thomas EJ. Types and origins of diagnostic errors in primary care settings. *JAMA Intern Med* 2013;173:418-25.
- 4) Graber ML, Franklin N, Gordon R. Diagnostic error in internal medicine. *Arch Intern Med* 2005;165:1493-9.
- 5) Kroenke K, Jackson JL. Outcome in general medical patients presenting with common symptoms: a prospective study with a 2-week and a 3-month follow-up. *Fam Pract* 1998;15:398-403.
- 6) Birtwhistle RV, Godwin MS, Delva MD, et al. Randomised equivalence trial comparing three month and six month follow up of patients with hypertension by family practitioners. *BMJ* 2004;328:204.
- 7) Pincavage AT, Ratner S, Prochaska ML, et al. Outcomes for resident-identified high-risk patients and resident perspectives of year-end continuity clinic handoffs. *J Gen Intern Med* 2012;27:1438-44.
- 8) Schillinger D, Piette J, Grumbach K, et al. Closing the loop: physician communication with diabetic patients who have low health literacy. *Arch Intern Med* 2003;163:83-90.
- 9) 国立国語研究所「病院の言葉」委員会. 「病院の言葉」を分かりやすくする提案. 2009.
- 10) 保険医療機関及び保険医療養担当規則（昭和32年厚生省令第15号）. 第16条、第16条の2.
- 11) Gandhi TK, Sittig DF, Franklin M, Sussman AJ, Fairchild DG, Bates DW. Communication breakdown in the outpatient referral process. *J Gen Intern Med* 2000;15:626-31.
- 12) 厚生労働省保険局医療課. 令和8年度診療報酬改定の概要 7. 外来医療の機能分化・強化等. 2026.
- 13) 医師法（昭和23年法律第201号）. 第19条、第20条.
- 14) 刑法（明治40年法律第45号）. 第160条.
- 15) 健康保険法（大正11年法律第70号）. 第99条.
- 16) 学校保健安全法施行規則（昭和33年文部省令第18号）. 第19条.
- 17) 厚生労働省. 令和7年度 急性呼吸器感染症（ARI）総合対策に関するQ&A（Q20、Q21）. 2026年10月4日閲覧.
- 18) 個人情報保護委員会, 厚生労働省. 医療・介護関係事業者における個人情報の適切な取扱いのためのガイドライン. 2017（2026年4月一部改正）. 同 Q&A（事例集）Q4-6、Q4-24、Q4-25.
- 19) 厚生労働省医政局長通知. 応招義務をはじめとした診療治療の求めに対する適切な対応の在り方等について（医政発1225第4号）. 令和元年12月25日.
- 20) 厚生労働省. オンライン診療の適切な実施に関する指針. 2018（2026年4月改訂）.

- 21) Heritage J, Robinson JD, Elliott MN, Beckett M, Wilkes M. Reducing patients' unmet concerns in primary care: the difference one word can make. J Gen Intern Med 2007;22:1429-33.
- 22) Mamede S, van Gog T, van den Berge K, et al. Effect of availability bias and reflective reasoning on diagnostic accuracy among internal medicine residents. JAMA 2010;304:1198-203.

1-2 外来処方の決まりと見直し

- 1) 厚生労働省. 内服薬処方せんの記載方法の在り方に関する検討会報告書. 2010.
- 2) 日本医療機能評価機構 医療事故情報収集等事業. 医療安全情報 No.167 抗リウマチ剤（メトトレキサート）の過剰投与に伴う骨髄抑制（第3報）. 2020.
- 3) 厚生労働省保険局医療課. 令和8年度診療報酬改定の概要（7. 外来医療の機能分化・強化等、調剤）. 2026.
- 4) 厚生労働省. 後発医薬品のある先発医薬品（長期収載品）の選定療養について. 2026年10月4日閲覧.
- 5) 保険医療機関及び保険医療養担当規則（昭和32年厚生省令第15号）. 第20条、第23条.
- 6) Tarn DM, Heritage J, Paterniti DA, Hays RD, Kravitz RL, Wenger NS. Physician communication when prescribing new medications. Arch Intern Med 2006;166:1855-62.
- 7) 療担規則及び薬担規則並びに療担基準に基づき厚生労働大臣が定める揭示事項等（平成18年厚生労働省告示第107号）. 第10.
- 8) 診療報酬の算定方法（平成20年厚生労働省告示第59号）別表第一 第2章第5部 投薬（通則、F100 処方料、F400 処方箋料）と、その留意事項通知（令和8年3月5日 保医発0305第6号）.
- 9) 厚生労働省 地方厚生局麻薬取締部. 麻薬・覚醒剤原料などを携帯して日本を出国する方へ（医療用向精神薬の携帯輸出入について）. 2026年10月7日閲覧.
- 10) 厚生労働省保険局医療課. 令和4年度調剤報酬改定の概要（調剤）. 2022.
- 11) 薬剤師法（昭和35年法律第146号）. 第24条.
- 12) 日本薬剤師会委託事業. 平成27年度全国薬局疑義照会調査報告書. 2016.
- 13) Tamblyn R, Egale T, Huang A, Winslade N, Doran P. The incidence and determinants of primary nonadherence with prescribed medication in primary care: a cohort study. Ann Intern Med 2014;160:441-50.
- 14) Claxton AJ, Cramer J, Pierce C. A systematic review of the associations between dose regimens and medication compliance. Clin Ther 2001;23:1296-310.
- 15) Gandhi TK, Weingart SN, Borus J, et al. Adverse drug events in ambulatory care. N Engl J Med 2003;348:1556-64.
- 16) Savage RD, Visentin JD, Bronskill SE, et al. Evaluation of a common prescribing cascade of calcium channel blockers and diuretics in older adults with hypertension. JAMA Intern Med 2020;180:643-51.
- 17) 厚生労働省. 高齢者の医薬品適正使用の指針（総論編）. 2018.
- 18) Kojima T, Akishita M, Nakamura T, et al. Polypharmacy as a risk for fall occurrence in geriatric outpatients. Geriatr Gerontol Int 2012;12:425-30.
- 19) Budnitz DS, Lovegrove MC, Shehab N, Richards CL. Emergency hospitalizations for adverse drug events in older Americans. N Engl J Med 2011;365:2002-12.
- 20) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認. アムロジン錠、ロキソニン錠、モーラステープ、エックスフォージ配合錠、リウマトレックスカプセル、メルカゾール錠、ユリノーム錠、ラミシール錠、リーマス錠）.
- 21) 厚生労働省. 医薬品副作用被害救済制度の概要と制度への協力をお願いについて. 医薬品・医療機器等安全性情報 No.337. 2016.
- 22) Bangalore S, Kamalakkannan G, Parkar S, Messerli FH. Fixed-dose combinations improve medication compliance: a meta-analysis. Am J Med 2007;120:713-9.

4 文献

- 23) Manzoli L, Flacco ME, Boccia S, et al. Generic versus brand-name drugs used in cardiovascular diseases. *Eur J Epidemiol* 2016;31:351-68.
- 24) Kesselheim AS, Misono AS, Lee JL, et al. Clinical equivalence of generic and brand-name drugs used in cardiovascular disease: a systematic review and meta-analysis. *JAMA* 2008;300:2514-26.
- 25) 厚生労働省医薬・生活衛生局医薬品審査管理課長通知. 後発医薬品の生物学的同等性試験ガイドライン等の一部改正について（薬生薬審発0319第1号）. 令和2年3月19日.
- 26) Kesselheim AS, Bykov K, Avorn J, Tong A, Doherty M, Choudhry NK. Burden of changes in pill appearance for patients receiving generic cardiovascular medications after myocardial infarction: cohort and nested case-control studies. *Ann Intern Med* 2014;161:96-103.

1-3 薬を出す前に確かめること

- 1) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）.
- 2) 日本産科婦人科学会, 日本産婦人科医会編. 産婦人科診療ガイドラインー産科編2023. 日本産科婦人科学会; 2023.
- 3) 国立成育医療研究センター 妊娠と薬情報センター. 妊娠と薬のご相談について／授乳中に安全に使用できると考えられる薬／授乳中の使用には適さないと考えられる薬（2025年8月改訂）. <https://www.ncchd.go.jp/kusuri/>（2026年10月に確認）
- 4) 厚生省児童家庭局母子保健課長・保健医療局地域保健・健康増進栄養課生活習慣病対策室長通知. 神経管閉鎖障害の発症リスク低減のための妊娠可能な年齢の女性等に対する葉酸の摂取に係る適切な情報提供の推進について（児母第72号, 健医地生発第78号）. 平成12年12月28日.
- 5) Shenoy ES, Macy E, Rowe T, Blumenthal KG. Evaluation and management of penicillin allergy: a review. *JAMA* 2019;321:188-99.
- 6) Picard M, Robitaille G, Karam F, et al. Cross-reactivity to cephalosporins and carbapenems in penicillin-allergic patients: two systematic reviews and meta-analyses. *J Allergy Clin Immunol Pract* 2019;7:2722-38.e5.
- 7) Gandhi S, Fleet JL, Bailey DG, et al. Calcium-channel blocker-clarithromycin drug interactions and acute kidney injury. *JAMA* 2013;310:2544-53.
- 8) 厚生労働省医薬食品局総務課長・安全対策課長通知. 医薬品服用中の自動車運転等の禁止等に関する患者への説明について（薬食総発0529第2号, 薬食安発0529第2号）. 平成25年5月29日.
- 9) 厚生労働省医薬・生活衛生局医薬安全対策課長通知. 抗コリン作用を有する薬剤における禁忌「緑内障」等に係る添付文書の「使用上の注意」改訂について（薬生安発0618第2号）. 令和元年6月18日.
- 10) 世界アンチ・ドーピング機構. 世界アンチ・ドーピング規程 2026年禁止表国際基準（日本アンチ・ドーピング機構による日本語版）. 2026.
- 11) Parekh TM, Raji M, Lin YL, Tan A, Kuo YF, Goodwin JS. Hypoglycemia after antimicrobial drug prescription for older patients using sulfonyleureas. *JAMA Intern Med* 2014;174:1605-12.
- 12) Ahlqvist VH, Sjöqvist H, Dalman C, et al. Acetaminophen use during pregnancy and children's risk of autism, ADHD, and intellectual disability. *JAMA* 2024;331:1205-14.
- 13) Okubo Y, Hayakawa I, Sugitate R, Nariai H. Maternal acetaminophen use and offspring's neurodevelopmental outcome: a nationwide birth cohort study. *Paediatr Perinat Epidemiol* 2026;40:70-9.
- 14) Sheikh J, Allotey J, Sobhy S, et al. Maternal paracetamol (acetaminophen) use during pregnancy and risk of autism spectrum disorder and attention deficit/hyperactivity disorder in offspring: umbrella review of systematic reviews. *BMJ* 2025;391:e088141.

1-4 検査の出し方と偶発所見

- 1) Pauker SG, Kassirer JP. The threshold approach to clinical decision making. *N Engl J Med* 1980;302:1109-17.

- 2) McGee S. Simplifying likelihood ratios. *J Gen Intern Med* 2002;17:646-9.
- 3) Leuchten N, Hoyer A, Brinks R, et al. Performance of antinuclear antibodies for classifying systemic lupus erythematosus: a systematic literature review and meta-regression of diagnostic data. *Arthritis Care Res (Hoboken)* 2018;70:428-38.
- 4) Tan EM, Feltkamp TE, Smolen JS, et al. Range of antinuclear antibodies in "healthy" individuals. *Arthritis Rheum* 1997;40:1601-11.
- 5) Kim JE, Lee KT, Lee JK, Paik SW, Rhee JC, Choi KW. Clinical usefulness of carbohydrate antigen 19-9 as a screening test for pancreatic cancer in an asymptomatic population. *J Gastroenterol Hepatol* 2004;19:182-6.
- 6) Ganguli I, Simpkin AL, Lupo C, et al. Cascades of care after incidental findings in a US national survey of physicians. *JAMA Netw Open* 2019;2:e1913325.
- 7) Lazo M, Selvin E, Clark JM. Brief communication: clinical implications of short-term variability in liver function test results. *Ann Intern Med* 2008;148:348-52.
- 8) Pieper-Bigelow C, Strocchi A, Levitt MD. Where does serum amylase come from and where does it go? *Gastroenterol Clin North Am* 1990;19:793-810.
- 9) Casalino LP, Dunham D, Chin MH, et al. Frequency of failure to inform patients of clinically significant outpatient test results. *Arch Intern Med* 2009;169:1123-9.
- 10) Callen JL, Westbrook JI, Georgiou A, Li J. Failure to follow-up test results for ambulatory patients: a systematic review. *J Gen Intern Med* 2012;27:1334-48.
- 11) 日本医療機能評価機構 医療事故情報収集等事業. 医療安全情報 No.138 画像診断報告書の確認不足 (第2報). 2018.
- 12) 日本医療機能評価機構 医療事故情報収集等事業. 医療安全情報 No.150 病理診断報告書の確認忘れ—上部消化管内視鏡検査—. 2019.
- 13) O'Sullivan JW, Muntinga T, Grigg S, Ioannidis JPA. Prevalence and outcomes of incidental imaging findings: umbrella review. *BMJ* 2018;361:k2387.
- 14) Ashizawa K, Maruyama Y, Kobayashi T, et al. Guidelines for the management of pulmonary nodules detected by low-dose CT lung cancer screening 6th edition: compiled by the Japanese Society of CT Screening. *Jpn J Radiol* 2025;43:333-46.
- 15) Fassnacht M, Tsagarakis S, Terzolo M, et al. European Society of Endocrinology clinical practice guidelines on the management of adrenal incidentalomas, in collaboration with the European Network for the Study of Adrenal Tumors. *Eur J Endocrinol* 2023;189:G1-42.
- 16) 日本循環器学会, 日本心臓血管外科学会, 日本胸部外科学会, 日本血管外科学会. 2020年改訂版 大動脈瘤・大動脈解離診療ガイドライン. 2020.
- 17) 日本消化器がん検診学会 超音波検診委員会, 日本超音波医学会, 日本人間ドック学会. 腹部超音波検診判定マニュアル改訂版 (2021年). 2021.
- 18) Foley KG, Lahaye MJ, Thoeni RF, et al. Management and follow-up of gallbladder polyps: updated joint guidelines between the ESGAR, EAES, EFISDS and ESGE. *Eur Radiol* 2022;32:3358-68.
- 19) Horeweg N, van Rosmalen J, Heuvelmans MA, et al. Lung cancer probability in patients with CT-detected pulmonary nodules: a prespecified analysis of data from the NELSON trial of low-dose CT screening. *Lancet Oncol* 2014;15:1332-41.
- 20) Ohtsuka T, Fernandez-Del Castillo C, Furukawa T, et al. International evidence-based Kyoto guidelines for the management of intraductal papillary mucinous neoplasm of the pancreas. *Pancreatol* 2024;24:255-70.
- 21) UCAS Japan Investigators, Morita A, Kirino T, et al. The natural course of unruptured cerebral aneurysms in a Japanese cohort. *N Engl J Med* 2012;366:2474-82.
- 22) Rolfe A, Burton C. Reassurance after diagnostic testing with a low pretest probability of serious disease: systematic review and meta-analysis. *JAMA Intern Med* 2013;173:407-16.

- 23) Nakamura H, Arakawa K, Itakura H, et al. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *Lancet* 2006;368:1155-63.
- 24) Akl EA, Oxman AD, Herrin J, et al. Using alternative statistical formats for presenting risks and risk reductions. *Cochrane Database Syst Rev* 2011;(3):CD006776.

1-5 手術を受ける患者の内科評価

- 1) 日本循環器学会, ほか. 2022年改訂版 非心臓手術における合併心疾患の評価と管理に関するガイドライン. 2022 (2023年9月更新) .
- 2) Lee TH, Marcantonio ER, Mangione CM, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation* 1999;100:1043-9.
- 3) Mills E, Eyawo O, Lockhart I, Kelly S, Wu P, Ebbert JO. Smoking cessation reduces postoperative complications: a systematic review and meta-analysis. *Am J Med* 2011;124:144-54.e8.
- 4) Myers K, Hajek P, Hinds C, McRobbie H. Stopping smoking shortly before surgery and postoperative complications: a systematic review and meta-analysis. *Arch Intern Med* 2011;171:983-9.
- 5) Smetana GW, Lawrence VA, Cornell JE. Preoperative pulmonary risk stratification for noncardiothoracic surgery: systematic review for the American College of Physicians. *Ann Intern Med* 2006;144:581-95.
- 6) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。メトグルコ錠、ジャディアンス錠、アマリール錠、トリキュラー錠、プレマリン錠、エビスタ錠、ビビアント錠、ミカルディス錠、メネシット配合錠、デパケン錠、イーケブラ錠) .
- 7) 日本糖尿病学会 SGLT2阻害薬の適正使用に関する委員会. 糖尿病治療におけるSGLT2阻害薬の適正使用に関するRecommendation. 2022年7月改訂.
- 8) Musallam KM, Tamim HM, Richards T, et al. Preoperative anaemia and postoperative outcomes in non-cardiac surgery: a retrospective cohort study. *Lancet* 2011;378:1396-407.
- 9) Mansour A, Watson W, Shayani V, Pickleman J. Abdominal operations in patients with cirrhosis: still a major surgical challenge. *Surgery* 1997;122:730-5.
- 10) Teh SH, Nagorney DM, Stevens SR, et al. Risk factors for mortality after surgery in patients with cirrhosis. *Gastroenterology* 2007;132:1261-9.
- 11) Jørgensen ME, Torp-Pedersen C, Gislason GH, et al. Time elapsed after ischemic stroke and risk of adverse cardiovascular events and mortality following elective noncardiac surgery. *JAMA* 2014;312:269-77.
- 12) Schein OD, Katz J, Bass EB, et al. The value of routine preoperative medical testing before cataract surgery. *N Engl J Med* 2000;342:168-75.
- 13) Keay L, Lindsley K, Tielsch J, Katz J, Schein O. Routine preoperative medical testing for cataract surgery. *Cochrane Database Syst Rev* 2019;1:CD007293.
- 14) Chung F, Yuan H, Yin L, Vairavanathan S, Wong DT. Elimination of preoperative testing in ambulatory surgery. *Anesth Analg* 2009;108:467-75.
- 15) McFalls EO, Ward HB, Moritz TE, et al. Coronary-artery revascularization before elective major vascular surgery. *N Engl J Med* 2004;351:2795-804.
- 16) Wijeyesundera DN, Beattie WS, Karkouti K, Neuman MD, Austin PC, Laupacis A. Association of echocardiography before major elective non-cardiac surgery with postoperative survival and length of hospital stay: population based cohort study. *BMJ* 2011;342:d3695.
- 17) Wijeyesundera DN, Beattie WS, Austin PC, Hux JE, Laupacis A. Non-invasive cardiac stress testing before elective major non-cardiac surgery: population based cohort study. *BMJ* 2010;340:b5526.

1-6 漢方薬を出すとき

- 1) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ツムラの医療用エキス顆粒：小柴胡湯、葛根湯、芍薬甘草湯、抑肝散、大建中湯、六君子湯、五苓散、補中益気湯、牛車腎気丸、半夏瀉心湯、麦門冬湯、小青竜湯、加味逍遙散、当歸芍薬散、桂枝茯苓丸、防風通聖散、黄連解毒湯、大黃甘草湯、麻子仁丸、八味地黄丸。ワーファリン錠）。
- 2) 厚生労働省。重篤副作用疾患別対応マニュアル 偽アルドステロン症。2006（2022年2月改定）。
- 3) 厚生労働省。重篤副作用疾患別対応マニュアル 間質性肺炎（肺臓炎、胞隔炎、肺線維症）。2006（2019年9月改定）。
- 4) 萬谷直樹，岡洋志，佐橋佳郎，ほか。甘草の使用量と偽アルドステロン症の頻度に関する文献的調査。日本東洋医学雑誌 2015;66:197-202。
- 5) Shimizu S, Kobayashi T, Tomioka H, Ohtsu K, Matsui T, Hibi T. Involvement of herbal medicine as a cause of mesenteric phlebosclerosis: results from a large-scale nationwide survey. J Gastroenterol 2017;52:308-14.
- 6) Ang-Lee MK, Moss J, Yuan CS. Herbal medicines and perioperative care. JAMA 2001;286:208-16.
- 7) Suzuki H, Matsuzaki J, Fukushima Y, et al. Randomized clinical trial: rikkunshito in the treatment of functional dyspepsia—a multicenter, double-blind, randomized, placebo-controlled study. Neurogastroenterol Motil 2014;26:950-61.
- 8) Tominaga K, Sakata Y, Kusunoki H, et al. Rikkunshito simultaneously improves dyspepsia correlated with anxiety in patients with functional dyspepsia: a randomized clinical trial (the DREAM study). Neurogastroenterol Motil 2018;30:e13319.
- 9) 日本消化器病学会編。機能性消化管疾患診療ガイドライン2021—機能性ディスぺプシア（FD）（改訂第2版）。南江堂；2021。
- 10) Yoshikawa K, Shimada M, Wakabayashi G, et al. Effect of daikenchuto, a traditional Japanese herbal medicine, after total gastrectomy for gastric cancer: a multicenter, randomized, double-blind, placebo-controlled, phase II trial. J Am Coll Surg 2015;221:571-8.
- 11) Lech GE, Moraes LBL, Sturmer CM, et al. Effects of daikenchuto (TJ-100) after gastrointestinal oncological surgery: an updated meta-analysis of randomized controlled trials. J Laparoendosc Adv Surg Tech A 2026;36:606-15.
- 12) 日本消化管学会編。便通異常症診療ガイドライン2023—慢性便秘症。南江堂；2023。
- 13) Furukawa K, Tomita N, Uematsu D, et al. Randomized double-blind placebo-controlled multicenter trial of Yokukansan for neuropsychiatric symptoms in Alzheimer's disease. Geriatr Gerontol Int 2017;17:211-8.
- 14) Matsunaga S, Kishi T, Iwata N. Yokukansan in the treatment of behavioral and psychological symptoms of dementia: an updated meta-analysis of randomized controlled trials. J Alzheimers Dis 2016;54:635-43.
- 15) 日本神経学会監修，「認知症疾患診療ガイドライン」作成委員会編。認知症疾患診療ガイドライン 2017。医学書院；2017。
- 16) Shimada S, Arai T, Tamaoka A, Homma M. Licorice-induced hypokalaemia in patients treated with Yokukansan preparations: identification of the risk factors in a retrospective cohort study. BMJ Open 2017;7:e014218.
- 17) 馬場駿吉，高坂知節，稲村直樹，ほか。小青竜湯の通年性鼻アレルギーに対する効果。耳鼻咽喉科臨床 1995;88:389-405。
- 18) Kurita J, Yonekura S, Iinuma T, et al. Evaluation of shoseiryuto for seasonal allergic rhinitis, using an environmental challenge chamber. World Allergy Organ J 2022;15:100636.

- 19) Irifune K, Hamada H, Ito R, et al. Antitussive effect of bakumondoto a fixed kampo medicine (six herbal components) for treatment of post-infectious prolonged cough: controlled clinical pilot study with 19 patients. *Phytomedicine* 2011;18:630-3.
- 20) Nishikawa K, Aoyama T, Oba MS, et al. The clinical impact of Hangeshashinto (TJ-14) in the treatment of chemotherapy-induced oral mucositis in gastric cancer and colorectal cancer: analyses of pooled data from two phase II randomized clinical trials (HANGESHA-G and HANGESHA-C). *J Cancer* 2018;9:1725-30.
- 21) Mori K, Kondo T, Kamiyama Y, Kano Y, Tominaga K. Preventive effect of Kampo medicine (Hangeshashin-to) against irinotecan-induced diarrhea in advanced non-small-cell lung cancer. *Cancer Chemother Pharmacol* 2003;51:403-6.
- 22) Oki E, Emi Y, Kojima H, et al. Preventive effect of Goshajinkigan on peripheral neurotoxicity of FOLFOX therapy (GENIUS trial): a placebo-controlled, double-blind, randomized phase III study. *Int J Clin Oncol* 2015;20:767-75.
- 23) Fujisawa N, Oya S, Yoshida S, et al. A prospective randomized study on the preventive effect of Japanese herbal Kampo medicine goreisan for recurrence of chronic subdural hematoma. *Neurol Med Chir (Tokyo)* 2021;61:12-20.
- 24) Satoh N, Sakai S, Kogure T, et al. A randomized double blind placebo-controlled clinical trial of Hochuekkito, a traditional herbal medicine, in the treatment of elderly patients with weakness N of one and responder restricted design. *Phytomedicine* 2005;12:549-54.
- 25) Ota K, Fukui K, Nakamura E, et al. Effect of Shakuyaku-kanzo-to in patients with muscle cramps: a systematic literature review. *J Gen Fam Med* 2020;21:56-62.

1-7 制度と書類

- 1) 厚生労働省. 高額療養費制度を利用される皆さまへ（2026年7月31日更新）. 2026年10月閲覧.
- 2) 難病情報センター. 指定難病患者への医療費助成制度のご案内. 2026年10月閲覧.
- 3) 厚生労働省. 自立支援医療制度の概要、自立支援医療（更生医療）の概要、自立支援医療（精神通院医療）の概要. 2026年10月閲覧.
- 4) 厚生省社会局長通知. 生活保護法による医療扶助運営要領について（社発第727号）. 昭和36年9月30日.
- 5) 全国健康保険協会. 病気やケガで会社を休んだとき（傷病手当金）. 2026年10月閲覧.
- 6) 厚生労働省. 治療と就業の両立支援指針（令和8年厚生労働省告示第28号）. 2026.
- 7) 厚生労働省. 身体障害者手帳制度の概要. 2026年10月閲覧.
- 8) 日本年金機構. 障害年金（障害基礎年金・障害厚生年金の受給要件・請求時期・年金額）. 2026年10月閲覧.
- 9) 厚生労働省. 令和8年4月時点の指定難病（告示番号1～348）. 2026年10月閲覧.
- 10) 難病の患者に対する医療等に関する法律（平成26年法律第50号）第6条、第7条.
- 11) 社会福祉法（昭和26年法律第45号）第2条第3項第9号.
- 12) 国民健康保険法（昭和33年法律第192号）第58条第2項.
- 13) 全国健康保険協会. 健康保険 傷病手当金支給申請書（療養担当者記入用）、および北海道支部「傷病手当金について」（2026年3月13日）. 2026年10月閲覧.
- 14) 厚生労働省. 改訂 心の健康問題により休業した労働者の職場復帰支援の手引き. 2009.
- 15) 厚生労働省労働基準局安全衛生部労働衛生課長. 治療と就業の両立支援に関する診療報酬の改定について（基安労発0305第3号）. 令和8年3月5日.
- 16) 身体障害者福祉法（昭和24年法律第283号）第15条.
- 17) 厚生労働省社会・援護局障害保健福祉部長通知. 身体障害者障害程度等級表の解説（身体障害認定基準）について（障発第0110001号）. 平成15年1月10日（改正後全文）.

- 18) 日本年金機構. 障害認定日（年金用語集）. 2026年10月閲覧.
- 19) 日本年金機構. 障害年金ガイド 令和8年度版. 2026.
- 20) 精神保健及び精神障害者福祉に関する法律（昭和25年法律第123号）第45条、同施行規則 第23条.
- 21) 厚生労働省老健局老人保健課長. 要介護認定における「認定調査票記入の手引き」、「主治医意見書記入の手引き」及び「特定疾病にかかる診断基準」について（平成21年9月30日 老老発0930第2号、令和8年4月改正）.
- 22) 介護保険法（平成9年法律第123号）第29条.
- 23) 道路交通法（昭和35年法律第105号）第101条の6、第103条、第117条の4.
- 24) 道路交通法施行令（昭和35年政令第270号）第33条の2の3.
- 25) 警察庁. 安全運転相談窓口について. 2026年10月閲覧.
- 26) 日本神経学会, 日本神経治療学会, 日本認知症学会, 日本老年医学会, 日本老年精神医学会. わが国における運転免許証に係る認知症等の診断の届出ガイドライン. 2014.
- 27) 医師法（昭和23年法律第201号）第20条.
- 28) 日本年金機構. 国民年金・厚生年金保険 障害認定基準（令和4年4月1日改正）.

第2部 外来でよくある訴え

2-1 かぜ症候群・咽頭痛・急性気管支炎

- 1) 厚生労働省健康・生活衛生局感染症対策部感染症対策課編. 抗微生物薬適正使用の手引き 第四版 医科・外来編. 2026.
- 2) 日本呼吸器学会成人肺炎診療ガイドライン2024作成委員会. 成人肺炎診療ガイドライン2024. 日本呼吸器学会；2024.
- 3) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認. タミフルカプセル、ゾフルーザ錠、パキロビッドバック、ゾコーバ錠、カロナール錠、トランサミン錠、サワシリンカプセル、オーグメンチン配合錠、ミノマイシン錠、ジスロマック錠、ケフレックスカプセル、ラゲブリオカプセル、PL配合顆粒、ツムラ葛根湯エキスパッケージ、ツムラ麻黄湯エキスパッケージ、メジコン錠）.
- 4) Kenealy T, Arroll B. Antibiotics for the common cold and acute purulent rhinitis. Cochrane Database Syst Rev 2025;11:CD000247.
- 5) Ebell MH, Lundgren J, Youngpairoj S. How long does a cough last? Comparing patients' expectations with data from a systematic review of the literature. Ann Fam Med 2013;11:5-13.
- 6) Smith SM, Fahey T, Smucny J, Becker LA. Antibiotics for acute bronchitis. Cochrane Database Syst Rev 2017;6:CD000245.
- 7) Cornia PB, Hersh AL, Lipsky BA, Newman TB, Gonzales R. Does this coughing adolescent or adult patient have pertussis? JAMA 2010;304:890-6.
- 8) Fine AM, Nizet V, Mandl KD. Large-scale validation of the Centor and McIsaac scores to predict group A streptococcal pharyngitis. Arch Intern Med 2012;172:847-52.
- 9) Stewart EH, Davis B, Clemans-Taylor BL, Littenberg B, Estrada CA, Centor RM. Rapid antigen group A streptococcus test to diagnose pharyngitis: a systematic review and meta-analysis. PLoS One 2014;9:e111727.
- 10) 日本感染症学会気道感染症抗菌薬適正使用委員会. 気道感染症の抗菌薬適正使用に関する提言（改訂版）. 感染症学雑誌 2022;96(Suppl):S1-22.
- 11) Spinks A, Glasziou PP, Del Mar CB. Antibiotics for treatment of sore throat in children and adults. Cochrane Database Syst Rev 2021;12:CD000023.
- 12) Ebell MH, Call M, Shinholser J, Gardner J. Does this patient have infectious mononucleosis? The rational clinical examination systematic review. JAMA 2016;315:1502-9.

- 13) Chartrand C, Leeflang MM, Minion J, Brewer T, Pai M. Accuracy of rapid influenza diagnostic tests: a meta-analysis. *Ann Intern Med* 2012;156:500-11.
- 14) Call SA, Vollenweider MA, Hornung CA, Simel DL, McKinney WP. Does this patient have influenza? *JAMA* 2005;293:987-97.
- 15) Dinnes J, Sharma P, Berhane S, et al. Rapid, point-of-care antigen tests for diagnosis of SARS-CoV-2 infection. *Cochrane Database Syst Rev* 2022;7:CD013705.
- 16) Dobson J, Whitley RJ, Pocock S, Monto AS. Oseltamivir treatment for influenza in adults: a meta-analysis of randomised controlled trials. *Lancet* 2015;385:1729-37.
- 17) Hayden FG, Sugaya N, Hirotsu N, et al. Baloxavir marboxil for uncomplicated influenza in adults and adolescents. *N Engl J Med* 2018;379:913-23.
- 18) Monto AS, Kuhlbusch K, Bernasconi C, et al. Efficacy of baloxavir treatment in preventing transmission of influenza. *N Engl J Med* 2025;392:1582-93.
- 19) Yoshino T, Arita R, Horiba Y, Watanabe K. The use of maoto (Ma-Huang-Tang), a traditional Japanese Kampo medicine, to alleviate flu symptoms: a systematic review and meta-analysis. *BMC Complement Altern Med* 2019;19:68.
- 20) Hammond J, Leister-Tebbe H, Gardner A, et al. Oral nirmatrelvir for high-risk, nonhospitalized adults with Covid-19. *N Engl J Med* 2022;386:1397-408.
- 21) Hammond J, Fountaine RJ, Yunis C, et al. Nirmatrelvir for vaccinated or unvaccinated adult outpatients with Covid-19. *N Engl J Med* 2024;390:1186-95.
- 22) Yotsuyanagi H, Ohmagari N, Doi Y, et al. Efficacy and safety of 5-day oral ensitrelvir for patients with mild to moderate COVID-19: the SCORPIO-SR randomized clinical trial. *JAMA Netw Open* 2024;7:e2354991.
- 23) Jayk Bernal A, Gomes da Silva MM, Musungaie DB, et al. Molnupiravir for oral treatment of Covid-19 in nonhospitalized patients. *N Engl J Med* 2022;386:509-20.
- 24) Smith SM, Schroeder K, Fahey T. Over-the-counter (OTC) medications for acute cough in children and adults in community settings. *Cochrane Database Syst Rev* 2014;(11):CD001831.
- 25) Abuelgasim H, Albury C, Lee J. Effectiveness of honey for symptomatic relief in upper respiratory tract infections: a systematic review and meta-analysis. *BMJ Evid Based Med* 2021;26:57-64.
- 26) 学校保健安全法施行規則（昭和33年文部省令第18号）第18条、第19条。
- 27) Spurling GK, Dooley L, Clark J, Askew DA. Immediate versus delayed versus no antibiotics for respiratory infections. *Cochrane Database Syst Rev* 2023;10:CD004417.
- 28) Okabayashi S, Goto M, Kawamura T, et al. Non-superiority of Kakkonto, a Japanese herbal medicine, to a representative multiple cold medicine with respect to anti-aggravation effects on the common cold: a randomized controlled trial. *Intern Med* 2014;53:949-56.
- 29) De Sutter AI, Eriksson L, van Driel ML. Oral antihistamine-decongestant-analgesic combinations for the common cold. *Cochrane Database Syst Rev* 2022;1:CD004976.

2-2 鼻炎・花粉症

- 1) 松原篤, 坂下雅文, 後藤穰, ほか. 鼻アレルギーの全国疫学調査2019（1998年, 2008年との比較）: 速報—耳鼻咽喉科医およびその家族を対象として. *日耳鼻* 2020;123:485-90.
- 2) Okubo K, Kurono Y, Ichimura K, et al. Japanese guidelines for allergic rhinitis 2020. *Allergol Int* 2020;69:331-45.（日本語の最新版は、日本耳鼻咽喉科免疫アレルギー感染症学会編. 鼻アレルギー診療ガイドライン—通年性鼻炎と花粉症—2024年版（改訂第10版）. 金原出版; 2024.）
- 3) 厚生労働省健康・生活衛生局感染症対策部感染症対策課編. 抗微生物薬適正使用の手引き 第四版 医科・外来編. 2026.

- 4) Tokunaga T, Sakashita M, Haruna T, et al. Novel scoring system and algorithm for classifying chronic rhinosinusitis: the JESREC Study. *Allergy* 2015;70:995-1003.
- 5) 厚生労働省. オマリズマブ（遺伝子組換え）製剤の最適使用推進ガイドライン（季節性アレルギー性鼻炎）. 令和6年5月改訂（医薬薬審発0521第2号）.
- 6) Weiner JM, Abramson MJ, Puy RM. Intranasal corticosteroids versus oral H1 receptor antagonists in allergic rhinitis: systematic review of randomised controlled trials. *BMJ* 1998;317:1624-9.
- 7) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。アレグラ錠、クラリチン錠、デザレックス錠、ピラノア錠、タリオン錠、アレジオン錠、ザイザル錠、アレロック錠、ルパフィン錠、ポララミン錠、ブリビナ液、ナゾネックス点鼻液、アラミスト点鼻液、ディレグラ配合錠、シングレア錠、アレジオンLX点眼液、シダキュアスギ花粉舌下錠、ミティキュアダニ舌下錠、ゾレア皮下注、フルメトロン点眼液、ハルナールD錠、ツムラ小竜竜湯エキス顆粒）.
- 8) Wilson AM, O'Byrne PM, Parameswaran K. Leukotriene receptor antagonists for allergic rhinitis: a systematic review and meta-analysis. *Am J Med* 2004;116:338-44.
- 9) Nurmatov U, van Schayck CP, Hurwitz B, Sheikh A. House dust mite avoidance measures for perennial allergic rhinitis: an updated Cochrane systematic review. *Allergy* 2012;67:158-65.
- 10) Head K, Snidvongs K, Glew S, et al. Saline irrigation for allergic rhinitis. *Cochrane Database Syst Rev* 2018;6:CD012597.
- 11) Yonekura S, Gotoh M, Kaneko S, Maekawa Y, Okubo K, Okamoto Y. Disease-modifying effect of Japanese cedar pollen sublingual immunotherapy tablets. *J Allergy Clin Immunol Pract* 2021;9:4103-16.e14.
- 12) Okubo K, Masuyama K, Imai T, et al. Efficacy and safety of the SQ house dust mite sublingual immunotherapy tablet in Japanese adults and adolescents with house dust mite-induced allergic rhinitis. *J Allergy Clin Immunol* 2017;139:1840-8.e10.
- 13) Okubo K, Okano M, Sato N, et al. Add-on omalizumab for inadequately controlled severe pollinosis despite standard-of-care: a randomized study. *J Allergy Clin Immunol Pract* 2020;8:3130-40.e2.
- 14) Seresirikachorn K, Chitsuthipakorn W, Kanjanawasee D, Khattiyawittayakun L, Snidvongs K. Effects of H1 antihistamine addition to intranasal corticosteroid for allergic rhinitis: a systematic review and meta-analysis. *Int Forum Allergy Rhinol* 2018;8:1083-92.
- 15) Howarth PH, Stern MA, Roi L, Reynolds R, Bousquet J. Double-blind, placebo-controlled study comparing the efficacy and safety of fexofenadine hydrochloride (120 and 180 mg once daily) and cetirizine in seasonal allergic rhinitis. *J Allergy Clin Immunol* 1999;104:927-33.
- 16) Hampel F, Ratner P, Mansfield L, Meeves S, Liao Y, Georges G. Fexofenadine hydrochloride, 180 mg, exhibits equivalent efficacy to cetirizine, 10 mg, with less drowsiness in patients with moderate-to-severe seasonal allergic rhinitis. *Ann Allergy Asthma Immunol* 2003;91:354-61.
- 17) Vieira RJ, Gil-Mata S, Ferreira A, et al. Efficacy and safety of oral antihistamines for allergic rhinitis: network meta-analysis. *J Allergy Clin Immunol Pract* 2026;14:1175-83.
- 18) Kawauchi H, Yanai K, Wang DY, Itahashi K, Okubo K. Antihistamines for allergic rhinitis treatment from the viewpoint of nonsedative properties. *Int J Mol Sci* 2019;20:213.

2-3 長引く咳

- 1) 厚生労働省健康・生活衛生局感染症対策部感染症対策課編. 抗微生物薬適正使用の手引き 第四版 医科・外来編. 2026.
- 2) National Institute for Health and Care Excellence. Suspected cancer: recognition and referral (NICE guideline NG12). 2015（2026年4月更新）.

12 文献

- 3) Griffith DE, Brown-Elliott BA, Langsjoen B, et al. Clinical and molecular analysis of macrolide resistance in *Mycobacterium avium* complex lung disease. *Am J Respir Crit Care Med* 2006;174:928-34.
- 4) 日本結核・非結核性抗酸菌症学会 非結核性抗酸菌症対策委員会, 日本呼吸器学会 感染症・結核学術部会. 肺非結核性抗酸菌症診断に関する指針—2024年改訂. *結核* 2024;99:267-70.
- 5) Daley CL, Iaccarino JM, Lange C, et al. Treatment of nontuberculous mycobacterial pulmonary disease: an official ATS/ERS/ESCMID/IDSA clinical practice guideline. *Clin Infect Dis* 2020;71:905-13.
- 6) Fujimura M, Abo M, Ogawa H, et al. Importance of atopic cough, cough variant asthma and sinobronchial syndrome as causes of chronic cough in the Hokuriku area of Japan. *Respirology* 2005;10:201-7.
- 7) Matsumoto H, Niimi A, Takemura M, et al. Prevalence and clinical manifestations of gastro-oesophageal reflux-associated chronic cough in the Japanese population. *Cough* 2007;3:1.
- 8) Dicpinigaitis PV. Angiotensin-converting enzyme inhibitor-induced cough: ACCP evidence-based clinical practice guidelines. *Chest* 2006;129(1 Suppl):169S-73S.
- 9) 日本アレルギー学会喘息ガイドライン専門部会監修. 喘息予防・管理ガイドライン2024. 協和企画; 2024.
- 10) Okubo K, Kurono Y, Ichimura K, et al. Japanese guidelines for allergic rhinitis 2020. *Allergol Int* 2020;69:331-45.
- 11) Raghu G, Remy-Jardin M, Ryerson CJ, et al. Diagnosis of hypersensitivity pneumonitis in adults: an official ATS/JRS/ALAT clinical practice guideline. *Am J Respir Crit Care Med* 2020;202:e36-69.
- 12) Moore A, Ashdown HF, Shinkins B, et al. Clinical characteristics of pertussis-associated cough in adults and children: a diagnostic systematic review and meta-analysis. *Chest* 2017;152:353-67.
- 13) 日本呼吸器学会 咳嗽・喀痰の診療ガイドライン2019作成委員会編. 咳嗽・喀痰の診療ガイドライン 2019. メディカルレビュー社; 2019.
- 14) Song WJ, Kim HJ, Shim JS, et al. Diagnostic accuracy of fractional exhaled nitric oxide measurement in predicting cough-variant asthma and eosinophilic bronchitis in adults with chronic cough: a systematic review and meta-analysis. *J Allergy Clin Immunol* 2017;140:701-9.
- 15) Matsunaga K, Hirano T, Akamatsu K, et al. Exhaled nitric oxide cutoff values for asthma diagnosis according to rhinitis and smoking status in Japanese subjects. *Allergol Int* 2011;60:331-7.
- 16) Matsunaga K, Hirano T, Kawayama T, et al. Reference ranges for exhaled nitric oxide fraction in healthy Japanese adult population. *Allergol Int* 2010;59:363-7.
- 17) Ebell MH, Lundgren J, Youngpairoj S. How long does a cough last? Comparing patients' expectations with data from a systematic review of the literature. *Ann Fam Med* 2013;11:5-13.
- 18) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。メジコン錠、メブチン錠、シムビコートタービュヘイラー、ネキシウムカプセル、リフヌア錠、ミカルディス錠、チモプトール点眼液、ムコダイン錠、コデインリン酸塩錠、エリスロシン錠、クラリス錠、ツムラ麦門冬湯エキス顆粒)。
- 19) Wang K, Birring SS, Taylor K, et al. Montelukast for postinfectious cough in adults: a double-blind randomised placebo-controlled trial. *Lancet Respir Med* 2014;2:35-43.
- 20) Fujimura M, Ogawa H, Nishizawa Y, Nishi K. Comparison of atopic cough with cough variant asthma: is atopic cough a precursor of asthma? *Thorax* 2003;58:14-8.
- 21) Kahrilas PJ, Howden CW, Hughes N, Molloy-Bland M. Response of chronic cough to acid-suppressive therapy in patients with gastroesophageal reflux disease. *Chest* 2013;143:605-12.
- 22) Smith SM, Schroeder K, Fahey T. Over-the-counter (OTC) medications for acute cough in children and adults in community settings. *Cochrane Database Syst Rev* 2014;(11):CD001831.

- 23) Zheng JP, Kang J, Huang SG, et al. Effect of carbocysteine on acute exacerbation of chronic obstructive pulmonary disease (PEACE Study): a randomised placebo-controlled study. *Lancet* 2008;371:2013-8.
- 24) McGarvey LP, Birring SS, Morice AH, et al. Efficacy and safety of gefapixant, a P2X3 receptor antagonist, in refractory chronic cough and unexplained chronic cough (COUGH-1 and COUGH-2): results from two double-blind, randomised, parallel-group, placebo-controlled, phase 3 trials. *Lancet* 2022;399:909-23.
- 25) Bradley SH, Abraham S, Callister ME, et al. Sensitivity of chest X-ray for detecting lung cancer in people presenting with symptoms: a systematic review. *Br J Gen Pract* 2019;69:e827-35.
- 26) Kudoh S, Azuma A, Yamamoto M, Izumi T, Ando M. Improvement of survival in patients with diffuse panbronchiolitis treated with low-dose erythromycin. *Am J Respir Crit Care Med* 1998;157:1829-32.
- 27) Hosono Y, Kitada S, Yano Y, et al. The association between erythromycin monotherapy for *Mycobacterium avium* complex lung disease and cross-resistance to clarithromycin: a retrospective case-series study. *J Infect Chemother* 2018;24:353-7.

2-4 続く発熱・倦怠感・体重減少

- 1) Stadje R, Dornieden K, Baum E, et al. The differential diagnosis of tiredness: a systematic review. *BMC Fam Pract* 2016;17:147.
- 2) Nicholson BD, Virdee P, Aveyard P, et al. Prioritising primary care patients with unexpected weight loss for cancer investigation: diagnostic accuracy study (update). *BMJ* 2024;387:e080199.
- 3) Metalidis C, Knockaert DC, Bobbaers H, Vanderschueren S. Involuntary weight loss. Does a negative baseline evaluation provide adequate reassurance? *Eur J Intern Med* 2008;19:345-9.
- 4) Bosch X, Monclús E, Escoda O, et al. Unintentional weight loss: clinical characteristics and outcomes in a prospective cohort of 2677 patients. *PLoS One* 2017;12:e0175125.
- 5) Koch H, van Bokhoven MA, ter Riet G, et al. Ordering blood tests for patients with unexplained fatigue in general practice: what does it yield? Results of the VAMPIRE trial. *Br J Gen Pract* 2009;59:e93-100.
- 6) Institute of Medicine. Beyond Myalgic Encephalomyelitis/Chronic Fatigue Syndrome: Redefining an Illness. Washington, DC: National Academies Press; 2015.
- 7) Naito T, Mizooka M, Mitsumoto F, et al. Diagnostic workup for fever of unknown origin: a multicenter collaborative retrospective study. *BMJ Open* 2013;3:e003971.
- 8) Yamaguchi M, Ohta A, Tsunematsu T, et al. Preliminary criteria for classification of adult Still's disease. *J Rheumatol* 1992;19:424-30.
- 9) 日本循環器学会ほか. 2026年改訂版 大型血管炎診療ガイドライン. 2026.
- 10) Kwong JS, Carignan S, Kang EY, Müller NL, FitzGerald JM. Miliary tuberculosis. Diagnostic accuracy of chest radiography. *Chest* 1996;110:339-42.
- 11) Fan L, Chen Z, Hao XH, Hu ZY, Xiao HP. Interferon-gamma release assays for the diagnosis of extrapulmonary tuberculosis: a systematic review and meta-analysis. *FEMS Immunol Med Microbiol* 2012;65:456-66.
- 12) Oka T. Psychogenic fever: how psychological stress affects body temperature in the clinical population. *Temperature (Austin)* 2015;2:368-78.
- 13) Wreghitt TG, Teare EL, Sule O, Devi R, Rice P. Cytomegalovirus infection in immunocompetent patients. *Clin Infect Dis* 2003;37:1603-6.
- 14) Murase T, Yamaguchi M, Suzuki R, et al. Intravascular large B-cell lymphoma (IVLBCL): a clinicopathologic study of 96 cases with special reference to the immunophenotypic heterogeneity of CD5. *Blood* 2007;109:478-85.

- 15) Fincher RM, Page MI. Clinical significance of extreme elevation of the erythrocyte sedimentation rate. *Arch Intern Med* 1986;146:1581-3.
- 16) Takeuchi M, Dahabreh IJ, Nihashi T, Iwata M, Varghese GM, Terasawa T. Nuclear imaging for classic fever of unknown origin: meta-analysis. *J Nucl Med* 2016;57:1913-9.
- 17) Buttgereit F, Dejaco C, Matteson EL, Dasgupta B. Polymyalgia rheumatica and giant cell arteritis: a systematic review. *JAMA* 2016;315:2442-58.
- 18) Dasgupta B, Cimmino MA, Maradit-Kremers H, et al. 2012 provisional classification criteria for polymyalgia rheumatica: a European League Against Rheumatism/American College of Rheumatology collaborative initiative. *Ann Rheum Dis* 2012;71:484-92.
- 19) Dejaco C, Singh YP, Perel P, et al. 2015 Recommendations for the management of polymyalgia rheumatica: a European League Against Rheumatism/American College of Rheumatology collaborative initiative. *Ann Rheum Dis* 2015;74:1799-807.
- 20) Knockaert DC, Dujardin KS, Bobbaers HJ. Long-term follow-up of patients with undiagnosed fever of unknown origin. *Arch Intern Med* 1996;156:618-20.
- 21) Moore C Jr, Ormseth M, Fuchs H. Causes and significance of markedly elevated serum ferritin levels in an academic medical center. *J Clin Rheumatol* 2013;19:324-8.
- 22) Schram AM, Campigotto F, Mullally A, et al. Marked hyperferritinemia does not predict for HLH in the adult population. *Blood* 2015;125:1548-52.
- 23) Fautrel B, Le Moël G, Saint-Marcoux B, et al. Diagnostic value of ferritin and glycosylated ferritin in adult onset Still's disease. *J Rheumatol* 2001;28:322-9.

2-5 リンパ節腫脹

- 1) Fijten GH, Blijham GH. Unexplained lymphadenopathy in family practice. An evaluation of the probability of malignant causes and the effectiveness of physicians' workup. *J Fam Pract* 1988;27:373-6.
- 2) Chau I, Kelleher MT, Cunningham D, et al. Rapid access multidisciplinary lymph node diagnostic clinic: analysis of 550 patients. *Br J Cancer* 2003;88:354-61.
- 3) Gaddey HL, Riegel AM. Unexplained lymphadenopathy: evaluation and differential diagnosis. *Am Fam Physician* 2016;94:896-903.
- 4) Ellison E, LaPuerta P, Martin SE. Supraclavicular masses: results of a series of 309 cases biopsied by fine needle aspiration. *Head Neck* 1999;21:239-46.
- 5) Vassilakopoulos TP, Pangalis GA. Application of a prediction rule to select which patients presenting with lymphadenopathy should undergo a lymph node biopsy. *Medicine (Baltimore)* 2000;79:338-47.
- 6) National Institute for Health and Care Excellence. Suspected cancer: recognition and referral (NICE guideline NG12). 2015 (2026年4月更新) .
- 7) Kanda N, Yamaguchi R, Yamamoto Y, Matsumura M, Hatakeyama S. Performance of serum soluble interleukin-2 receptor as a diagnostic marker for lymphoma in patients with fever. *Sci Rep* 2023;13:18784.
- 8) Chen X, Li Y, Ding X, et al. The significance of serum levels of soluble interleukin-2 receptor in patients undergoing maintenance hemodialysis. *Ren Fail* 2020;42:419-27.
- 9) Steinkamp HJ, Cornehl M, Hosten N, Pegios W, Vogl T, Felix R. Cervical lymphadenopathy: ratio of long- to short-axis diameter as a predictor of malignancy. *Br J Radiol* 1995;68:266-70.
- 10) Ahuja AT, Ying M. Sonographic evaluation of cervical lymph nodes. *AJR Am J Roentgenol* 2005;184:1691-9.
- 11) Ebell MH, Call M, Shinholser J, Gardner J. Does this patient have infectious mononucleosis? The rational clinical examination systematic review. *JAMA* 2016;315:1502-9.

- 12) Kucukardali Y, Solmazgul E, Kunter E, Oncul O, Yildirim S, Kaplan M. Kikuchi-Fujimoto disease: analysis of 244 cases. *Clin Rheumatol* 2007;26:50-4.
- 13) 日本血液学会編. 造血器腫瘍診療ガイドライン 第3.1版 (2024年版). 第II章 リンパ腫 悪性リンパ腫 総論.
- 14) Bartlett A, Williams R, Hilton M. Splenic rupture in infectious mononucleosis: a systematic review of published case reports. *Injury* 2016;47:531-8.
- 15) Fontanilla JM, Barnes A, von Reyn CF. Current diagnosis and management of peripheral tuberculous lymphadenitis. *Clin Infect Dis* 2011;53:555-62.
- 16) Frederiksen JK, Sharma M, Casulo C, Burack WR. Systematic review of the effectiveness of fine-needle aspiration and/or core needle biopsy for subclassifying lymphoma. *Arch Pathol Lab Med* 2015;139:245-51.
- 17) Kuramoto N, Saito S, Fujii T, et al. Characteristics of rheumatoid arthritis with immunodeficiency-associated lymphoproliferative disorders to regress spontaneously by the withdrawal of methotrexate and their clinical course: a retrospective, multicenter, case-control study. *Mod Rheumatol* 2022;32:24-31.
- 18) セファレキシンの電子添文 (KEGG MEDICUSで2026年10月に確認。ケフレックスカプセル) .
- 19) Wolfson S, Kim E, Plaunova A, et al. Axillary adenopathy after COVID-19 vaccine: no reason to delay screening mammogram. *Radiology* 2022;303:297-9.
- 20) Bass JW, Freitas BC, Freitas AD, et al. Prospective randomized double blind placebo-controlled evaluation of azithromycin for treatment of cat-scratch disease. *Pediatr Infect Dis J* 1998;17:447-52.
- 21) Stevens DL, Bisno AL, Chambers HF, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2014;59:e10-52.
- 22) Pynnonen MA, Gillespie MB, Roman B, et al. Clinical practice guideline: evaluation of the neck mass in adults. *Otolaryngol Head Neck Surg* 2017;157(2 Suppl):S1-30.
- 23) Kühnl A, Cunningham D, Hutka M, et al. Rapid access clinic for unexplained lymphadenopathy and suspected malignancy: prospective analysis of 1000 patients. *BMC Hematol* 2018;18:19.

2-6 貧血

- 1) Guralnik JM, Eisenstaedt RS, Ferrucci L, Klein HG, Woodman RC. Prevalence of anemia in persons 65 years and older in the United States: evidence for a high rate of unexplained anemia. *Blood* 2004;104:2263-8.
- 2) World Health Organization. Guideline on haemoglobin cutoffs to define anaemia in individuals and populations. Geneva: WHO; 2024.
- 3) 日本鉄バイオサイエンス学会編. 鉄欠乏性貧血の診療指針. フジメディカル出版 ; 2024.
- 4) Sheth TN, Choudhry NK, Bowes M, Detsky AS. The relation of conjunctival pallor to the presence of anemia. *J Gen Intern Med* 1997;12:102-6.
- 5) Rockey DC, Altayar O, Falck-Ytter Y, Kalmaz D. AGA technical review on gastrointestinal evaluation of iron deficiency anemia. *Gastroenterology* 2020;159:1097-119.
- 6) Ko CW, Siddique SM, Patel A, et al. AGA clinical practice guidelines on the gastrointestinal evaluation of iron deficiency anemia. *Gastroenterology* 2020;159:1085-94.
- 7) Snook J, Bhala N, Beales ILP, et al. British Society of Gastroenterology guidelines for the management of iron deficiency anaemia in adults. *Gut* 2021;70:2030-51.
- 8) Hoffmann JJ, Urrechaga E, Aguirre U. Discriminant indices for distinguishing thalassemia and iron deficiency in patients with microcytic anemia: a meta-analysis. *Clin Chem Lab Med* 2015;53:1883-94.

- 9) Graham DY, Opekun AR, Willingham FF, Qureshi WA. Visible small-intestinal mucosal injury in chronic NSAID users. *Clin Gastroenterol Hepatol* 2005;3:55-9.
- 10) Komatsu N, Arita K, Mitsui H, Nemoto T, Hanaki K. Efficacy and safety of ferric citrate hydrate compared with sodium ferrous citrate in Japanese patients with iron deficiency anemia: a randomized, double-blind, phase 3 non-inferiority study. *Int J Hematol* 2021;114:8-17.
- 11) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。フェロミア錠、リオナ錠、ノベルジン錠、フェロ・グラデュメット錠）。
- 12) Okam MM, Koch TA, Tran MH. Iron supplementation, response in iron-deficiency anemia: analysis of five trials. *Am J Med* 2017;130:991.e1-8.
- 13) Lam JR, Schneider JL, Zhao W, Corley DA. Proton pump inhibitor and histamine 2 receptor antagonist use and vitamin B12 deficiency. *JAMA* 2013;310:2435-42.
- 14) Vannella L, Lahner E, Osborn J, Annibale B. Systematic review: gastric cancer incidence in pernicious anaemia. *Aliment Pharmacol Ther* 2013;37:375-82.
- 15) 日本透析医学会. 2015年版 慢性腎臓病患者における腎性貧血治療のガイドライン. 透析会誌 2016;49:89-158.
- 16) Rajkumar SV, Dimopoulos MA, Palumbo A, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol* 2014;15:e538-48.
- 17) Kyle RA, Larson DR, Therneau TM, et al. Long-term follow-up of monoclonal gammopathy of undetermined significance. *N Engl J Med* 2018;378:241-9.
- 18) Gangat N, Szuber N, Tefferi A. JAK2 unmutated erythrocytosis: 2026 update on diagnosis and management. *Am J Hematol* 2025;100:2333-54.
- 19) Barbui T, Thiele J, Gisslinger H, et al. The 2016 WHO classification and diagnostic criteria for myeloproliferative neoplasms: document summary and in-depth discussion. *Blood Cancer J* 2018;8:15.
- 20) Tefferi A, Barbui T. Polycythemia vera: 2024 update on diagnosis, risk-stratification, and management. *Am J Hematol* 2023;98:1465-87.
- 21) Marchioli R, Finazzi G, Specchia G, et al. Cardiovascular events and intensity of treatment in polycythemia vera. *N Engl J Med* 2013;368:22-33.
- 22) Griesshammer M, Bangertner M, Sauer T, Wennauer R, Bergmann L, Heimpel H. Aetiology and clinical significance of thrombocytosis: analysis of 732 patients with an elevated platelet count. *J Intern Med* 1999;245:295-300.
- 23) Bailey SE, Ukoumunne OC, Shephard EA, Hamilton W. Clinical relevance of thrombocytosis in primary care: a prospective cohort study of cancer incidence using English electronic medical records and cancer registry data. *Br J Gen Pract* 2017;67:e405-13.
- 24) Tefferi A, Vannucchi AM, Barbui T. Essential thrombocythemia: 2024 update on diagnosis, risk stratification, and management. *Am J Hematol* 2024;99:697-718.
- 25) 柏木浩和, 桑名正隆, 村田満, ほか. 成人免疫性血小板減少症診断参照ガイド 2023年版. 臨床血液 2023;64:1245-57.
- 26) 柏木浩和, 桑名正隆, 羽藤高明, ほか. 成人特発性血小板減少性紫斑病治療の参照ガイド 2019改訂版. 臨床血液 2019;60:877-96.
- 27) Valent P, Klion AD, Horny HP, et al. Contemporary consensus proposal on criteria and classification of eosinophilic disorders and related syndromes. *J Allergy Clin Immunol* 2012;130:607-12.e9.
- 28) Shomali W, Gotlib J. World Health Organization-defined eosinophilic disorders: 2022 update on diagnosis, risk stratification, and management. *Am J Hematol* 2022;97:129-48.
- 29) Vaucher P, Druais PL, Waldvogel S, Favrat B. Effect of iron supplementation on fatigue in nonanemic menstruating women with low ferritin: a randomized controlled trial. *CMAJ* 2012;184:1247-54.

- 30) Arnold DM, Bernotas A, Nazi I, et al. Platelet count response to H. pylori treatment in patients with immune thrombocytopenic purpura with and without H. pylori infection: a systematic review. *Haematologica* 2009;94:850-6.

2-7 頭痛（繰り返す頭痛）

- 1) Wall M, Kupersmith MJ, Kieburtz KD, et al. The idiopathic intracranial hypertension treatment trial: clinical profile at baseline. *JAMA Neurol* 2014;71:693-701.
- 2) Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition. *Cephalalgia* 2018;38:1-211.
- 3) Detsky ME, McDonald DR, Baerlocher MO, Tomlinson GA, McCrory DC, Booth CM. Does this patient with headache have a migraine or need neuroimaging? *JAMA* 2006;296:1274-83.
- 4) Lipton RB, Dodick D, Sadovsky R, et al. A self-administered screener for migraine in primary care: the ID Migraine validation study. *Neurology* 2003;61:375-82.
- 5) 日本神経学会, 日本頭痛学会, 日本神経治療学会監修. 頭痛の診療ガイドライン2021. 医学書院; 2021.
- 6) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）.
- 7) Carlsen LN, Munksgaard SB, Nielsen M, et al. Comparison of 3 treatment strategies for medication overuse headache: a randomized clinical trial. *JAMA Neurol* 2020;77:1069-78.
- 8) 日本頭痛学会理事会. 成人の片頭痛予防療法におけるカルシトニン遺伝子関連ペプチド（CGRP）関連抗体薬の使用に関する日本頭痛学会のポジションステートメント. 2025.
- 9) Lipton RB, Stewart WF, Stone AM, Láinez MJ, Sawyer JP. Stratified care vs step care strategies for migraine: the Disability in Strategies of Care (DISC) Study: a randomized trial. *JAMA* 2000;284:2599-605.
- 10) Karlsson WK, Ostinelli EG, Zhuang ZA, et al. Comparative effects of drug interventions for the acute management of migraine episodes in adults: systematic review and network meta-analysis. *BMJ* 2024;386:e080107.

2-8 めまい・ふらつき

- 1) Kroenke K, Hoffman RM, Einstadter D. How common are various causes of dizziness? A critical review. *South Med J* 2000;93:160-7.
- 2) Maarsingh OR, Dros J, Schellevis FG, et al. Causes of persistent dizziness in elderly patients in primary care. *Ann Fam Med* 2010;8:196-205.
- 3) Lempert T, Olesen J, Furman J, et al. Vestibular migraine: diagnostic criteria. *J Vestib Res* 2022;32:1-6.
- 4) Lopez-Escamez JA, Carey J, Chung WH, et al. Diagnostic criteria for Menière's disease. *J Vestib Res* 2015;25:1-7.
- 5) Staab JP, Eckhardt-Henn A, Horii A, et al. Diagnostic criteria for persistent postural-perceptual dizziness (PPPD): consensus document of the Committee for the Classification of Vestibular Disorders of the Bárány Society. *J Vestib Res* 2017;27:191-208.
- 6) Radtke A, von Brevern M, Tiel-Wilck K, Mainz-Perchalla A, Neuhauser H, Lempert T. Self-treatment of benign paroxysmal positional vertigo: Semont maneuver vs Epley procedure. *Neurology* 2004;63:150-2.
- 7) Adrion C, Fischer CS, Wagner J, et al. Efficacy and safety of betahistine treatment in patients with Meniere's disease: primary results of a long term, multicentre, double blind, randomised, placebo controlled, dose defining trial (BEMED trial). *BMJ* 2016;352:h6816.
- 8) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。メトリジン錠、メリスロン錠、セファドール錠、イソバイドシロップ）.

- 9) McDonnell MN, Hillier SL. Vestibular rehabilitation for unilateral peripheral vestibular dysfunction. *Cochrane Database Syst Rev* 2015;(1):CD005397.
- 10) Yardley L, Barker F, Muller I, et al. Clinical and cost effectiveness of booklet based vestibular rehabilitation for chronic dizziness in primary care: single blind, parallel group, pragmatic, randomised controlled trial. *BMJ* 2012;344:e2237.
- 11) Gibbons CH, Schmidt P, Biaggioni I, et al. The recommendations of a consensus panel for the screening, diagnosis, and treatment of neurogenic orthostatic hypotension and associated supine hypertension. *J Neurol* 2017;264:1567-82.
- 12) Juraschek SP, Hu JR, Cluett JL, et al. Effects of intensive blood pressure treatment on orthostatic hypotension: a systematic review and individual participant-based meta-analysis. *Ann Intern Med* 2021;174:58-68.
- 13) Sheldon RS, Grubb BP, Olshansky B, et al. 2015 Heart Rhythm Society expert consensus statement on the diagnosis and treatment of postural tachycardia syndrome, inappropriate sinus tachycardia, and vasovagal syncope. *Heart Rhythm* 2015;12:e41-63.
- 14) Vernino S, Bourne KM, Stiles LE, et al. Postural orthostatic tachycardia syndrome (POTS): state of the science and clinical care from a 2019 National Institutes of Health Expert Consensus Meeting - Part 1. *Auton Neurosci* 2021;235:102828.
- 15) Bourne KM, Sheldon RS, Hall J, et al. Compression garment reduces orthostatic tachycardia and symptoms in patients with postural orthostatic tachycardia syndrome. *J Am Coll Cardiol* 2021;77:285-96.
- 16) 日本小児心身医学会. 起立性調節障害 (OD) . 学会ウェブサイト (2026年10月閲覧) .
- 17) Jeong SH, Kim JS, Kim HJ, et al. Prevention of benign paroxysmal positional vertigo with vitamin D supplementation: a randomized trial. *Neurology* 2020;95:e1117-25.

2-9 しびれ

- 1) Callaghan BC, Price RS, Feldman EL. Distal symmetric polyneuropathy: a review. *JAMA* 2015;314:2172-81.
- 2) Leonhard SE, Mandarakas MR, Gondim FAA, et al. Diagnosis and management of Guillain-Barré syndrome in ten steps. *Nat Rev Neurol* 2019;15:671-83.
- 3) Chen WH. Cheiro-oral syndrome: a clinical analysis and review of literature. *Yonsei Med J* 2009;50:777-83.
- 4) Tong HC, Haig AJ, Yamakawa K. The Spurling test and cervical radiculopathy. *Spine* 2002;27:156-9.
- 5) Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy. A population-based study from Rochester, Minnesota, 1976 through 1990. *Brain* 1994;117:325-35.
- 6) Ono K, Ebara S, Fuji T, Yonenobu K, Fujiwara K, Yamashita K. Myelopathy hand. New clinical signs of cervical cord damage. *J Bone Joint Surg Br* 1987;69:215-9.
- 7) Lossos A, Siegal T. Numb chin syndrome in cancer patients: etiology, response to treatment, and prognostic significance. *Neurology* 1992;42:1181-4.
- 8) Gwathmey KG. Sensory neuronopathies. *Muscle Nerve* 2016;53:8-19.
- 9) Camdessanché JP, Jousserand G, Ferraud K, et al. The pattern and diagnostic criteria of sensory neuropathy: a case-control study. *Brain* 2009;132:1723-33.
- 10) Schaumburg H, Kaplan J, Windebank A, et al. Sensory neuropathy from pyridoxine abuse. A new megavitamin syndrome. *N Engl J Med* 1983;309:445-8.
- 11) D'Arcy CA, McGee S. The rational clinical examination. Does this patient have carpal tunnel syndrome? *JAMA* 2000;283:3110-7.

- 12) Dabbagh A, MacDermid JC, Yong J, et al. Diagnostic test accuracy of provocative maneuvers for the diagnosis of carpal tunnel syndrome: a systematic review and meta-analysis. *Phys Ther* 2023;103:pzad029.
- 13) Chesterton LS, Blagojevic-Bucknall M, Burton C, et al. The clinical and cost-effectiveness of corticosteroid injection versus night splints for carpal tunnel syndrome (INSTINCTS trial): an open-label, parallel group, randomised controlled trial. *Lancet* 2018;392:1423-33.
- 14) Burton C, Rathod-Mistry T, Blackburn S, et al. The effectiveness of corticosteroid injection versus night splints for carpal tunnel syndrome: 24-month follow-up of a randomized trial. *Rheumatology (Oxford)* 2023;62:546-54.
- 15) Gerritsen AA, de Vet HC, Scholten RJ, et al. Splinting vs surgery in the treatment of carpal tunnel syndrome: a randomized controlled trial. *JAMA* 2002;288:1245-51.
- 16) Sperry BW, Reyes BA, Ikram A, et al. Tenosynovial and cardiac amyloidosis in patients undergoing carpal tunnel release. *J Am Coll Cardiol* 2018;72:2040-50.
- 17) Novak CB, Lee GW, Mackinnon SE, Lay L. Provocative testing for cubital tunnel syndrome. *J Hand Surg Am* 1994;19:817-20.
- 18) Visser NA, Notermans NC, Linssen RS, van den Berg LH, Vrancken AF. Incidence of polyneuropathy in Utrecht, the Netherlands. *Neurology* 2015;84:259-64.
- 19) England JD, Gronseth GS, Franklin G, et al. Practice parameter: evaluation of distal symmetric polyneuropathy: role of laboratory and genetic testing (an evidence-based review). *Neurology* 2009;72:185-92.
- 20) Notermans NC, Wokke JH, van der Graaf Y, Franssen H, van Dijk GW, Jennekens FG. Chronic idiopathic axonal polyneuropathy: a five year follow up. *J Neurol Neurosurg Psychiatry* 1994;57:1525-7.
- 21) Finnerup NB, Attal N, Haroutounian S, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. *Lancet Neurol* 2015;14:162-73.
- 22) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 23) 日本ペインクリニック学会 神経障害性疼痛薬物療法ガイドライン改訂版作成ワーキンググループ編。神経障害性疼痛薬物療法ガイドライン 改訂第2版。真興交易医書出版部；2016。
- 24) Smith EM, Pang H, Cirrincione C, et al. Effect of duloxetine on pain, function, and quality of life among patients with chemotherapy-induced painful peripheral neuropathy: a randomized clinical trial. *JAMA* 2013;309:1359-67.
- 25) de Jager J, Kooy A, Lehter P, et al. Long term treatment with metformin in patients with type 2 diabetes and risk of vitamin B-12 deficiency: randomised placebo controlled trial. *BMJ* 2010;340:c2181.
- 26) Aroda VR, Edelstein SL, Goldberg RB, et al. Long-term metformin use and vitamin B12 deficiency in the Diabetes Prevention Program Outcomes Study. *J Clin Endocrinol Metab* 2016;101:1754-61.
- 27) Sawangjit R, Thongphui S, Chaichompu W, Phumart P. Efficacy and safety of mecobalamin on peripheral neuropathy: a systematic review and meta-analysis of randomized controlled trials. *J Altern Complement Med* 2020;26:1117-29.
- 28) Ang CD, Alviar MJ, Dans AL, et al. Vitamin B for treating peripheral neuropathy. *Cochrane Database Syst Rev* 2008;(3):CD004573.
- 29) Didangelos T, Karlafti E, Kotzakioulafi E, et al. Vitamin B12 supplementation in diabetic neuropathy: a 1-year, randomized, double-blind, placebo-controlled trial. *Nutrients* 2021;13:395.

2-10 動悸・労作時の胸部症状

- 1) Weber BE, Kapoor WN. Evaluation and outcomes of patients with palpitations. *Am J Med* 1996;100:138-48.
- 2) 日本循環器学会, ほか. 急性冠症候群ガイドライン (2018年改訂版). 2019.
- 3) Thavendiranathan P, Bagai A, Khoo C, Dorian P, Choudhry NK. Does this patient with palpitations have a cardiac arrhythmia? *JAMA* 2009;302:2135-43.
- 4) Lessmeier TJ, Gamperling D, Johnson-Liddon V, et al. Unrecognized paroxysmal supraventricular tachycardia: potential for misdiagnosis as panic disorder. *Arch Intern Med* 1997;157:537-43.
- 5) Baman TS, Lange DC, Ilg KJ, et al. Relationship between burden of premature ventricular complexes and left ventricular function. *Heart Rhythm* 2010;7:865-9.
- 6) Alboni P, Botto GL, Baldi N, et al. Outpatient treatment of recent-onset atrial fibrillation with the "pill-in-the-pocket" approach. *N Engl J Med* 2004;351:2384-91.
- 7) Echt DS, Liebson PR, Mitchell LB, et al. Mortality and morbidity in patients receiving encainide, flecainide, or placebo. The Cardiac Arrhythmia Suppression Trial. *N Engl J Med* 1991;324:781-8.
- 8) 日本循環器学会, ほか. 2022年JCSガイドライン フォーカスアップデート版 安定冠動脈疾患の診断と治療. 2022.
- 9) Bösner S, Haasenritter J, Becker A, et al. Ruling out coronary artery disease in primary care: development and validation of a simple prediction rule. *CMAJ* 2010;182:1295-300.
- 10) Haasenritter J, Bösner S, Vaucher P, et al. Ruling out coronary heart disease in primary care: external validation of a clinical prediction rule. *Br J Gen Pract* 2012;62:e415-21.
- 11) Bösner S, Becker A, Hani MA, et al. Chest wall syndrome in primary care patients with chest pain: presentation, associated features and diagnosis. *Fam Pract* 2010;27:363-9.
- 12) Proulx AM, Zryd TW. Costochondritis: diagnosis and treatment. *Am Fam Physician* 2009;80:617-20.
- 13) Knuuti J, Ballo H, Juarez-Orozco LE, et al. The performance of non-invasive tests to rule-in and rule-out significant coronary artery stenosis in patients with stable angina: a meta-analysis focused on post-test disease probability. *Eur Heart J* 2018;39:3322-30.
- 14) SCOT-HEART Investigators. Coronary CT angiography and 5-year risk of myocardial infarction. *N Engl J Med* 2018;379:924-33.
- 15) Maron DJ, Hochman JS, Reynolds HR, et al. Initial invasive or conservative strategy for stable coronary disease. *N Engl J Med* 2020;382:1395-407.
- 16) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。ニトロペン舌下錠、バイアスピリン錠、ミオコルスプレー、メインテート錠、ノルバスク錠、コニール錠)。
- 17) Henrikson CA, Howell EE, Bush DE, et al. Chest pain relief by nitroglycerin does not predict active coronary artery disease. *Ann Intern Med* 2003;139:979-86.
- 18) Etchells E, Bell C, Robb K. Does this patient have an abnormal systolic murmur? *JAMA* 1997;277:564-71.
- 19) Etchells E, Glens V, Shadowitz S, Bell C, Siu S. A bedside clinical prediction rule for detecting moderate or severe aortic stenosis. *J Gen Intern Med* 1998;13:699-704.
- 20) 日本循環器学会, ほか. 2020年改訂版 弁膜症治療のガイドライン. 2020.
- 21) Leon MB, Smith CR, Mack M, et al. Transcatheter aortic-valve implantation for aortic stenosis in patients who cannot undergo surgery. *N Engl J Med* 2010;363:1597-607.
- 22) 日本循環器学会. 2026年改訂版 感染性心内膜炎診療ガイドライン. 2026.
- 23) Perez MV, Mahaffey KW, Hedlin H, et al. Large-scale assessment of a smartwatch to identify atrial fibrillation. *N Engl J Med* 2019;381:1909-17.

- 24) Ataklte F, Erqou S, Laukkanen J, Kaptoge S. Meta-analysis of ventricular premature complexes and their relation to cardiac mortality in general populations. *Am J Cardiol* 2013;112:1263-70.

2-11 浮腫

- 1) Blankfield RP, Finkelhor RS, Alexander JJ, et al. Etiology and diagnosis of bilateral leg edema in primary care. *Am J Med* 1998;105:192-7.
- 2) 日本循環器学会, 日本肺高血圧・肺循環学会. 肺血栓塞栓症・深部静脈血栓症および肺高血圧症に関するガイドライン (2025年改訂版). 2025.
- 3) Geersing GJ, Zuithoff NP, Kearon C, et al. Exclusion of deep vein thrombosis using the Wells rule in clinically important subgroups: individual patient data meta-analysis. *BMJ* 2014;348:g1340.
- 4) Goss JA, Greene AK. Sensitivity and specificity of the Stemmer sign for lymphedema: a clinical lymphoscintigraphic study. *Plast Reconstr Surg Glob Open* 2019;7:e2295.
- 5) Mant J, Doust J, Roalfe A, et al. Systematic review and individual patient data meta-analysis of diagnosis of heart failure, with modelling of implications of different diagnostic strategies in primary care. *Health Technol Assess* 2009;13(32):1-207.
- 6) Chikama R, Hosokawa M, Miyazawa T, Miura R, Suzuki T, Tagami H. Nonepisodic angioedema associated with eosinophilia: report of 4 cases and review of 33 young female patients reported in Japan. *Dermatology* 1998;197:321-5.
- 7) Matsuda M, Fushimi T, Nakamura A, Ikeda S. Nonepisodic angioedema with eosinophilia: a report of two cases and a review of the literature. *Clin Rheumatol* 2006;25:422-5.
- 8) Makani H, Bangalore S, Romero J, et al. Peripheral edema associated with calcium channel blockers: incidence and withdrawal rate—a meta-analysis of randomized trials. *J Hypertens* 2011;29:1270-80.
- 9) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。ノルバスク錠、リリカ、タリージェ錠、アクトス錠、ラシックス錠、ダイアート錠、ルブラック錠、アルダクトンA、ビ・シフロール錠、シンメトレル錠、リフレックス錠)。
- 10) Makani H, Bangalore S, Romero J, Wever-Pinzon O, Messerli FH. Effect of renin-angiotensin system blockade on calcium channel blocker-associated peripheral edema. *Am J Med* 2011;124:128-35.
- 11) Savage RD, Visentin JD, Bronskill SE, et al. Evaluation of a common prescribing cascade of calcium channel blockers and diuretics in older adults with hypertension. *JAMA Intern Med* 2020;180:643-51.
- 12) White SL, Yu R, Craig JC, Polkinghorne KR, Atkins RC, Chadban SJ. Diagnostic accuracy of urine dipsticks for detection of albuminuria in the general community. *Am J Kidney Dis* 2011;58:19-28.
- 13) 厚生労働科学研究費補助金難治性疾患政策研究事業 難治性腎障害に関する調査研究班編. エビデンスに基づくネフローゼ症候群診療ガイドライン2020. 東京医学社 ; 2020.
- 14) 日本循環器学会, 日本心不全学会. 2025年改訂版 心不全診療ガイドライン. 2025.
- 15) Taylor CJ, Ordóñez-Mena JM, Lay-Flurrie SL, et al. Natriuretic peptide testing and heart failure diagnosis in primary care: diagnostic accuracy study. *Br J Gen Pract* 2023;73:e1-8.
- 16) 日本腎臓学会 (作成), 日本医師会 (監修). かかりつけ医から腎臓専門医・専門医療機関への紹介基準. 2018.
- 17) Blankfield RP, Hudgel DW, Tapolyai AA, Zyzanski SJ. Bilateral leg edema, obesity, pulmonary hypertension, and obstructive sleep apnea. *Arch Intern Med* 2000;160:2357-62.

2-12 胸やけ・胃もたれ・心窩部痛

- 1) Nasseri-Moghaddam S, Mousavian AH, Kasaeian A, et al. What is the prevalence of clinically significant endoscopic findings in subjects with dyspepsia? Updated systematic review and meta-analysis. Clin Gastroenterol Hepatol 2023;21:1739-49.e2.
- 2) 日本消化器病学会編. 機能性消化管疾患診療ガイドライン2021—機能性ディスペプシア (FD) (改訂第2版). 南江堂; 2021.
- 3) Vakil N, Moayyedi P, Fennerty MB, Talley NJ. Limited value of alarm features in the diagnosis of upper gastrointestinal malignancy: systematic review and meta-analysis. Gastroenterology 2006;131:390-401.
- 4) 厚生労働省保険局医療課長通知. ヘリコバクター・ピロリ感染の診断及び治療に関する取扱いについて (平成12年10月31日保険発第180号. 平成25年2月21日, 令和4年10月31日一部改正).
- 5) Numans ME, Lau J, de Wit NJ, Bonis PA. Short-term treatment with proton-pump inhibitors as a test for gastroesophageal reflux disease: a meta-analysis of diagnostic test characteristics. Ann Intern Med 2004;140:518-27.
- 6) 日本消化器病学会編. 胃食道逆流症 (GERD) 診療ガイドライン2021 (改訂第3版). 南江堂; 2021.
- 7) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. ネキシウムカプセル、タケブロンOD錠、パリエット錠、ガスター錠、タケキャブ錠、ボノサップバック、ボノピオンパック、アコファイド錠).
- 8) Ferwana M, Abdulmajeed I, Alhajiahmed A, et al. Accuracy of urea breath test in *Helicobacter pylori* infection: meta-analysis. World J Gastroenterol 2015;21:1305-14.
- 9) Gisbert JP, de la Morena F, Abairra V. Accuracy of monoclonal stool antigen test for the diagnosis of *H. pylori* infection: a systematic review and meta-analysis. Am J Gastroenterol 2006;101:1921-30.
- 10) Murakami K, Sakurai Y, Shiino M, Funao N, Nishimura A, Asaka M. Vonoprazan, a novel potassium-competitive acid blocker, as a component of first-line and second-line triple therapy for *Helicobacter pylori* eradication: a phase III, randomised, double-blind study. Gut 2016;65:1439-46.
- 11) Sugimoto M, Kamada T, Ito M, et al. Evidence-based clinical practice guidelines for peptic ulcer disease 2026. J Gastroenterol 2026;61:1361-81. (日本消化器病学会 消化性潰瘍診療ガイドライン2026の英語版)
- 12) Ford AC, Yuan Y, Moayyedi P. *Helicobacter pylori* eradication therapy to prevent gastric cancer: systematic review and meta-analysis. Gut 2020;69:2113-21.
- 13) Matsueda K, Hongo M, Tack J, Saito Y, Kato H. A placebo-controlled trial of acotiamide for meal-related symptoms of functional dyspepsia. Gut 2012;61:821-8.
- 14) Moayyedi P, Eikelboom JW, Bosch J, et al. Safety of proton pump inhibitors based on a large, multi-year, randomized trial of patients receiving rivaroxaban or aspirin. Gastroenterology 2019;157:682-91.e2.
- 15) Rostom A, Dube C, Wells G, et al. Prevention of NSAID-induced gastroduodenal ulcers. Cochrane Database Syst Rev 2002;(4):CD002296.
- 16) Reimer C, Søndergaard B, Hilsted L, Bytzer P. Proton-pump inhibitor therapy induces acid-related symptoms in healthy volunteers after withdrawal of therapy. Gastroenterology 2009;137:80-7.
- 17) Ashida K, Sakurai Y, Hori T, et al. Randomised clinical trial: vonoprazan, a novel potassium-competitive acid blocker, vs. lansoprazole for the healing of erosive oesophagitis. Aliment Pharmacol Ther 2016;43:240-51.
- 18) Laine L, DeVault K, Katz P, et al. Vonoprazan versus lansoprazole for healing and maintenance of healing of erosive esophagitis: a randomized trial. Gastroenterology 2023;164:61-71.

2-13 腹痛・嘔吐

- 1) Viniol A, Keunecke C, Biroga T, et al. Studies of the symptom abdominal pain—a systematic review and meta-analysis. *Fam Pract* 2014;31:517-29.
- 2) 厚生労働省健康・生活衛生局感染症対策部感染症対策課. 抗微生物薬適正使用の手引き 第四版 医科・外来編. 2026.
- 3) Merrett ND, Wilson RB, Cosman P, Biankin AV. Superior mesenteric artery syndrome: diagnosis and treatment strategies. *J Gastrointest Surg* 2009;13:287-92.
- 4) Egerton-Warburton D, Meek R, Mee MJ, Braitberg G. Antiemetic use for nausea and vomiting in adult emergency department patients: randomized controlled trial comparing ondansetron, metoclopramide, and placebo. *Ann Emerg Med* 2014;64:526-32.
- 5) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。コロナール錠、ブスコパン錠、プリンペラン錠、ナウゼリン錠、ボルタレンサボ、イリボー錠）。
- 6) Ohle R, O'Reilly F, O'Brien KK, Fahey T, Dimitrov BD. The Alvarado score for predicting acute appendicitis: a systematic review. *BMC Med* 2011;9:139.
- 7) Shea JA, Berlin JA, Escarce JJ, et al. Revised estimates of diagnostic test sensitivity and specificity in suspected biliary tract disease. *Arch Intern Med* 1994;154:2573-81.
- 8) 日本消化器病学会編. 胆石症診療ガイドライン2021（改訂第3版）. 南江堂；2021.
- 9) 日本泌尿器科学会, 日本尿路結石症学会, 日本泌尿器内視鏡・ロボティクス学会編. 尿路結石症診療ガイドライン 第3版. 2023.
- 10) Hollingsworth JM, Canales BK, Rogers MAM, et al. Alpha blockers for treatment of ureteric stones: systematic review and meta-analysis. *BMJ* 2016;355:i6112.
- 11) Peery AF, Shaukat A, Strate LL. AGA clinical practice update on medical management of colonic diverticulitis: expert review. *Gastroenterology* 2021;160:906-11.e1.
- 12) Fitzgibbons RJ Jr, Giobbie-Hurder A, Gibbs JO, et al. Watchful waiting vs repair of inguinal hernia in minimally symptomatic men: a randomized clinical trial. *JAMA* 2006;295:285-92.
- 13) Fukudo S, Ida M, Akiho H, Nakashima Y, Matsueda K. Effect of ramosetron on stool consistency in male patients with irritable bowel syndrome with diarrhea. *Clin Gastroenterol Hepatol* 2014;12:953-9.
- 14) Mol FMU, Maatman RC, De Joode LEGH, Van Eerten P, Scheltinga MR, Roumen R. Characteristics of 1116 consecutive patients diagnosed with anterior cutaneous nerve entrapment syndrome (ACNES). *Ann Surg* 2021;273:373-8.
- 15) Boelens OB, Scheltinga MR, Houterman S, Roumen RM. Randomized clinical trial of trigger point infiltration with lidocaine to diagnose anterior cutaneous nerve entrapment syndrome. *Br J Surg* 2013;100:217-21.
- 16) Shimizu K, Ito T, Irisawa A, et al. Evidence-based clinical practice guidelines for chronic pancreatitis 2021. *J Gastroenterol* 2022;57:709-24.
- 17) Fraquelli M, Casazza G, Conte D, Colli A. Non-steroid anti-inflammatory drugs for biliary colic. *Cochrane Database Syst Rev* 2016;9:CD006390.

2-14 下痢・便秘

- 1) 厚生労働省健康・生活衛生局感染症対策部感染症対策課. 抗微生物薬適正使用の手引き 第四版 医科・外来編. 2026.
- 2) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ロペミンカプセル、マグミット錠、新レシカルボン坐剤、テレミンソフト坐薬、モビコール配合内用剤、ラグノスNF経口ゼリー、アミティーザカプセル、リンゼス錠、グーフイス錠、ブルゼニド錠、イリボー錠、ポリフル錠、ピオフェルミンR錠、ミヤBM錠）。

- 3) McCarthy N, Giesecke J. Incidence of Guillain-Barré syndrome following infection with *Campylobacter jejuni*. *Am J Epidemiol* 2001;153:610-4.
- 4) Buchwald DS, Blaser MJ. A review of human salmonellosis: II. Duration of excretion following infection with nontyphi *Salmonella*. *Rev Infect Dis* 1984;6:345-56.
- 5) 日本消化管学会編. 便通異常症診療ガイドライン2023—慢性下痢症. 南江堂；2023.
- 6) Bonderup OK, Nielsen GL, Dall M, Pottegård A, Hallas J. Significant association between the use of different proton pump inhibitors and microscopic colitis: a nationwide Danish case-control study. *Aliment Pharmacol Ther* 2018;48:618-25.
- 7) van Rhee PF, Van de Vijver E, Fidler V. Faecal calprotectin for screening of patients with suspected inflammatory bowel disease: diagnostic meta-analysis. *BMJ* 2010;341:c3369.
- 8) 日本消化器病学会編. 機能的消化管疾患診療ガイドライン2020—過敏性腸症候群（IBS）（改訂第2版）. 南江堂；2020.
- 9) 日本消化管学会編. 便通異常症診療ガイドライン2023—慢性便秘症. 南江堂；2023.
- 10) Ford AC, Veldhuyzen van Zanten SJ, Rodgers CC, Talley NJ, Vakil NB, Moayyedi P. Diagnostic utility of alarm features for colorectal cancer: systematic review and meta-analysis. *Gut* 2008;57:1545-53.
- 11) Lee-Robichaud H, Thomas K, Morgan J, Nelson RL. Lactulose versus polyethylene glycol for chronic constipation. *Cochrane Database Syst Rev* 2010;(7):CD007570.
- 12) 厚生労働省保険局医療課長通知. 使用薬剤の薬価（薬価基準）の一部改正等について（保医発0821第1号）. 2018年8月21日.
- 13) Wakai E, Ikemura K, Sugimoto H, Iwamoto T, Okuda M. Risk factors for the development of hypermagnesemia in patients prescribed magnesium oxide: a retrospective cohort study. *J Pharm Health Care Sci* 2019;5:4.
- 14) Thabane M, Kottachchi DT, Marshall JK. Systematic review and meta-analysis: the incidence and prognosis of post-infectious irritable bowel syndrome. *Aliment Pharmacol Ther* 2007;26:535-44.
- 15) Power AM, Talley NJ, Ford AC. Association between constipation and colorectal cancer: systematic review and meta-analysis of observational studies. *Am J Gastroenterol* 2013;108:894-903.
- 16) Yamasaki M, Funakoshi S, Matsuda S, et al. Interaction of magnesium oxide with gastric acid secretion inhibitors in clinical pharmacotherapy. *Eur J Clin Pharmacol* 2014;70:921-4.
- 17) Ibuka H, Ishihara M, Suzuki A, et al. Antacid attenuates the laxative action of magnesium in cancer patients receiving opioid analgesic. *J Pharm Pharmacol* 2016;68:1214-21.

2-15 頻尿・尿失禁・排尿困難

- 1) 日本泌尿器科学会編. 男性下部尿路症状・前立腺肥大症診療ガイドライン. リッチヒルメディカル；2017.
- 2) 日本排尿機能学会, 日本泌尿器科学会編. 過活動膀胱診療ガイドライン [第3版]. リッチヒルメディカル；2022.
- 3) Sakakibara R, Kishi M, Tsuyusaki Y, et al. "Meningitis-retention syndrome": a review. *Neurourol Urodyn* 2013;32:19-23.
- 4) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ベタニス錠、ベオーバ錠、ベシケア錠、ハルナールD錠、ユリーフ錠、ザルティア錠、アボルブカプセル、ミニリンメルトOD錠、ケフラールカプセル、クラビット錠）.
- 5) Coupland CAC, Hill T, Denning T, Morriss R, Moore M, Hippisley-Cox J. Anticholinergic drug exposure and the risk of dementia: a nested case-control study. *JAMA Intern Med* 2019;179:1084-93.

- 6) Roehrborn CG, Siami P, Barkin J, et al. The effects of combination therapy with dutasteride and tamsulosin on clinical outcomes in men with symptomatic benign prostatic hyperplasia: 4-year results from the CombAT study. *Eur Urol* 2010;57:123-31.
- 7) Washino S, Ugata Y, Saito K, Miyagawa T. Calcium channel blockers are associated with nocturia in men aged 40 years or older. *J Clin Med* 2021;10:1603.
- 8) 日本排尿機能学会, 日本泌尿器科学会編. 夜間頻尿診療ガイドライン [第2版]. リッチヒルメディカル; 2020.
- 9) Pesonen JS, Vernooij RWM, Cartwright R, et al. The impact of nocturia on falls and fractures: a systematic review and meta-analysis. *J Urol* 2020;203:674-83.
- 10) Dumoulin C, Cacciari LP, Hay-Smith EJC. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Cochrane Database Syst Rev* 2018;10:CD005654.
- 11) Bent S, Nallamothu BK, Simel DL, Fihn SD, Saint S. Does this woman have an acute uncomplicated urinary tract infection? *JAMA* 2002;287:2701-10.
- 12) 日本感染症学会, 日本化学療法学会. JAID/JSC感染症治療ガイドライン2015—尿路感染症・男性性器感染症—. *日化療会誌* 2016;64:1-30.
- 13) Hooton TM, Vecchio M, Iroz A, et al. Effect of increased daily water intake in premenopausal women with recurrent urinary tract infections: a randomized clinical trial. *JAMA Intern Med* 2018;178:1509-15.
- 14) Williams G, Hahn D, Stephens JH, Craig JC, Hodson EM. Cranberries for preventing urinary tract infections. *Cochrane Database Syst Rev* 2023;4:CD001321.
- 15) Gágyor I, Bleidorn J, Kochen MM, Schmiemann G, Wegscheider K, Hummers-Pradier E. Ibuprofen versus fosfomycin for uncomplicated urinary tract infection in women: randomised controlled trial. *BMJ* 2015;351:h6544.
- 16) Vik I, Bollestad M, Grude N, et al. Ibuprofen versus pivmecillinam for uncomplicated urinary tract infection in women—a double-blind, randomized non-inferiority trial. *PLoS Med* 2018;15:e1002569.
- 17) Kronenberg A, Bütikofer L, Odutayo A, et al. Symptomatic treatment of uncomplicated lower urinary tract infections in the ambulatory setting: randomised, double blind trial. *BMJ* 2017;359:j4784.

2-16 性感染症

- 1) 厚生労働省健康・生活衛生局感染症対策部感染症対策課. 日本の性感染症の発生動向 (第97回厚生科学審議会感染症部会 参考資料2, 令和7年8月7日). 2025.
- 2) 厚生労働省エイズ動向委員会. 第167回エイズ動向委員会 委員長コメント: 令和7年 HIV感染者・エイズ患者の年間新規報告数 (確定値). 2026.
- 3) Workowski KA, Bachmann LH, Chan PA, et al. Sexually transmitted infections treatment guidelines, 2021. *MMWR Recomm Rep* 2021;70(4):1-187.
- 4) 内閣府男女共同参画局. 性犯罪・性暴力被害者のためのワンストップ支援センター一覧. 2026.
- 5) 日本性感染症学会. 性感染症 診断・治療 ガイドライン2020 梅毒 (2023年6月 一部改訂). 2023.
- 6) 厚生労働省保険局医療課長. 診療報酬の算定方法の一部改正に伴う実施上の留意事項について (令和8年3月5日保医発0305第6号). 別添1 D012 感染症免疫学的検査. 2026.
- 7) 日本性感染症学会. 性感染症 診断・治療 ガイドライン2016. *日性感染症会誌* 2016;27(1 Suppl).
- 8) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. ロセフィン静注用、ジスロマック錠、サワシリン、ステルイズ水、性懸濁筋注、ミノマイシン、ピブライマイシン錠、バルトレックス錠、アメナリーフ錠、ベセルナクリーム、ツルパダ配合錠).

- 9) Dombrowski JC, Wierzbicki MR, Newman LM, et al. Doxycycline versus azithromycin for the treatment of rectal chlamydia in men who have sex with men: a randomized controlled trial. *Clin Infect Dis* 2021;73:824-31.
- 10) 日本性感染症学会 梅毒委員会. ペンジルペニシリンベンザチン水和物製剤およびアモキシシリン水和物製剤 供給制限下における適正使用・適正発注へのご協力のお願い (2026年8月24日). 2026.
- 11) HIV感染症および血友病におけるチーム医療の構築と医療水準の向上を目指した研究班. 抗HIV治療ガイドライン (2026年3月版). 2026.
- 12) Ikeuchi K, Fukushima K, Tanaka M, Yajima K, Imamura A. Clinical efficacy and tolerability of 1.5 g/day oral amoxicillin therapy without probenecid for the treatment of syphilis. *Sex Transm Infect* 2022;98:173-7.
- 13) Ando N, Mizushima D, Omata K, et al. Combination of amoxicillin 3000 mg and probenecid versus 1500 mg amoxicillin monotherapy for treating syphilis in patients with human immunodeficiency virus: an open-label, randomized, controlled, non-inferiority trial. *Clin Infect Dis* 2023;77:779-87.
- 14) Nishijima T, Kawana K, Fukasawa I, et al. Effectiveness and tolerability of oral amoxicillin in pregnant women with active syphilis, Japan, 2010-2018. *Emerg Infect Dis* 2020;26:1192-200.
- 15) Geisler WM, Uniyal A, Lee JY, et al. Azithromycin versus doxycycline for urogenital Chlamydia trachomatis infection. *N Engl J Med* 2015;373:2512-21.
- 16) Kong FY, Tabrizi SN, Law M, et al. Azithromycin versus doxycycline for the treatment of genital chlamydia infection: a meta-analysis of randomized controlled trials. *Clin Infect Dis* 2014;59:193-205.

2-17 月経・更年期

- 1) 日本産科婦人科学会・日本産婦人科医会編. 産婦人科診療ガイドライン—婦人科外来編2023. 日本産科婦人科学会; 2023.
- 2) 厚生労働省. 要指導医薬品である緊急避妊薬の販売が可能な薬局・店舗販売業の店舗 一覧. 2026.
- 3) Shankar M, Lee CA, Sabin CA, Economides DL, Kadir RA. von Willebrand disease in women with menorrhagia: a systematic review. *BJOG* 2004;111:734-40.
- 4) Jespersen C, Lauritsen MP, Frokjaer VG, Schroll JB. Selective serotonin reuptake inhibitors for premenstrual syndrome and premenstrual dysphoric disorder. *Cochrane Database Syst Rev* 2024;8:CD001396.
- 5) Marjoribanks J, Ayeleke RO, Farquhar C, Proctor M. Nonsteroidal anti-inflammatory drugs for dysmenorrhoea. *Cochrane Database Syst Rev* 2015;(7):CD001751.
- 6) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。ブルフェン錠、ナイキサン錠、ポンタールカプセル、ロキソニン錠、ジュリナ錠、ツムラ加味逍遙散、ツムラ当帰芍薬散、ツムラ桂枝茯苓丸)。
- 7) Bastian LA, Smith CM, Nanda K. Is this woman perimenopausal? *JAMA* 2003;289:895-902.
- 8) MacLennan AH, Broadbent JL, Lester S, Moore V. Oral oestrogen and combined oestrogen/progestogen therapy versus placebo for hot flushes. *Cochrane Database Syst Rev* 2004;(4):CD002978.
- 9) Takamatsu K, Ogawa M, Higuchi T, Takeda T, Hayashi K, Mizunuma H. Effects of kamishoyosan, a traditional Japanese medicine, on menopausal symptoms: a randomized, placebo-controlled, double-blind clinical trial. *Evid Based Complement Alternat Med* 2020;2020:9285317.
- 10) Clarke MA, Long BJ, Del Mar Morillo A, Arbyn M, Bakkum-Gamez JN, Wentzensen N. Association of endometrial cancer risk with postmenopausal bleeding in women: a systematic review and meta-analysis. *JAMA Intern Med* 2018;178:1210-22.

- 11) Perrotta C, Aznar M, Mejia R, Albert X, Ng CW. Oestrogens for preventing recurrent urinary tract infection in postmenopausal women. *Cochrane Database Syst Rev* 2008;(2):CD005131.
- 12) 日本医療研究開発機構 障害者対策総合研究開発事業「摂食障害の治療支援ネットワークの指針と簡易治療プログラムの開発」(研究開発代表者 安藤哲也, 分担 須藤信行). 神経性やせ症(AN) 初期診療の手引き 第1版. 2019.
- 13) Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321-33.
- 14) Manson JE, Aragaki AK, Rossouw JE, et al. Menopausal hormone therapy and long-term all-cause and cause-specific mortality: the Women's Health Initiative randomized trials. *JAMA* 2017;318:927-38.
- 15) Manson JE, Chlebowski RT, Stefanick ML, et al. Menopausal hormone therapy and health outcomes during the intervention and extended poststopping phases of the Women's Health Initiative randomized trials. *JAMA* 2013;310:1353-68.

2-18 腰痛・関節痛

- 1) Henschke N, Maher CG, Refshauge KM, et al. Prevalence of and screening for serious spinal pathology in patients presenting to primary care settings with acute low back pain. *Arthritis Rheum* 2009;60:3072-80.
- 2) 日本整形外科学会, 日本腰痛学会監修. 腰痛診療ガイドライン2019 (改訂第2版). 南江堂; 2019.
- 3) Downie A, Williams CM, Henschke N, et al. Red flags to screen for malignancy and fracture in patients with low back pain: systematic review. *BMJ* 2013;347:f7095.
- 4) Langdon J, Way A, Heaton S, Bernard J, Molloy S. Vertebral compression fractures--new clinical signs to aid diagnosis. *Ann R Coll Surg Engl* 2010;92:163-6.
- 5) Geusens P, Kendler DL, Fahrleitner-Pammer A, López-Romero P, Marin F. Distribution of prevalent and incident vertebral fractures and their association with bone mineral density in postmenopausal women in the teriparatide versus risedronate VERO clinical trial. *Calcif Tissue Int* 2020;106:646-54.
- 6) Chou R, Fu R, Carrino JA, Deyo RA. Imaging strategies for low-back pain: systematic review and meta-analysis. *Lancet* 2009;373:463-72.
- 7) Brinjikji W, Luetmer PH, Comstock B, et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol* 2015;36:811-6.
- 8) da C Menezes Costa L, Maher CG, Hancock MJ, McAuley JH, Herbert RD, Costa LO. The prognosis of acute and persistent low-back pain: a meta-analysis. *CMAJ* 2012;184:E613-24.
- 9) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。ロキソニン錠・テープ、セレコックス錠、サインバルタカプセル、カロナール錠、オパルモン錠、プレタール錠)。
- 10) van der Gaag WH, Roelofs PD, Enthoven WT, van Tulder MW, Koes BW. Non-steroidal anti-inflammatory drugs for acute low back pain. *Cochrane Database Syst Rev* 2020;4:CD013581.
- 11) Williams CM, Maher CG, Latimer J, et al. Efficacy of paracetamol for acute low-back pain: a double-blind, randomised controlled trial. *Lancet* 2014;384:1586-96.
- 12) Sieper J, van der Heijde D, Landewé R, et al. New criteria for inflammatory back pain in patients with chronic back pain: a real patient exercise by experts from the Assessment of SpondyloArthritis international Society (ASAS). *Ann Rheum Dis* 2009;68:784-8.
- 13) Poddubnyy D, van Tubergen A, Landewé R, Sieper J, van der Heijde D. Development of an ASAS-endorsed recommendation for the early referral of patients with a suspicion of axial spondyloarthritis. *Ann Rheum Dis* 2015;74:1483-7.

- 14) Steffens D, Maher CG, Pereira LS, et al. Prevention of low back pain: a systematic review and meta-analysis. *JAMA Intern Med* 2016;176:199-208.
- 15) van der Windt DA, Simons E, Riphagen II, et al. Physical examination for lumbar radiculopathy due to disc herniation in patients with low-back pain. *Cochrane Database Syst Rev* 2010; (2):CD007431.
- 16) Mathieson S, Maher CG, McLachlan AJ, et al. Trial of pregabalin for acute and chronic sciatica. *N Engl J Med* 2017;376:1111-20.
- 17) Peul WC, van Houwelingen HC, van den Hout WB, et al. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med* 2007;356:2245-56.
- 18) Suri P, Rainville J, Kalichman L, Katz JN. Does this older adult with lower extremity pain have the clinical syndrome of lumbar spinal stenosis? *JAMA* 2010;304:2628-36.
- 19) 日本循環器学会, 日本血管外科学会. 2022年改訂版 末梢動脈疾患ガイドライン. 2022.
- 20) 日本整形外科学会, 日本脊椎脊髄病学会監修. 腰部脊柱管狭窄症診療ガイドライン2021 (改訂第2版). 南江堂; 2021.
- 21) Aletaha D, Neogi T, Silman AJ, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum* 2010;62:2569-81.
- 22) Nishimura K, Sugiyama D, Kogata Y, et al. Meta-analysis: diagnostic accuracy of anti-cyclic citrullinated peptide antibody and rheumatoid factor for rheumatoid arthritis. *Ann Intern Med* 2007;146:797-808.
- 23) Combe B, Landewe R, Daien CI, et al. 2016 update of the EULAR recommendations for the management of early arthritis. *Ann Rheum Dis* 2017;76:948-59.
- 24) Ribbens C, Martin y Porras M, Franchimont N, et al. Increased matrix metalloproteinase-3 serum levels in rheumatic diseases: relationship with synovitis and steroid treatment. *Ann Rheum Dis* 2002;61:161-6.
- 25) Derry S, Conaghan P, Da Silva JA, Wiffen PJ, Moore RA. Topical NSAIDs for chronic musculoskeletal pain in adults. *Cochrane Database Syst Rev* 2016;4:CD007400.
- 26) Nakamura J, Oinuma K, Ohtori S, et al. Distribution of hip pain in osteoarthritis patients secondary to developmental dysplasia of the hip. *Mod Rheumatol* 2013;23:119-24.
- 27) Taylor W, Gladman D, Helliwell P, et al. Classification criteria for psoriatic arthritis: development of new criteria from a large international study. *Arthritis Rheum* 2006;54:2665-73.
- 28) Hannu T. Reactive arthritis. *Best Pract Res Clin Rheumatol* 2011;25:347-57.
- 29) Yamamoto T, Hiraiwa T, Tobita R, et al. Characteristics of Japanese patients with pustulotic arthro-osteitis associated with palmoplantar pustulosis: a multicenter study. *Int J Dermatol* 2020;59:441-4.
- 30) Hand C, Clipsham K, Rees JL, Carr AJ. Long-term outcome of frozen shoulder. *J Shoulder Elbow Surg* 2008;17:231-6.
- 31) Hermans J, Luime JJ, Meuffels DE, Reijman M, Simel DL, Bierma-Zeinstra SM. Does this patient with shoulder pain have rotator cuff disease? The Rational Clinical Examination systematic review. *JAMA* 2013;310:837-47.
- 32) Minagawa H, Yamamoto N, Abe H, et al. Prevalence of symptomatic and asymptomatic rotator cuff tears in the general population: from mass-screening in one village. *J Orthop* 2013;10:8-12.
- 33) Jones CMP, Day RO, Koes BW, et al. Opioid analgesia for acute low back pain and neck pain (the OPAL trial): a randomised placebo-controlled trial. *Lancet* 2023;402:304-12.
- 34) Pereira TV, Jüni P, Saadat P, et al. Viscosupplementation for knee osteoarthritis: systematic review and meta-analysis. *BMJ* 2022;378:e069722.

- 35) 日本整形外科学会監修，日本整形外科学会診療ガイドライン委員会，変形性膝関節症診療ガイドライン策定委員会編．変形性膝関節症診療ガイドライン2023．南江堂；2023．

2-19 発疹・かゆみ

- 1) 国立健康危機管理研究機構 感染症情報提供サイト．麻疹．2026年10月に閲覧．
- 2) 学校保健安全法施行規則（昭和33年文部省令第18号）第19条．e-Gov法令検索．2026年10月に閲覧．
- 3) 国立健康危機管理研究機構 感染症情報提供サイト．手足口病（詳細版）．2014年10月17日改訂．2026年10月に閲覧．
- 4) 蕁麻疹診療ガイドライン策定委員会．蕁麻疹診療ガイドライン2026（第4版）．日皮会誌 2026;136:375-493．
- 5) 田中稔彦，亀好良一，秀 道広．広島大学皮膚科外来での蕁麻疹の病型別患者数．アレルギー 2006;55:134-9．
- 6) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認．レミッチ、アレグラ錠、ピラノア錠、バルトレックス錠、ファムビル錠、アメナリーフ錠、リリカ、タリージェ錠、ストロメクトール錠、スミスリンローション、カロナール錠、リンデロン-V軟膏、ヒルドイドローション、モーラステープ）．
- 7) 日本小児アレルギー学会．食物アレルギー診療ガイドライン2021．協和企画；2021．
- 8) Morita E, Matsuo H, Chinuki Y, Takahashi H, Dahlström J, Tanaka A. Food-dependent exercise-induced anaphylaxis: importance of omega-5 gliadin and HMW-glutenin as causative antigens for wheat-dependent exercise-induced anaphylaxis. *Allergol Int* 2009;58:493-8.
- 9) 帯状疱疹診療ガイドライン策定委員会．帯状疱疹診療ガイドライン2025．日皮会誌 2025;135:527-56．
- 10) Takao Y, Miyazaki Y, Okeda M, et al. Incidences of herpes zoster and postherpetic neuralgia in Japanese adults aged 50 years and older from a community-based prospective cohort study: the SHEZ study. *J Epidemiol* 2015;25:617-25．
- 11) Blakely KM, Drucker AM, Rosen CF. Drug-induced photosensitivity—an update: culprit drugs, prevention and management. *Drug Saf* 2019;42:827-47．
- 12) Anderson HJ, Lee JB. A review of fixed drug eruption with a special focus on generalized bullous fixed drug eruption. *Medicina (Kaunas)* 2021;57:925．
- 13) 佐藤貴浩，横関博雄，室田浩之，ほか．皮膚瘙痒症診療ガイドライン2020．日皮会誌 2020;130:1589-606．
- 14) 日本皮膚科学会疥癬診療ガイドライン策定委員会．疥癬診療ガイドライン（第3版）．日皮会誌 2015;125:2023-48．
- 15) 日本皮膚科学会・日本医真菌学会合同皮膚真菌症診療ガイドライン策定委員会．皮膚真菌症診療ガイドライン2025．日皮会誌 2025;135:2511-66．
- 16) 佐伯秀久，大矢幸弘，荒川浩一，ほか．アトピー性皮膚炎診療ガイドライン2024．日皮会誌 2024;134:2741-843．
- 17) Long CC, Finlay AY. The finger-tip unit—a new practical measure. *Clin Exp Dermatol* 1991;16:444-7．
- 18) Mihalache A, Yu P, Tao BK, Xie JS, Margolin EA, Micieli JA. Hutchinson's sign in herpes zoster ophthalmicus: a systematic review and meta-analysis. *Ophthalmology* 2026;133:411-22．
- 19) Chen N, Li Q, Yang J, Zhou M, Zhou D, He L. Antiviral treatment for preventing postherpetic neuralgia. *Cochrane Database Syst Rev* 2014;(2):CD006866．
- 20) Jiang X, Li Y, Chen N, Zhou M, He L. Corticosteroids for preventing postherpetic neuralgia. *Cochrane Database Syst Rev* 2023;12:CD005582．

2-20 不眠・不安・抑うつ

- 1) Arroll B, Khin N, Kerse N. Screening for depression in primary care with two verbally asked questions: cross sectional study. *BMJ* 2003;327:1144-6．

- 2) Levis B, Benedetti A, Thombs BD; DEPRESSion Screening Data (DEPRESSD) Collaboration. Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. *BMJ* 2019;365:l1476.
- 3) Dazzi T, Gribble R, Wessely S, Fear NT. Does asking about suicide and related behaviours induce suicidal ideation? What is the evidence? *Psychol Med* 2014;44:3361-3.
- 4) Nagappa M, Liao P, Wong J, et al. Validation of the STOP-Bang questionnaire as a screening tool for obstructive sleep apnea among different populations: a systematic review and meta-analysis. *PLoS One* 2015;10:e0143697.
- 5) 厚生労働省 健康づくりのための睡眠指針の改訂に関する検討会. 健康づくりのための睡眠ガイド2023. 2024年2月.
- 6) 厚生労働省. 健康に配慮した飲酒に関するガイドライン. 2024年2月.
- 7) Trauer JM, Qian MY, Doyle JS, Rajaratnam SM, Cunnington D. Cognitive behavioral therapy for chronic insomnia: a systematic review and meta-analysis. *Ann Intern Med* 2015;163:191-204.
- 8) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。デエビゴ錠、クービピック錠、ジェイゾロフト錠、レクサプロ錠、リフレックス錠）.
- 9) 厚生労働科学研究・障害者対策総合研究事業「睡眠薬の適正使用及び減量・中止のための診療ガイドラインに関する研究班」, 日本睡眠学会・睡眠薬使用ガイドライン作成ワーキンググループ編. 睡眠薬の適正な使用と休薬のための診療ガイドライン. 2013.
- 10) Tannenbaum C, Martin P, Tamblyn R, Benedetti A, Ahmed S. Reduction of inappropriate benzodiazepine prescriptions among older adults through direct patient education: the EMPOWER cluster randomized trial. *JAMA Intern Med* 2014;174:890-8.
- 11) Takaesu Y, Utsumi T, Okajima I, et al. Psychosocial intervention for discontinuing benzodiazepine hypnotics in patients with chronic insomnia: a systematic review and meta-analysis. *Sleep Med Rev* 2019;48:101214.
- 12) 日本うつ病学会 うつ病診療ガイドライン作成ワーキンググループ. 日本うつ病学会 うつ病診療ガイドライン2025. 2025.
- 13) Stone M, Laughren T, Jones ML, et al. Risk of suicidality in clinical trials of antidepressants in adults: analysis of proprietary data submitted to US Food and Drug Administration. *BMJ* 2009;339:b2880.
- 14) Cipriani A, Furukawa TA, Salanti G, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. *Lancet* 2018;391:1357-66.
- 15) Mitchell AJ, Vaze A, Rao S. Clinical diagnosis of depression in primary care: a meta-analysis. *Lancet* 2009;374:609-19.
- 16) Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606-13.
- 17) Callahan M. Hypoxic hazards of traditional paper bag rebreathing in hyperventilating patients. *Ann Emerg Med* 1989;18:622-8.
- 18) Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166:1092-7.
- 19) Kyle SD, Siriwardena AN, Espie CA, et al. Clinical and cost-effectiveness of nurse-delivered sleep restriction therapy for insomnia in primary care (HABIT): a pragmatic, superiority, open-label, randomised controlled trial. *Lancet* 2023;402:975-87.
- 20) Fournier JC, DeRubeis RJ, Hollon SD, et al. Antidepressant drug effects and depression severity: a patient-level meta-analysis. *JAMA* 2010;303:47-53.

2-21 睡眠時無呼吸

- 1) Benjafield AV, Ayas NT, Eastwood PR, et al. Estimation of the global prevalence and burden of obstructive sleep apnoea: a literature-based analysis. *Lancet Respir Med* 2019;7:687-98.
- 2) 日本循環器学会, ほか. 2023年改訂版 循環器領域における睡眠呼吸障害の診断・治療に関するガイドライン. 2023.
- 3) Myers KA, Mrkobrada M, Simel DL. Does this patient have obstructive sleep apnea?: The Rational Clinical Examination systematic review. *JAMA* 2013;310:731-41.
- 4) 日本呼吸器学会, 厚生労働科学研究費補助金難治性疾患政策研究事業「難治性呼吸器疾患・肺高血圧症に関する調査研究」班 監修. 睡眠時無呼吸症候群 (SAS) の診療ガイドライン2020. 南江堂; 2020.
- 5) Mokhlesi B, Masa JF, Brozek JL, et al. Evaluation and management of obesity hypoventilation syndrome. An official American Thoracic Society clinical practice guideline. *Am J Respir Crit Care Med* 2019;200:e6-e24.
- 6) 厚生労働省保険局医療課長. 診療報酬の算定方法の一部改正に伴う実施上の留意事項について (令和8年3月5日保医発0305第6号). 別添1 C107-2 在宅持続陽圧呼吸療法指導管理料, D237 終夜睡眠ポリグラフイー. 2026.
- 7) McEvoy RD, Antic NA, Heeley E, et al. CPAP for prevention of cardiovascular events in obstructive sleep apnea. *N Engl J Med* 2016;375:919-31.
- 8) Andrade RG, Piccin VS, Nascimento JA, Viana FM, Genta PR, Lorenzi-Filho G. Impact of the type of mask on the effectiveness of and adherence to continuous positive airway pressure treatment for obstructive sleep apnea. *J Bras Pneumol* 2014;40:658-68.
- 9) Askland K, Wright L, Wozniak DR, Emmanuel T, Caston J, Smith I. Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev* 2020;4:CD007736.
- 10) モディオダール錠100 mgの電子添文 (KEGG MEDICUSで2026年10月に確認) .
- 11) Foster GD, Borradaile KE, Sanders MH, et al. A randomized study on the effect of weight loss on obstructive sleep apnea among obese patients with type 2 diabetes: the Sleep AHEAD study. *Arch Intern Med* 2009;169:1619-26.
- 12) Drager LF, Brunoni AR, Jenner R, Lorenzi-Filho G, Benseñor IM, Lotufo PA. Effects of CPAP on body weight in patients with obstructive sleep apnoea: a meta-analysis of randomised trials. *Thorax* 2015;70:258-64.
- 13) Malhotra A, Grunstein RR, Fietze I, et al. Tirzepatide for the treatment of obstructive sleep apnea and obesity. *N Engl J Med* 2024;391:1193-205.
- 14) Srijithesh PR, Aghoram R, Goel A, Dhanya J. Positional therapy for obstructive sleep apnoea. *Cochrane Database Syst Rev* 2019;5:CD010990.
- 15) 道路交通法 (昭和35年法律第105号) 第66条, 第89条, 第101条, 第117条の4. 道路交通法施行令 (昭和35年政令第270号) 第33条の2の3.
- 16) Tregear S, Reston J, Schoelles K, Phillips B. Obstructive sleep apnea and risk of motor vehicle crash: systematic review and meta-analysis. *J Clin Sleep Med* 2009;5:573-81.
- 17) Tregear S, Reston J, Schoelles K, Phillips B. Continuous positive airway pressure reduces risk of motor vehicle crash among drivers with obstructive sleep apnea: systematic review and meta-analysis. *Sleep* 2010;33:1373-80.
- 18) Martínez-García MA, Capote F, Campos-Rodríguez F, et al. Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *JAMA* 2013;310:2407-15.
- 19) Cowie MR, Woehrle H, Wegscheider K, et al. Adaptive servo-ventilation for central sleep apnea in systolic heart failure. *N Engl J Med* 2015;373:1095-105.

- 20) Sánchez-de-la-Torre M, Gracia-Lavedan E, Benitez ID, et al. Adherence to CPAP treatment and the risk of recurrent cardiovascular events: a meta-analysis. *JAMA* 2023;330:1255-65.

2-22 検査で説明のつかない症状

- 1) Kroenke K, Mangelsdorff AD. Common symptoms in ambulatory care: incidence, evaluation, therapy, and outcome. *Am J Med* 1989;86:262-6.
- 2) Haller H, Cramer H, Lauche R, Dobos G. Somatoform disorders and medically unexplained symptoms in primary care. *Dtsch Arztebl Int* 2015;112:279-87.
- 3) Eikelboom EM, Tak LM, Roest AM, Rosmalen JGM. A systematic review and meta-analysis of the percentage of revised diagnoses in functional somatic symptoms. *J Psychosom Res* 2016;88:60-7.
- 4) Roenneberg C, Sattel H, Schaefer R, Henningsen P, Hausteiner-Wiehle C. Functional somatic symptoms. *Dtsch Arztebl Int* 2019;116:553-60.
- 5) Kroenke K, Spitzer RL, Williams JB, et al. Physical symptoms in primary care. Predictors of psychiatric disorders and functional impairment. *Arch Fam Med* 1994;3:774-9.
- 6) Wolfe F, Clauw DJ, Fitzcharles MA, et al. 2016 Revisions to the 2010/2011 fibromyalgia diagnostic criteria. *Semin Arthritis Rheum* 2016;46:319-29.
- 7) Bidonde J, Busch AJ, Schachter CL, et al. Aerobic exercise training for adults with fibromyalgia. *Cochrane Database Syst Rev* 2017;6:CD012700.
- 8) リリカ、サインバルタの電子添文（2026年10月に確認）。
- 9) McWhirter L, Stone J, Sandercock P, Whiteley W. Hoover's sign for the diagnosis of functional weakness: a prospective unblinded cohort study in patients with suspected stroke. *J Psychosom Res* 2011;71:384-6.
- 10) Löwe B, Levenson J, Depping M, et al. Somatic symptom disorder: a scoping review on the empirical evidence of a new diagnosis. *Psychol Med* 2022;52:632-48.
- 11) Gierk B, Kohlmann S, Kroenke K, et al. The somatic symptom scale-8 (SSS-8): a brief measure of somatic symptom burden. *JAMA Intern Med* 2014;174:399-407.
- 12) olde Hartman TC, Borghuis MS, Lucassen PL, van de Laar FA, Speckens AE, van Weel C. Medically unexplained symptoms, somatisation disorder and hypochondriasis: course and prognosis. A systematic review. *J Psychosom Res* 2009;66:363-77.
- 13) Kingdon C, Lowe A, Shepherd C, Nacul L. What primary care practitioners need to know about the new NICE guideline for myalgic encephalomyelitis/chronic fatigue syndrome in adults. *Healthcare (Basel)* 2022;10:2438.
- 14) Thomas KB. General practice consultations: is there any point in being positive? *Br Med J (Clin Res Ed)* 1987;294:1200-2.
- 15) Petrie KJ, Müller JT, Schirmbeck F, et al. Effect of providing information about normal test results on patients' reassurance: randomised controlled trial. *BMJ* 2007;334:352.
- 16) Hoedeman R, Blankenstein AH, van der Feltz-Cornelis CM, Krol B, Stewart R, Groothoff JW. Consultation letters for medically unexplained physical symptoms in primary care. *Cochrane Database Syst Rev* 2010;(12):CD006524.
- 17) Smith GR Jr, Monson RA, Ray DC. Psychiatric consultation in somatization disorder. A randomized controlled study. *N Engl J Med* 1986;314:1407-13.
- 18) Ring A, Dowrick C, Humphris G, Salmon P. Do patients with unexplained physical symptoms pressurise general practitioners for somatic treatment? A qualitative study. *BMJ* 2004;328:1057.
- 19) Kleinstäuber M, Witthöft M, Steffanowski A, van Marwijk H, Hiller W, Lambert MJ. Pharmacological interventions for somatoform disorders in adults. *Cochrane Database Syst Rev* 2014;(11):CD010628.

- 20) van Dessel N, den Boeft M, van der Wouden JC, et al. Non-pharmacological interventions for somatoform disorders and medically unexplained physical symptoms (MUPS) in adults. Cochrane Database Syst Rev 2014;(11):CD011142.

2-23 もの忘れ

- 1) Holsinger T, Deveau J, Boustani M, Williams JW Jr. Does this patient have dementia? JAMA 2007;297:2391-404.
- 2) Satyadev N, Tipton PW, Martens Y, et al. Improving early recognition of treatment-responsive causes of rapidly progressive dementia: the STAM3P score. Ann Neurol 2024;95:237-48.
- 3) Creavin ST, Wisniewski S, Noel-Storr AH, et al. Mini-Mental State Examination (MMSE) for the detection of dementia in clinically unevaluated people aged 65 and over in community and primary care populations. Cochrane Database Syst Rev 2016;(1):CD011145.
- 4) 日本神経学会監修, 「認知症疾患診療ガイドライン」作成委員会編. 認知症疾患診療ガイドライン 2017. 医学書院; 2017.
- 5) Fujiwara Y, Suzuki H, Yasunaga M, et al. Brief screening tool for mild cognitive impairment in older Japanese: validation of the Japanese version of the Montreal Cognitive Assessment. Geriatr Gerontol Int 2010;10:225-32.
- 6) Seitz DP, Chan CC, Newton HT, et al. Mini-Cog for the detection of dementia within a primary care setting. Cochrane Database Syst Rev 2021;7:CD011145.
- 7) Ngandu T, Lehtisalo J, Solomon A, et al. A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. Lancet 2015;385:2255-63.
- 8) Clarfield AM. The decreasing prevalence of reversible dementias: an updated meta-analysis. Arch Intern Med 2003;163:2219-29.
- 9) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。アリセプト、メマリー、イクセロンパッチ、レミニール、レケンピ点滴静注、ケサンラ点滴静注)。
- 10) Birks JS, Harvey RJ. Donepezil for dementia due to Alzheimer's disease. Cochrane Database Syst Rev 2018;6:CD001190.
- 11) Gill SS, Anderson GM, Fischer HD, et al. Syncope and its consequences in patients with dementia receiving cholinesterase inhibitors: a population-based cohort study. Arch Intern Med 2009;169:867-73.
- 12) Woods B, Rai HK, Elliott E, Aguirre E, Orrell M, Spector A. Cognitive stimulation to improve cognitive functioning in people with dementia. Cochrane Database Syst Rev 2023;1:CD005562.
- 13) van Dyck CH, Swanson CJ, Aisen P, et al. Lecanemab in early Alzheimer's disease. N Engl J Med 2023;388:9-21.
- 14) 厚生労働省. 最適使用推進ガイドライン レカネマブ (遺伝子組換え) (令和5年12月)、ドナネマブ (遺伝子組換え) (令和6年11月、令和7年8月改訂)。
- 15) Zimmer JA, Ardayfio P, Wang H, et al. Amyloid-related imaging abnormalities with donanemab in early symptomatic Alzheimer disease: secondary analysis of the TRAILBLAZER-ALZ and ALZ 2 randomized clinical trials. JAMA Neurol 2025;82:461-9.
- 16) 道路交通法 (昭和35年法律第105号) 第101条の4、第101条の6、第102条、第103条。
- 17) 警察庁. 認知機能検査について (ウェブサイト、2026年10月閲覧)。
- 18) 高齢者虐待の防止、高齢者の養護者に対する支援等に関する法律 (平成17年法律第124号) 第7条。
- 19) Howard R, McShane R, Lindsay J, et al. Donepezil and memantine for moderate-to-severe Alzheimer's disease. N Engl J Med 2012;366:893-903.

- 20) Howard R, McShane R, Lindesay J, et al. Nursing home placement in the Donepezil and Memantine in Moderate to Severe Alzheimer's Disease (DOMINO-AD) trial: secondary and post-hoc analyses. *Lancet Neurol* 2015;14:1171-81.
- 21) Andrews JS, Desai U, Kirson NY, Zichlin ML, Ball DE, Matthews BR. Disease severity and minimal clinically important differences in clinical outcome assessments for Alzheimer's disease clinical trials. *Alzheimers Dement (N Y)* 2019;5:354-63.
- 22) Ebell MH, Barry HC, Baduni K, Grasso G. Clinically important benefits and harms of monoclonal antibodies targeting amyloid for the treatment of Alzheimer disease: a systematic review and meta-analysis. *Ann Fam Med* 2024;22:50-62.

2-24 高齢者の外来

- 1) Kojima G, Iliffe S, Taniguchi Y, Shimada H, Rakugi H, Walters K. Prevalence of frailty in Japan: a systematic review and meta-analysis. *J Epidemiol* 2017;27:347-53.
- 2) Briggs R, McDonough A, Ellis G, Bennett K, O'Neill D, Robinson D. Comprehensive geriatric assessment for community-dwelling, high-risk, frail, older people. *Cochrane Database Syst Rev* 2022;5:CD012705.
- 3) Bagai A, Thavendiranathan P, Detsky AS. Does this patient have hearing impairment? *JAMA* 2006;295:416-28.
- 4) Lin FR, Pike JR, Albert MS, et al. Hearing intervention versus health education control to reduce cognitive decline in older adults with hearing loss in the USA (ACHIEVE): a multicentre, randomised controlled trial. *Lancet* 2023;402:786-97.
- 5) Harwood RH, Foss AJ, Osborn F, Gregson RM, Zaman A, Masud T. Falls and health status in elderly women following first eye cataract surgery: a randomised controlled trial. *Br J Ophthalmol* 2005;89:53-9.
- 6) Tanaka T, Takahashi K, Hirano H, et al. Oral frailty as a risk factor for physical frailty and mortality in community-dwelling elderly. *J Gerontol A Biol Sci Med Sci* 2018;73:1661-7.
- 7) Satake S, Arai H. The revised Japanese version of the Cardiovascular Health Study criteria (revised J-CHS criteria). *Geriatr Gerontol Int* 2020;20:992-3.
- 8) Chen LK, Woo J, Assantachai P, et al. Asian Working Group for Sarcopenia: 2019 consensus update on sarcopenia diagnosis and treatment. *J Am Med Dir Assoc* 2020;21:300-7.e2.
- 9) Chen LK, Hsiao FY, Akishita M, et al. A focus shift from sarcopenia to muscle health in the Asian Working Group for Sarcopenia 2025 consensus update. *Nat Aging* 2025;5:2164-75.
- 10) Tanaka T, Takahashi K, Akishita M, Tsuji T, Iijima K. "Yubi-wakka" (finger-ring) test: a practical self-screening method for sarcopenia, and a predictor of disability and mortality among Japanese community-dwelling older adults. *Geriatr Gerontol Int* 2018;18:224-32.
- 11) Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: the LIFE study randomized clinical trial. *JAMA* 2014;311:2387-96.
- 12) 厚生労働省. 「日本人の食事摂取基準（2025年版）」策定検討会報告書. 2024.
- 13) Milne AC, Potter J, Vivanti A, Avenell A. Protein and energy supplementation in elderly people at risk from malnutrition. *Cochrane Database Syst Rev* 2009;(2):CD003288.
- 14) Montero-Odasso M, van der Velde N, Martin FC, et al. World guidelines for falls prevention and management for older adults: a global initiative. *Age Ageing* 2022;51:afac205.
- 15) Ganz DA, Bao Y, Shekelle PG, Rubenstein LZ. Will my patient fall? *JAMA* 2007;297:77-86.
- 16) Sherrington C, Fairhall NJ, Wallbank GK, et al. Exercise for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2019;1:CD012424.

- 17) Clemson L, Stark S, Pighills AC, et al. Environmental interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2023;3:CD013258.
- 18) Studenski S, Perera S, Patel K, et al. Gait speed and survival in older adults. *JAMA* 2011;305:50-8.
- 19) 介護保険法施行令 第2条（特定疾病）。
- 20) 厚生省老人保健福祉局企画課長通知. 指定居宅サービスに要する費用の額の算定に関する基準（訪問通所サービス、居宅療養管理指導及び福祉用具貸与に係る部分）及び指定居宅介護支援に要する費用の額の算定に関する基準の制定に伴う実施上の留意事項について（平成12年3月1日 老企第36号）。
- 21) 警察庁. 令和5年版 警察白書 第5章 安全かつ快適な交通の確保. 2023.
- 22) Chihuri S, Mielenz TJ, DiMaggio CJ, et al. Driving cessation and health outcomes in older adults. *J Am Geriatr Soc* 2016;64:332-41.
- 23) Hirai H, Ichikawa M, Kondo N, Kondo K. The risk of functional limitations after driving cessation among older Japanese adults: the JAGES cohort study. *J Epidemiol* 2020;30:332-7.
- 24) Yourman LC, Cenzer IS, Boscardin WJ, et al. Evaluation of time to benefit of statins for the primary prevention of cardiovascular events in adults aged 50 to 75 years: a meta-analysis. *JAMA Intern Med* 2021;181:179-85.
- 25) Chen T, Shao F, Chen K, et al. Time to clinical benefit of intensive blood pressure lowering in patients 60 years and older with hypertension: a secondary analysis of randomized clinical trials. *JAMA Intern Med* 2022;182:660-7.
- 26) Deardorff WJ, Cenzer I, Nguyen B, Lee SJ. Time to benefit of bisphosphonate therapy for the prevention of fractures among postmenopausal women with osteoporosis: a meta-analysis of randomized clinical trials. *JAMA Intern Med* 2022;182:33-41.
- 27) Sheppard JP, Burt J, Lown M, et al. Effect of antihypertensive medication reduction vs usual care on short-term blood pressure control in patients with hypertension aged 80 years and older: the OPTIMISE randomized clinical trial. *JAMA* 2020;323:2039-51.
- 28) Sheppard JP, Temple E, Wang A, et al. Effect of antihypertensive deprescribing on hospitalisation and mortality: long-term follow-up of the OPTiMISE randomised controlled trial. *Lancet Healthy Longev* 2024;5:e563-73.

2-25 外来の小さな訴え

- 1) Garrison SR, Dormuth CR, Morrow RL, Carney GA, Khan KM. Nocturnal leg cramps and prescription use that precedes them: a sequence symmetry analysis. *Arch Intern Med* 2012;172:120-6.
- 2) Hallegraeff JM, van der Schans CP, de Ruiter R, de Greef MH. Stretching before sleep reduces the frequency and severity of nocturnal leg cramps in older adults: a randomised trial. *J Physiother* 2012;58:17-22.
- 3) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ツムラ芍薬甘草湯エキス顆粒、コントミン糖衣錠、ギャバロン錠、プリンペラン錠、シグマート錠、オルテクサー口腔用軟膏、デキサメタゾン口腔用軟膏、アフタッチ口腔用貼付剤、フロリッドゲル経口用、サリグレンカプセル、サラジェン錠、フェロミア錠、ビ・シフロール錠、ニュープロバッチ、レグナイト錠、アロチノロール塩酸塩錠、インデルール錠、プリミドン錠）。
- 4) 厚生労働省. 重篤副作用疾患別対応マニュアル 偽アルドステロン症. 2006（2022年2月改定）。
- 5) Garrison SR, Korownyk CS, Kolber MR, et al. Magnesium for skeletal muscle cramps. *Cochrane Database Syst Rev* 2020;9:CD009402.
- 6) El-Tawil S, Al Musa T, Valli H, et al. Quinine for muscle cramps. *Cochrane Database Syst Rev* 2015;(4):CD005044.
- 7) Chang FY, Lu CL. Hiccup: mystery, nature and treatment. *J Neurogastroenterol Motil* 2012;18:123-30.

- 8) Zhang C, Zhang R, Zhang S, Xu M, Zhang S. Baclofen for stroke patients with persistent hiccups: a randomized, double-blind, placebo-controlled trial. *Trials* 2014;15:295.
- 9) Wang T, Wang D. Metoclopramide for patients with intractable hiccups: a multicentre, randomised, controlled pilot study. *Intern Med J* 2014;44:1205-9.
- 10) Edgar NR, Saleh D, Miller RA. Recurrent aphthous stomatitis: a review. *J Clin Aesthet Dermatol* 2017;10(3):26-36.
- 11) 日本口腔外科学会（監修）．チェアサイドで行う口腔がん検診（スクリーニング）．2019.
- 12) Tan ECK, Lexomboon D, Sandborgh-Englund G, Haasum Y, Johnell K. Medications that cause dry mouth as an adverse effect in older people: a systematic review and metaanalysis. *J Am Geriatr Soc* 2018;66:76-84.
- 13) 厚生労働省．2022（令和4）年 国民生活基礎調査の概況．2023.
- 14) Goto S, Umehara J, Aizawa T, Kokubun S. Crowned dens syndrome. *J Bone Joint Surg Am* 2007;89:2732-6.
- 15) Gross A, Kay TM, Paquin JP, et al. Exercises for mechanical neck disorders. *Cochrane Database Syst Rev* 2015;1:CD004250.
- 16) Tunkel DE, Bauer CA, Sun GH, et al. Clinical practice guideline: tinnitus. *Otolaryngol Head Neck Surg* 2014;151(2 Suppl):S1-40.
- 17) Chandrasekhar SS, Tsai Do BS, Schwartz SR, et al. Clinical practice guideline: sudden hearing loss (update). *Otolaryngol Head Neck Surg* 2019;161(1_suppl):S1-45.
- 18) Winkelman JW, Berkowski JA, DelRosso LM, et al. Treatment of restless legs syndrome and periodic limb movement disorder: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med* 2025;21:137-52.
- 19) Allen RP, Picchietti DL, Auerbach M, et al. Evidence-based and consensus clinical practice guidelines for the iron treatment of restless legs syndrome/Willis-Ekbom disease in adults and children: an IRLSSG task force report. *Sleep Med* 2018;41:27-44.
- 20) Baizabal-Carvallo JF, Morgan JC. Drug-induced tremor, clinical features, diagnostic approach and management. *J Neurol Sci* 2022;435:120192.
- 21) Bhatia KP, Bain P, Bajaj N, et al. Consensus statement on the classification of tremors. From the task force on tremor of the International Parkinson and Movement Disorder Society. *Mov Disord* 2018;33:75-87.
- 22) Zesiewicz TA, Elble RJ, Louis ED, et al. Evidence-based guideline update: treatment of essential tremor: report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2011;77:1752-5.
- 23) Postuma RB, Berg D, Stern M, et al. MDS clinical diagnostic criteria for Parkinson's disease. *Mov Disord* 2015;30:1591-601.
- 24) 厚生労働省．重篤副作用疾患別対応マニュアル 薬剤性パーキンソニズム．2006（2022年2月改定）．
- 25) Schrag A, Horsfall L, Walters K, Noyce A, Petersen I. Prediagnostic presentations of Parkinson's disease in primary care: a case-control study. *Lancet Neurol* 2015;14:57-64.

2-26 膠原病・血管炎を疑ったとき

- 1) Tan EM, Feltkamp TE, Smolen JS, et al. Range of antinuclear antibodies in "healthy" individuals. *Arthritis Rheum* 1997;40:1601-11.
- 2) Sato S, Hoshino K, Satoh T, et al. RNA helicase encoded by melanoma differentiation-associated gene 5 is a major autoantigen in patients with clinically amyopathic dermatomyositis: association with rapidly progressive interstitial lung disease. *Arthritis Rheum* 2009;60:2193-200.
- 3) Tsuji H, Nakashima R, Hosono Y, et al. Multicenter prospective study of the efficacy and safety of combined immunosuppressive therapy with high-dose glucocorticoid, tacrolimus, and

- cyclophosphamide in interstitial lung diseases accompanied by anti-melanoma differentiation-associated gene 5-positive dermatomyositis. *Arthritis Rheumatol* 2020;72:488-98.
- 4) 日本循環器学会ほか. 2026年改訂版 大型血管炎診療ガイドライン. 2026.
 - 5) Mahr A, Batteux F, Tubiana S, et al. Brief report: prevalence of antineutrophil cytoplasmic antibodies in infective endocarditis. *Arthritis Rheumatol* 2014;66:1672-7.
 - 6) Aringer M, Costenbader K, Daikh D, et al. 2019 European League Against Rheumatism/ American College of Rheumatology classification criteria for systemic lupus erythematosus. *Ann Rheum Dis* 2019;78:1151-9.
 - 7) Damoiseaux J, Andrade LEC, Carballo OG, et al. Clinical relevance of HEp-2 indirect immunofluorescent patterns: the International Consensus on ANA patterns (ICAP) perspective. *Ann Rheum Dis* 2019;78:879-89.
 - 8) Salazar GA, Assassi S, Wigley F, et al. Antinuclear antibody-negative systemic sclerosis. *Semin Arthritis Rheum* 2015;44:680-6.
 - 9) Grygiel-Górniak B, Rogacka N, Puszczewicz M. Antinuclear antibodies in healthy people and non-rheumatic diseases - diagnostic and clinical implications. *Reumatologia* 2018;56:243-8.
 - 10) Abeles AM, Abeles M. The clinical utility of a positive antinuclear antibody test result. *Am J Med* 2013;126:342-8.
 - 11) Yazdany J, Schmajak G, Robbins M, et al. Choosing wisely: the American College of Rheumatology's Top 5 list of things physicians and patients should question. *Arthritis Care Res (Hoboken)* 2013;65:329-39.
 - 12) Guchelaar NAD, Waling MM, Adhin AA, van Daele PLA, Schreurs MWJ, Rombach SM. The value of anti-neutrophil cytoplasmic antibodies (ANCA) testing for the diagnosis of ANCA-associated vasculitis, a systematic review and meta-analysis. *Autoimmun Rev* 2021;20:102716.
 - 13) Sada KE, Yamamura M, Harigai M, et al. Classification and characteristics of Japanese patients with antineutrophil cytoplasmic antibody-associated vasculitis in a nationwide, prospective, inception cohort study. *Arthritis Res Ther* 2014;16:R101.
 - 14) 厚生労働省保険局医療課長. 診療報酬の算定方法の一部改正に伴う実施上の留意事項について（令和8年3月5日 保医発0305第6号）. 別添1 D014 自己抗体検査. 2026.
 - 15) Sada KE, Amano K, Uehara R, et al. A nationwide survey on the epidemiology and clinical features of eosinophilic granulomatosis with polyangiitis (Churg-Strauss) in Japan. *Mod Rheumatol* 2014;24:640-4.
 - 16) Trallero-Araguás E, Rodrigo-Pendás JÁ, Selva-O'Callaghan A, et al. Usefulness of anti-p155 autoantibody for diagnosing cancer-associated dermatomyositis: a systematic review and meta-analysis. *Arthritis Rheum* 2012;64:523-32.
 - 17) Boddaert J, Huong DLT, Amoura Z, Wechsler B, Godeau P, Piette JC. Late-onset systemic lupus erythematosus: a personal series of 47 patients and pooled analysis of 714 cases in the literature. *Medicine (Baltimore)* 2004;83:348-59.
 - 18) アプレゾリン、アミサリン、イスコチン、ミノマイシン、レミケードの電子添文（KEGG MEDICUSで2026年10月に確認）.
 - 19) Arnaud L, Mertz P, Gavand PE, et al. Drug-induced systemic lupus: revisiting the ever-changing spectrum of the disease using the WHO pharmacovigilance database. *Ann Rheum Dis* 2019;78:504-8.
 - 20) Shiboski CH, Shiboski SC, Seror R, et al. 2016 American College of Rheumatology/European League Against Rheumatism classification criteria for primary Sjögren's syndrome: a consensus and data-driven methodology involving three international patient cohorts. *Ann Rheum Dis* 2017;76:9-16.

- 21) van den Hoogen F, Khanna D, Fransen J, et al. 2013 classification criteria for systemic sclerosis: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Ann Rheum Dis* 2013;72:1747-55.
- 22) Koenig M, Joyal F, Fritzler MJ, et al. Autoantibodies and microvascular damage are independent predictive factors for the progression of Raynaud's phenomenon to systemic sclerosis: a twenty-year prospective study of 586 patients, with validation of proposed criteria for early systemic sclerosis. *Arthritis Rheum* 2008;58:3902-12.
- 23) Spencer-Green G, Alter D, Welch HG. Test performance in systemic sclerosis: anti-centromere and anti-Scl-70 antibodies. *Am J Med* 1997;103:242-8.
- 24) Maverakis E, Patel F, Kronenberg DG, et al. International consensus criteria for the diagnosis of Raynaud's phenomenon. *J Autoimmun* 2014;48-49:60-5.
- 25) Tanaka Y, Kuwana M, Fujii T, et al. 2019 Diagnostic criteria for mixed connective tissue disease (MCTD): from the Japan research committee of the ministry of health, labor, and welfare for systemic autoimmune diseases. *Mod Rheumatol* 2021;31:29-33.
- 26) Harabuchi Y, Kishibe B, Tateyama K, et al. Clinical features and treatment outcomes of otitis media with antineutrophil cytoplasmic antibody (ANCA)-associated vasculitis (OMAAV): a retrospective analysis of 235 patients from a nationwide survey in Japan. *Mod Rheumatol* 2017;27:87-94.
- 27) Grayson PC, Ponte C, Suppiah R, et al. 2022 American College of Rheumatology/EULAR classification criteria for Takayasu arteritis. *Ann Rheum Dis* 2022;81:1654-60.
- 28) International Team for the Revision of the International Criteria for Behçet's Disease (ITR-ICBD). The International Criteria for Behçet's Disease (ICBD): a collaborative study of 27 countries on the sensitivity and specificity of the new criteria. *J Eur Acad Dermatol Venereol* 2014;28:338-47.
- 29) Umehara H, Okazaki K, Masaki Y, et al. Comprehensive diagnostic criteria for IgG4-related disease (IgG4-RD), 2011. *Mod Rheumatol* 2012;22:21-30.
- 30) Barbhaiya M, Zuily S, Naden R, et al. 2023 ACR/EULAR antiphospholipid syndrome classification criteria. *Ann Rheum Dis* 2023;82:1258-70.
- 31) リウマトレックスカプセルの電子添文（KEGG MEDICUSで2026年10月に確認）。
- 32) Kameda H, Yamaoka K, Yamanishi Y, et al. Japan College of Rheumatology guidance for the use of methotrexate in patients with rheumatoid arthritis: secondary publication. *Mod Rheumatol* 2024;34:1-10.
- 33) Nielsen SF, Bojesen SE, Schnohr P, Nordestgaard BG. Elevated rheumatoid factor and long term risk of rheumatoid arthritis: a prospective cohort study. *BMJ* 2012;345:e5244.

第3部 生活習慣病と慢性疾患の継続診療

3-1 高血圧

- 1) Ettehad D, Emdin CA, Kiran A, et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 2016;387:957-67.
- 2) 日本高血圧学会高血圧管理・治療ガイドライン委員会編. 高血圧管理・治療ガイドライン2025. ライフサイエンス出版；2025.（英語版：Hypertens Res 2026;49:9-235）
- 3) Clark CE, Taylor RS, Shore AC, Ukoumunne OC, Campbell JL. Association of a difference in systolic blood pressure between arms with vascular disease and mortality: a systematic review and meta-analysis. *Lancet* 2012;379:905-14.

- 4) Hodgkinson J, Mant J, Martin U, et al. Relative effectiveness of clinic and home blood pressure monitoring compared with ambulatory blood pressure monitoring in diagnosis of hypertension: systematic review. *BMJ* 2011;342:d3621.
- 5) Nieman LK, Biller BM, Findling JW, et al. The diagnosis of Cushing's syndrome: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2008;93:1526-40.
- 6) Zhang W, Zhang S, Deng Y, et al. Trial of intensive blood-pressure control in older patients with hypertension. *N Engl J Med* 2021;385:1268-79.
- 7) He FJ, Li J, MacGregor GA. Effect of longer term modest salt reduction on blood pressure: Cochrane systematic review and meta-analysis of randomised trials. *BMJ* 2013;346:f1325.
- 8) Neal B, Wu Y, Feng X, et al. Effect of salt substitution on cardiovascular events and death. *N Engl J Med* 2021;385:1067-77.
- 9) Wald DS, Law M, Morris JK, Bestwick JP, Wald NJ. Combination therapy versus monotherapy in reducing blood pressure: meta-analysis on 11,000 participants from 42 trials. *Am J Med* 2009;122:290-300.
- 10) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ノルバスク錠、ミカルディス錠、ナトリックス錠、アルダクトンA錠、ミネプロ錠、エンレスト錠、クラビット錠、レニベース錠、メインテート錠）。
- 11) Law MR, Wald NJ, Morris JK, Jordan RE. Value of low dose combination treatment with blood pressure lowering drugs: analysis of 354 randomised trials. *BMJ* 2003;326:1427.
- 12) Williams B, MacDonald TM, Morant S, et al. Spironolactone versus placebo, bisoprolol, and doxazosin to determine the optimal treatment for drug-resistant hypertension (PATHWAY-2): a randomised, double-blind, crossover trial. *Lancet* 2015;386:2059-68.
- 13) 日本腎臓学会 CKD患者の血圧管理に関する検討小委員会. 慢性腎臓病（CKD）患者の血圧管理に関するステートメント. *日腎会誌* 2026;68:253-8.
- 14) 日本循環器学会, 日本心臓血管外科学会, 日本胸部外科学会, 日本血管外科学会. 2020年改訂版 大動脈瘤・大動脈解離診療ガイドライン. 2020.
- 15) Propanolol Aneurysm Trial Investigators. Propranolol for small abdominal aortic aneurysms: results of a randomized trial. *J Vasc Surg* 2002;35:72-9.
- 16) Sweeting MJ, Thompson SG, Brown LC, Powell JT; RESCAN collaborators. Meta-analysis of individual patient data to examine factors affecting growth and rupture of small abdominal aortic aneurysms. *Br J Surg* 2012;99:655-65.
- 17) Pasternak B, Inghammar M, Svanström H. Fluoroquinolone use and risk of aortic aneurysm and dissection: nationwide cohort study. *BMJ* 2018;360:k678.
- 18) Mackenzie IS, Rogers A, Poulter NR, et al. Cardiovascular outcomes in adults with hypertension with evening versus morning dosing of usual antihypertensives in the UK (TIME study): a prospective, randomised, open-label, blinded-endpoint clinical trial. *Lancet* 2022;400:1417-25.
- 19) Makani H, Bangalore S, Romero J, et al. Peripheral edema associated with calcium channel blockers: incidence and withdrawal rate—a meta-analysis of randomized trials. *J Hypertens* 2011;29:1270-80.
- 20) Makani H, Bangalore S, Romero J, Wever-Pinzon O, Messerli FH. Effect of renin-angiotensin system blockade on calcium channel blocker-associated peripheral edema. *Am J Med* 2011;124:128-35.
- 21) Sica D. Calcium channel blocker-related peripheral edema: can it be resolved? *J Clin Hypertens (Greenwich)* 2003;5:291-4, 297.
- 22) van Hamersvelt HW, Kloke HJ, de Jong DJ, Koene RA, Huysmans FT. Oedema formation with the vasodilators nifedipine and diazoxide: direct local effect or sodium retention? *J Hypertens* 1996;14:1041-5.

3-2 糖尿病

- 1) UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837-53.
- 2) Action to Control Cardiovascular Risk in Diabetes Study Group. Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59.
- 3) 日本糖尿病学会編・著. 糖尿病診療ガイドライン2024. 南江堂；2024.
- 4) NCD Risk Factor Collaboration (NCD-RisC). Effects of diabetes definition on global surveillance of diabetes prevalence and diagnosis: a pooled analysis of 96 population-based studies with 331,288 participants. *Lancet Diabetes Endocrinol* 2015;3:624-37.
- 5) 日本糖尿病対策推進会議編. 糖尿病治療のエッセンス 2022年版. 2022. (日本医師会のサイトで公開)
- 6) 坊内良太郎, 近藤龍也, 太田康晴, ほか. 2型糖尿病の薬物療法のアルゴリズム (第2版). *糖尿病* 2023;66:715-33.
- 7) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. メトグルコ錠、ジャディアンス錠、フォーシーガ錠、リベルサス錠、オゼンピック皮下注、マンジャロ皮下注、アクトス錠、トラゼンタ錠、ジャヌビア錠、アマリール錠、ランタス注ソロスター、オングリザ錠) .
- 8) UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 1998;352:854-65.
- 9) Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med* 2015;373:2117-28.
- 10) Nuffield Department of Population Health Renal Studies Group, SGLT2 inhibitor Meta-Analysis Cardio-Renal Trialists' Consortium. Impact of diabetes on the effects of sodium glucose co-transporter-2 inhibitors on kidney outcomes: collaborative meta-analysis of large placebo-controlled trials. *Lancet* 2022;400:1788-801.
- 11) Sattar N, Lee MMY, Kristensen SL, et al. Cardiovascular, mortality, and kidney outcomes with GLP-1 receptor agonists in patients with type 2 diabetes: a systematic review and meta-analysis of randomised trials. *Lancet Diabetes Endocrinol* 2021;9:653-62.
- 12) 日本糖尿病学会 SGLT2阻害薬の適正使用に関する委員会. 糖尿病治療におけるSGLT2阻害薬の適正使用に関するRecommendation. 2022年7月26日改訂. (学会のサイトで公開)
- 13) American Diabetes Association Professional Practice Committee. 9. Pharmacologic approaches to glycemic treatment: Standards of Care in Diabetes—2026. *Diabetes Care* 2026;49(Suppl 1):S183-215.
- 14) サノフィ. 患者向医薬品ガイド ランタス注ソロスター. 2020年9月更新. (医薬品医療機器総合機構のサイトで公開)
- 15) Blanco M, Hernández MT, Strauss KW, Amaya M. Prevalence and risk factors of lipohypertrophy in insulin-injecting patients with diabetes. *Diabetes Metab* 2013;39:445-53.
- 16) 日本腎臓学会編. CKD診療ガイド2024. 東京医学社；2024.
- 17) 島田朗, 川崎英二, 阿比留教生, ほか. 緩徐進行1型糖尿病疑い例への治療介入に関するステートメント. *糖尿病* 2023;66:807-14.
- 18) Schaper NC, van Netten JJ, Apelqvist J, et al. Practical guidelines on the prevention and management of diabetes-related foot disease (IWGDF 2023 update). *Diabetes Metab Res Rev* 2024;40:e3657.
- 19) Gaede P, Vedel P, Larsen N, Jensen GV, Parving HH, Pedersen O. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003;348:383-93.

- 20) 日本糖尿病学会（監修：日本医師会）．かかりつけ医から糖尿病専門医・専門医療機関への紹介基準．2018．（学会のサイトで公開）
- 21) Lean ME, Leslie WS, Barnes AC, et al. Primary care-led weight management for remission of type 2 diabetes (DiRECT): an open-label, cluster-randomised trial. *Lancet* 2018;391:541-51.
- 22) Scirica BM, Bhatt DL, Braunwald E, et al. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* 2013;369:1317-26.
- 23) Green JB, Bethel MA, Armstrong PW, et al. Effect of sitagliptin on cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2015;373:232-42.
- 24) Rosenstock J, Perkovic V, Johansen OE, et al. Effect of linagliptin vs placebo on major cardiovascular events in adults with type 2 diabetes and high cardiovascular and renal risk: the CARMELINA randomized clinical trial. *JAMA* 2019;321:69-79.
- 25) Shi Q, Nong K, Vandvik PO, et al. Benefits and harms of drug treatment for type 2 diabetes: systematic review and network meta-analysis of randomised controlled trials. *BMJ* 2023;381:e074068.

3-3 脂質異常症

- 1) Cholesterol Treatment Trialists' (CTT) Collaboration. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. *Lancet* 2010;376:1670-81.
- 2) 日本動脈硬化学会．動脈硬化性疾患予防ガイドライン2022年版．2022．
- 3) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認．クレストール錠、リビトール錠、リパロ錠、ゼチーア錠、パルモディXR錠、メバロチン錠、レパーサ皮下注、レクビオ皮下注、ネクセトール錠、エバデールS）．
- 4) Li MY, Chen JH, Chen C, Kang YN. Association between egg consumption and cholesterol concentration: a systematic review and meta-analysis of randomized controlled trials. *Nutrients* 2020;12:1995.
- 5) Nakamura H, Arakawa K, Itakura H, et al. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *Lancet* 2006;368:1155-63.
- 6) Nicholls SJ, Brandrup-Wognsen G, Palmer M, Barter PJ. Meta-analysis of comparative efficacy of increasing dose of atorvastatin versus rosuvastatin versus simvastatin on lowering levels of atherogenic lipids (from VOYAGER). *Am J Cardiol* 2010;105:69-76.
- 7) Cannon CP, Blazing MA, Giugliano RP, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *N Engl J Med* 2015;372:2387-97.
- 8) Sabatine MS, Giugliano RP, Keech AC, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *N Engl J Med* 2017;376:1713-22.
- 9) Nissen SE, Lincoff AM, Brennan D, et al. Bempedoic acid and cardiovascular outcomes in statin-intolerant patients. *N Engl J Med* 2023;388:1353-64.
- 10) Ouchi Y, Sasaki J, Arai H, et al. Ezetimibe lipid-lowering trial on prevention of atherosclerotic cardiovascular disease in 75 or older (EWTPIA 75): a randomized, controlled trial. *Circulation* 2019;140:992-1003.
- 11) Cholesterol Treatment Trialists' Collaboration. Effect of statin therapy on muscle symptoms: an individual participant data meta-analysis of large-scale, randomised, double-blind trials. *Lancet* 2022;400:832-45.
- 12) 日本動脈硬化学会．スタチン不耐に関する診療指針2018（2022年12月改訂）．

- 13) Mammen AL, Chung T, Christopher-Stine L, et al. Autoantibodies against 3-hydroxy-3-methylglutaryl-coenzyme A reductase in patients with statin-associated autoimmune myopathy. *Arthritis Rheum* 2011;63:713-21.
- 14) Das Pradhan A, Glynn RJ, Fruchart JC, et al. Triglyceride lowering with pemafibrate to reduce cardiovascular risk. *N Engl J Med* 2022;387:1923-34.
- 15) Yokoyama M, Origasa H, Matsuzaki M, et al. Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomised open-label, blinded endpoint analysis. *Lancet* 2007;369:1090-8.
- 16) Giral P, Neumann A, Weill A, Coste J. Cardiovascular effect of discontinuing statins for primary prevention at the age of 75 years: a nationwide population-based cohort study in France. *Eur Heart J* 2019;40:3516-25.
- 17) Howard JP, Wood FA, Finegold JA, et al. Side effect patterns in a crossover trial of statin, placebo, and no treatment. *J Am Coll Cardiol* 2021;78:1210-22.
- 18) Giugliano RP, Pedersen TR, Park JG, et al. Clinical efficacy and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab: a prespecified secondary analysis of the FOURIER trial. *Lancet* 2017;390:1962-71.
- 19) Gaba P, O'Donoghue ML, Park JG, et al. Association between achieved low-density lipoprotein cholesterol levels and long-term cardiovascular and safety outcomes: an analysis of FOURIER-OLE. *Circulation* 2023;147:1192-203.
- 20) Giugliano RP, Mach F, Zavitz K, et al. Cognitive function in a randomized trial of evolocumab. *N Engl J Med* 2017;377:633-43.
- 21) Lee M, Cheng CY, Wu YL, Lee JD, Hsu CY, Ovbiagele B. Association between intensity of low-density lipoprotein cholesterol reduction with statin-based therapies and secondary stroke prevention: a meta-analysis of randomized clinical trials. *JAMA Neurol* 2022;79:349-58.

3-4 肥満症・脂肪性肝疾患と肝機能異常

- 1) 日本肝臓学会. 奈良宣言2023. 2023年6月15日.
- 2) Hernaez R, Lazo M, Bonekamp S, et al. Diagnostic accuracy and reliability of ultrasonography for the detection of fatty liver: a meta-analysis. *Hepatology* 2011;54:1082-90.
- 3) 厚生労働省難治性疾患政策研究事業「難治性の肝・胆道疾患に関する調査研究」班. 自己免疫性肝炎(AIH) 診療ガイドライン (2021年). 2022.
- 4) 国立健康危機管理研究機構 感染症情報提供サイト. A型肝炎(詳細版). <https://id-info.jihs.go.jp/> (2026年10月に確認)
- 5) 厚生労働省. E型肝炎ウイルスの感染事例・E型肝炎Q&A (2005年8月現在). <https://www.mhlw.go.jp/houdou/2003/08/h0819-2a.html> (2026年10月に確認)
- 6) Lleo A, Wang GQ, Gershwin ME, Hirschfield GM. Primary biliary cholangitis. *Lancet* 2020;396:1915-26.
- 7) 厚生労働省. 肝炎対策の推進に関する基本的な指針(平成23年5月16日策定、令和4年3月7日改正).
- 8) 日本肝臓学会 肝炎診療ガイドライン作成委員会編. C型肝炎治療ガイドライン(第8.4版). 2025.
- 9) 日本肝臓学会 肝炎診療ガイドライン作成委員会編. B型肝炎治療ガイドライン(第5版). 2026.
- 10) Rinella ME, Lazarus JV, Ratziu V, et al. A multisociety Delphi consensus statement on new fatty liver disease nomenclature. *Hepatology* 2023;78:1966-86.
- 11) Akuta N, Kogiso T, Ikejima K, et al. Evidence-based clinical practice guidelines for metabolic dysfunction-associated steatotic liver disease (MASLD) 2026. *J Gastroenterol* 2026;61:693-710. (日本消化器病学会・日本肝臓学会 MASLD診療ガイドライン2026の英語版)
- 12) 日本消化器病学会・日本肝臓学会編. NAFLD/NASH診療ガイドライン2020(改訂第2版). 南江堂; 2020.

- 13) Shah AG, Lydecker A, Murray K, et al. Comparison of noninvasive markers of fibrosis in patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2009;7:1104-12.
- 14) McPherson S, Hardy T, Dufour JF, et al. Age as a confounding factor for the accurate non-invasive diagnosis of advanced NAFLD fibrosis. *Am J Gastroenterol* 2017;112:740-51.
- 15) Vilar-Gomez E, Martinez-Perez Y, Calzadilla-Bertot L, et al. Weight loss through lifestyle modification significantly reduces features of nonalcoholic steatohepatitis. *Gastroenterology* 2015;149:367-78.e5.
- 16) 日本肥満学会編. 肥満症診療ガイドライン2022. ライフサイエンス出版；2022.
- 17) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ウゴービ皮下注、ゼップバウンド皮下注、サノレックス錠、ジャディアンス錠）。
- 18) Kadowaki T, Isendahl J, Khalid U, et al. Semaglutide once a week in adults with overweight or obesity, with or without type 2 diabetes in an east Asian population (STEP 6): a randomised, double-blind, double-dummy, placebo-controlled, phase 3a trial. *Lancet Diabetes Endocrinol* 2022;10:193-206.
- 19) Kadowaki T, Kiyosue A, Shingaki T, Oura T, Yokote K. Efficacy and safety of once-weekly tirzepatide in Japanese patients with obesity disease (SURMOUNT-J): a multicentre, randomised, double-blind, placebo-controlled phase 3 trial. *Lancet Diabetes Endocrinol* 2025;13:384-96.
- 20) 厚生労働省. 最適使用推進ガイドライン セマグルチド（遺伝子組換え）（販売名：ウゴービ皮下注）～肥満症～. 令和5年11月（令和8年6月一部改正）. 同 チルゼパチド（販売名：ゼップバウンド皮下注）～肥満症～. 令和7年3月（令和8年5月一部改正）. 同 セマグルチド～代謝機能障害関連脂肪肝炎～. 令和8年6月. 同 チルゼパチド～閉塞性睡眠時無呼吸症候群～. 令和8年5月.
- 21) Wilding JPH, Batterham RL, Davies M, et al. Weight regain and cardiometabolic effects after withdrawal of semaglutide: the STEP 1 trial extension. *Diabetes Obes Metab* 2022;24:1553-64.
- 22) Sanyal AJ, Newsome PN, Kliers I, et al. Phase 3 trial of semaglutide in metabolic dysfunction-associated steatohepatitis. *N Engl J Med* 2025;392:2089-99.
- 23) 日本消化器病学会・日本肝臓学会編. 肝硬変診療ガイドライン2020（改訂第3版）. 南江堂；2020.
- 24) de Franchis R, Bosch J, Garcia-Tsao G, Reiberger T, Ripoll C; Baveno VII Faculty. Baveno VII – renewing consensus in portal hypertension. *J Hepatol* 2022;76:959-74.
- 25) 厚生労働省. ビブリオ・バルニフィカスに関するQ&A. 2006（2016年11月更新）.
- 26) Steinberg DM, Tate DF, Bennett GG, Ennett S, Samuel-Hodge C, Ward DS. The efficacy of a daily self-weighing weight loss intervention using smart scales and e-mail. *Obesity* (Silver Spring) 2013;21:1789-97.
- 27) Gardner CD, Trepanowski JF, Del Gobbo LC, et al. Effect of low-fat vs low-carbohydrate diet on 12-month weight loss in overweight adults and the association with genotype pattern or insulin secretion: the DIETFITS randomized clinical trial. *JAMA* 2018;319:667-79.
- 28) Ge L, Sadeghirad B, Ball GDC, et al. Comparison of dietary macronutrient patterns of 14 popular named dietary programmes for weight and cardiovascular risk factor reduction in adults: systematic review and network meta-analysis of randomised trials. *BMJ* 2020;369:m696.
- 29) Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA* 2005;293:43-53.

3-5 高尿酸血症・痛風と尿路結石

- 1) 日本痛風・核酸代謝学会 ガイドライン改訂委員会編. 高尿酸血症・痛風の治療ガイドライン 第3版. 診断と治療社；2018.

- 2) FitzGerald JD, Dalbeth N, Mikuls T, et al. 2020 American College of Rheumatology guideline for the management of gout. *Arthritis Care Res (Hoboken)* 2020;72:744-60.
- 3) Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Alcohol intake and risk of incident gout in men: a prospective study. *Lancet* 2004;363:1277-81.
- 4) Choi HK, Curhan G. Soft drinks, fructose consumption, and the risk of gout in men: prospective cohort study. *BMJ* 2008;336:309-12.
- 5) Choi HK, Atkinson K, Karlson EW, Willett W, Curhan G. Purine-rich foods, dairy and protein intake, and the risk of gout in men. *N Engl J Med* 2004;350:1093-103.
- 6) Juraschek SP, Gelber AC, Choi HK, Appel LJ, Miller ER 3rd. Effects of the Dietary Approaches to Stop Hypertension (DASH) diet and sodium intake on serum uric acid. *Arthritis Rheumatol* 2016;68:3002-9.
- 7) Nielsen SM, Bartels EM, Henriksen M, et al. Weight loss for overweight and obese individuals with gout: a systematic review of longitudinal studies. *Ann Rheum Dis* 2017;76:1870-82.
- 8) Campion EW, Glynn RJ, DeLabry LO. Asymptomatic hyperuricemia. Risks and consequences in the Normative Aging Study. *Am J Med* 1987;82:421-6.
- 9) Badve SV, Pascoe EM, Tikun A, et al. Effects of allopurinol on the progression of chronic kidney disease. *N Engl J Med* 2020;382:2504-13.
- 10) Kimura K, Hosoya T, Uchida S, et al. Febuxostat therapy for patients with stage 3 CKD and asymptomatic hyperuricemia: a randomized trial. *Am J Kidney Dis* 2018;72:798-810.
- 11) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。フェブリック、アロプリノール錠、ユリス、ユリノーム、コルヒチン錠、ウラリット、トピロリック）。
- 12) 日本腎臓学会編。CKD診療ガイド2024。東京医学社；2024。
- 13) Stamp LK, Taylor WJ, Jones PB, et al. Starting dose is a risk factor for allopurinol hypersensitivity syndrome: a proposed safe starting dose of allopurinol. *Arthritis Rheum* 2012;64:2529-36.
- 14) White WB, Saag KG, Becker MA, et al. Cardiovascular safety of febuxostat or allopurinol in patients with gout. *N Engl J Med* 2018;378:1200-10.
- 15) Mackenzie IS, Ford I, Nuki G, et al. Long-term cardiovascular safety of febuxostat compared with allopurinol in patients with gout (FAST): a multicentre, prospective, randomised, open-label, non-inferiority trial. *Lancet* 2020;396:1745-57.
- 16) Borstad GC, Bryant LR, Abel MP, Scroggie DA, Harris MD, Alloway JA. Colchicine for prophylaxis of acute flares when initiating allopurinol for chronic gouty arthritis. *J Rheumatol* 2004;31:2429-32.
- 17) 日本泌尿器科学会，日本尿路結石症学会，日本泌尿器内視鏡・ロボティクス学会編。尿路結石症診療ガイドライン 第3版。2023。
- 18) Borghi L, Meschi T, Amato F, Briganti A, Novarini A, Giannini A. Urinary volume, water and recurrences in idiopathic calcium nephrolithiasis: a 5-year randomized prospective study. *J Urol* 1996;155:839-43.
- 19) Borghi L, Schianchi T, Meschi T, et al. Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria. *N Engl J Med* 2002;346:77-84.
- 20) Ferraro PM, Curhan GC, Gambaro G, Taylor EN. Total, dietary, and supplemental vitamin C intake and risk of incident kidney stones. *Am J Kidney Dis* 2016;67:400-7.
- 21) Dhayat NA, Bonny O, Roth B, et al. Hydrochlorothiazide and prevention of kidney-stone recurrence. *N Engl J Med* 2023;388:781-91.
- 22) Doherty M, Jenkins W, Richardson H, et al. Efficacy and cost-effectiveness of nurse-led care involving education and engagement of patients and a treat-to-target urate-lowering strategy versus usual care for gout: a randomised controlled trial. *Lancet* 2018;392:1403-12.

- 23) Beslon V, Moreau P, Maruani A, Maisonneuve H, Giraudeau B, Fournier JP. Effects of discontinuation of urate-lowering therapy: a systematic review. *J Gen Intern Med* 2018;33:358-66.

3-6 甲状腺機能の異常と甲状腺結節

- 1) 日本甲状腺学会. 甲状腺疾患診断ガイドライン2024 (2024年11月21日改定) .
- 2) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。チラーゼンS錠、メルカゾール錠、チウラジール錠、インデラル錠、アンカロン錠、リーマス錠) .
- 3) Barroso-Sousa R, Barry WT, Garrido-Castro AC, et al. Incidence of endocrine dysfunction following the use of different immune checkpoint inhibitor regimens: a systematic review and meta-analysis. *JAMA Oncol* 2018;4:173-82.
- 4) Li D, Radulescu A, Shrestha RT, et al. Association of biotin ingestion with performance of hormone and nonhormone assays in healthy adults. *JAMA* 2017;318:1150-60.
- 5) Vanderpump MP, Tunbridge WM, French JM, et al. The incidence of thyroid disorders in the community: a twenty-year follow-up of the Whickham Survey. *Clin Endocrinol (Oxf)* 1995;43:55-68.
- 6) Pearce SH, Brabant G, Duntas LH, et al. 2013 ETA guideline: management of subclinical hypothyroidism. *Eur Thyroid J* 2013;2:215-28.
- 7) Stott DJ, Rodondi N, Kearney PM, et al. Thyroid hormone therapy for older adults with subclinical hypothyroidism. *N Engl J Med* 2017;376:2534-44.
- 8) Kasagi K, Iwata M, Misaki T, Konishi J. Effect of iodine restriction on thyroid function in patients with primary hypothyroidism. *Thyroid* 2003;13:561-7.
- 9) Bolk N, Visser TJ, Nijman J, Jongste IJ, Tijssen JG, Berghout A. Effects of evening vs morning levothyroxine intake: a randomized double-blind crossover trial. *Arch Intern Med* 2010;170:1996-2003.
- 10) Flynn RW, Bonellie SR, Jung RT, MacDonald TM, Morris AD, Leese GP. Serum thyroid-stimulating hormone concentration and morbidity from cardiovascular disease and fractures in patients on long-term thyroxine therapy. *J Clin Endocrinol Metab* 2010;95:186-93.
- 11) Alexander EK, Marqusee E, Lawrence J, Jarolim P, Fischer GA, Larsen PR. Timing and magnitude of increases in levothyroxine requirements during pregnancy in women with hypothyroidism. *N Engl J Med* 2004;351:241-9.
- 12) Yassa L, Marqusee E, Fawcett R, Alexander EK. Thyroid hormone early adjustment in pregnancy (the THERAPY) trial. *J Clin Endocrinol Metab* 2010;95:3234-41.
- 13) Boelaert K, Torlinska B, Holder RL, Franklyn JA. Older subjects with hyperthyroidism present with a paucity of symptoms and signs: a large cross-sectional study. *J Clin Endocrinol Metab* 2010;95:2715-26.
- 14) Tozzoli R, Bagnasco M, Giavarina D, Bizzaro N. TSH receptor autoantibody immunoassay in patients with Graves' disease: improvement of diagnostic accuracy over different generations of methods. Systematic review and meta-analysis. *Autoimmun Rev* 2012;12:107-13.
- 15) Kubota S, Nishihara E, Kudo T, Ito M, Amino N, Miyauchi A. Initial treatment with 15 mg of prednisolone daily is sufficient for most patients with subacute thyroiditis in Japan. *Thyroid* 2013;23:269-72.
- 16) Sawin CT, Geller A, Wolf PA, et al. Low serum thyrotropin concentrations as a risk factor for atrial fibrillation in older persons. *N Engl J Med* 1994;331:1249-52.
- 17) Biondi B, Bartalena L, Cooper DS, Hegedüs L, Laurberg P, Kahaly GJ. The 2015 European Thyroid Association guidelines on diagnosis and treatment of endogenous subclinical hyperthyroidism. *Eur Thyroid J* 2015;4:149-63.

- 18) Nakamura H, Noh JY, Itoh K, Fukata S, Miyauchi A, Hamada N. Comparison of methimazole and propylthiouracil in patients with hyperthyroidism caused by Graves' disease. *J Clin Endocrinol Metab* 2007;92:2157-62.
- 19) Nakamura H, Miyauchi A, Miyawaki N, Imagawa J. Analysis of 754 cases of antithyroid drug-induced agranulocytosis over 30 years in Japan. *J Clin Endocrinol Metab* 2013;98:4776-83.
- 20) Kahaly GJ, Bartalena L, Hegedüs L, Leenhardt L, Poppe K, Pearce SH. 2018 European Thyroid Association guideline for the management of Graves' hyperthyroidism. *Eur Thyroid J* 2018;7:167-86.
- 21) Bartalena L, Kahaly GJ, Baldeschi L, et al. The 2021 European Group on Graves' orbitopathy (EUGOGO) clinical practice guidelines for the medical management of Graves' orbitopathy. *Eur J Endocrinol* 2021;185:G43-67.
- 22) 日本産科婦人科学会・日本産婦人科医会編. 産婦人科診療ガイドラインー産科編2020. 2020.
- 23) Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2016;26:1-133.
- 24) Brito JP, Gionfriddo MR, Al Nofal A, et al. The accuracy of thyroid nodule ultrasound to predict thyroid cancer: systematic review and meta-analysis. *J Clin Endocrinol Metab* 2014;99:1253-63.
- 25) US Preventive Services Task Force. Screening for thyroid cancer: US Preventive Services Task Force recommendation statement. *JAMA* 2017;317:1882-7.
- 26) Kamijo K. Study on cutoff value setting for differential diagnosis between Graves' disease and painless thyroiditis using the TRAb (Elecsys TRAb) measurement via the fully automated electrochemiluminescence immunoassay system. *Endocr J* 2010;57:895-902.
- 27) Ota H, Amino N, Morita S, et al. Quantitative measurement of thyroid blood flow for differentiation of painless thyroiditis from Graves' disease. *Clin Endocrinol (Oxf)* 2007;67:41-5.
- 28) Yoshimura Noh J, Momotani N, Fukada S, Ito K, Miyauchi A, Amino N. Ratio of serum free triiodothyronine to free thyroxine in Graves' hyperthyroidism and thyrotoxicosis caused by painless thyroiditis. *Endocr J* 2005;52:537-42.
- 29) Furumura Y, Ito M, Monno I, et al. Differentiating TRAb-positive painless thyroiditis from Graves' disease using biochemical and ultrasound findings. *Endocr J* 2026;73:1017-26.

3-7 骨粗鬆症

- 1) 骨粗鬆症の予防と治療ガイドライン作成委員会編. 骨粗鬆症の予防と治療ガイドライン2025年版. ライフサイエンス出版；2025.
- 2) Green AD, Colón-Emeric CS, Bastian L, Drake MT, Lyles KW. Does this woman have osteoporosis? *JAMA* 2004;292:2890-900.
- 3) Hagino H, Sawaguchi T, Endo N, Ito Y, Nakano T, Watanabe Y. The risk of a second hip fracture in patients after their first hip fracture. *Calcif Tissue Int* 2012;90:14-21.
- 4) Lyles KW, Colón-Emeric CS, Magaziner JS, et al. Zoledronic acid and clinical fractures and mortality after hip fracture. *N Engl J Med* 2007;357:1799-809.
- 5) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ボナロン、アクトネル、リクラスト、ブラリア、イベニティ、フォルテオ、エビスタ、エディロール）。
- 6) Black DM, Cummings SR, Karpf DB, et al. Randomised trial of effect of alendronate on risk of fracture in women with existing vertebral fractures. *Lancet* 1996;348:1535-41.
- 7) Cummings SR, San Martin J, McClung MR, et al. Denosumab for prevention of fractures in postmenopausal women with osteoporosis. *N Engl J Med* 2009;361:756-65.
- 8) Saag KG, Petersen J, Brandi ML, et al. Romosozumab or alendronate for fracture prevention in women with osteoporosis. *N Engl J Med* 2017;377:1417-27.

- 9) Kendler DL, Marin F, Zerbini CAF, et al. Effects of teriparatide and risedronate on new fractures in post-menopausal women with severe osteoporosis (VERO): a multicentre, double-blind, double-dummy, randomised controlled trial. *Lancet* 2018;391:230-40.
- 10) 日本透析医学会. 慢性腎臓病に伴う骨・ミネラル代謝異常の診療ガイドライン (2025年改訂版). 透析会誌 2026;59:127-224.
- 11) 日本腎臓学会編. エビデンスに基づくCKD診療ガイドライン2023. 東京医学社; 2023.
- 12) Bird ST, Smith ER, Gelperin K, et al. Severe hypocalcemia with denosumab among older female dialysis-dependent patients. *JAMA* 2024;331:491-9.
- 13) Black DM, Geiger EJ, Eastell R, et al. Atypical femur fracture risk versus fragility fracture prevention with bisphosphonates. *N Engl J Med* 2020;383:743-53.
- 14) Black DM, Schwartz AV, Ensrud KE, et al. Effects of continuing or stopping alendronate after 5 years of treatment: the Fracture Intervention Trial Long-term Extension (FLEX): a randomized trial. *JAMA* 2006;296:2927-38.
- 15) Cummings SR, Ferrari S, Eastell R, et al. Vertebral fractures after discontinuation of denosumab: a post hoc analysis of the randomized placebo-controlled FREEDOM trial and its extension. *J Bone Miner Res* 2018;33:190-8.
- 16) Lyu H, Yoshida K, Zhao SS, et al. Delayed denosumab injections and fracture risk among patients with osteoporosis: a population-based cohort study. *Ann Intern Med* 2020;173:516-26.
- 17) 顎骨壊死検討委員会. 薬剤関連顎骨壊死の病態と管理: 顎骨壊死検討委員会ポジションペーパー2023. 日本口腔外科学会ほか; 2023.
- 18) Gourlay ML, Fine JP, Preisser JS, et al. Bone-density testing interval and transition to osteoporosis in older women. *N Engl J Med* 2012;366:225-33.
- 19) Tsourdi E, Zillikens MC, Meier C, et al. Fracture risk and management of discontinuation of denosumab therapy: a systematic review and position statement by ECTS. *J Clin Endocrinol Metab* 2021;106:264-81.
- 20) LeBoff MS, Chou SH, Ratliff KA, et al. Supplemental vitamin D and incident fractures in midlife and older adults. *N Engl J Med* 2022;387:299-309.
- 21) Lyu H, Jundi B, Xu C, et al. Comparison of denosumab and bisphosphonates in patients with osteoporosis: a meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab* 2019;104:1753-65.
- 22) Curtis JR, Arora T, Liu Y, et al. Comparative effectiveness of denosumab vs alendronate among postmenopausal women with osteoporosis. *J Bone Miner Res* 2024;39:826-34.

3-8 喘息・COPDの吸入療法

- 1) Sanchis J, Gich I, Pedersen S; Aerosol Drug Management Improvement Team (ADMIT). Systematic review of errors in inhaler use: has patient technique improved over time? *Chest* 2016;150:394-406.
- 2) Global Initiative for Asthma. Global strategy for asthma management and prevention, 2026 update. 2026.
- 3) Niimi A, Fukunaga K, Taniguchi M, et al. Executive summary: Japanese guidelines for adult asthma (JGL) 2021. *Allergol Int* 2023;72:207-26.
- 4) 日本呼吸器学会COPDガイドライン第6版作成委員会編. COPD (慢性閉塞性肺疾患) 診断と治療のためのガイドライン2022 [第6版]. メディカルレビュー社; 2022.
- 5) Karrasch S, Linde K, Rücker G, et al. Accuracy of FENO for diagnosing asthma: a systematic review. *Thorax* 2017;72:109-16.
- 6) Straus SE, McAlister FA, Sackett DL, Deeks JJ. The accuracy of patient history, wheezing, and laryngeal measurements in diagnosing obstructive airway disease. *JAMA* 2000;283:1853-7.

- 7) Fukuchi Y, Nishimura M, Ichinose M, et al. COPD in Japan: the Nippon COPD Epidemiology study. *Respirology* 2004;9:458-65.
- 8) Kerstjens HA, Engel M, Dahl R, et al. Tiotropium in asthma poorly controlled with standard combination therapy. *N Engl J Med* 2012;367:1198-207.
- 9) Sobieraj DM, Weeda ER, Nguyen E, et al. Association of inhaled corticosteroids and long-acting β -agonists as controller and quick relief therapy with exacerbations and symptom control in persistent asthma: a systematic review and meta-analysis. *JAMA* 2018;319:1485-96.
- 10) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 11) Vogelmeier C, Hederer B, Glaab T, et al. Tiotropium versus salmeterol for the prevention of exacerbations of COPD. *N Engl J Med* 2011;364:1093-103.
- 12) Lipson DA, Barnhart F, Brealey N, et al. Once-daily single-inhaler triple versus dual therapy in patients with COPD. *N Engl J Med* 2018;378:1671-80.
- 13) Global Initiative for Chronic Obstructive Lung Disease. Global strategy for prevention, diagnosis and management of COPD: 2026 report. 2025.
- 14) Long-Term Oxygen Treatment Trial Research Group. A randomized trial of long-term oxygen for COPD with moderate desaturation. *N Engl J Med* 2016;375:1617-27.
- 15) Anthonisen NR, Connett JE, Kiley JP, et al. Effects of smoking intervention and the use of an inhaled anticholinergic bronchodilator on the rate of decline of FEV1: the Lung Health Study. *JAMA* 1994;272:1497-505.
- 16) Anthonisen NR, Skeans MA, Wise RA, et al. The effects of a smoking cessation intervention on 14.5-year mortality: a randomized clinical trial. *Ann Intern Med* 2005;142:233-9.
- 17) Ni Chroinin M, Greenstone I, Lasserson TJ, Ducharme FM. Addition of inhaled long-acting beta2-agonists to inhaled steroids as first line therapy for persistent asthma in steroid-naïve adults and children. *Cochrane Database Syst Rev* 2009;(4):CD005307.

3-9 抗血栓薬の外来管理

- 1) Davidson BL, Verheijen S, Lensing AW, et al. Bleeding risk of patients with acute venous thromboembolism taking nonsteroidal anti-inflammatory drugs or aspirin. *JAMA Intern Med* 2014;174:947-53.
- 2) Chai-Adisaksopha C, Hillis C, Monreal M, Witt DM, Crowther M. Thromboembolic events, recurrent bleeding and mortality after resuming anticoagulant following gastrointestinal bleeding: a meta-analysis. *Thromb Haemost* 2015;114:819-25.
- 3) Yao X, Shah ND, Sangaralingham LR, Gersh BJ, Noseworthy PA. Non-vitamin K antagonist oral anticoagulant dosing in patients with atrial fibrillation and renal dysfunction. *J Am Coll Cardiol* 2017;69:2779-90.
- 4) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。エリキュース錠、プラザキサカプセル、リクシアナ錠、イグザレルト錠、ワーファリン錠、プレタールOD錠、ブラビックス錠）。
- 5) Connolly SJ, Karthikeyan G, Ntsekhe M, et al. Rivaroxaban in rheumatic heart disease-associated atrial fibrillation. *N Engl J Med* 2022;387:978-88.
- 6) Pengo V, Denas G, Zoppellaro G, et al. Rivaroxaban vs warfarin in high-risk patients with antiphospholipid syndrome. *Blood* 2018;132:1365-71.
- 7) Holbrook A, Schulman S, Witt DM, et al. Evidence-based management of anticoagulant therapy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest* 2012;141(2 Suppl):e152S-84S.
- 8) 日本循環器学会, 日本不整脈心電学会. 2024年JCS/JHRSガイドライン フォーカスアップデート版 不整脈治療. 2024.

- 9) Khan F, Rahman A, Carrier M, et al. Long term risk of symptomatic recurrent venous thromboembolism after discontinuation of anticoagulant treatment for first unprovoked venous thromboembolism event: systematic review and meta-analysis. *BMJ* 2019;366:l4363.
- 10) 日本循環器学会, 日本肺高血圧・肺循環学会. 2025年改訂版 肺血栓塞栓症・深部静脈血栓症および肺高血圧症に関するガイドライン. 2025.
- 11) Carrier M, Lazo-Langner A, Shivakumar S, et al. Screening for occult cancer in unprovoked venous thromboembolism. *N Engl J Med* 2015;373:697-704.
- 12) Martinelli I, Lensing AW, Middeldorp S, et al. Recurrent venous thromboembolism and abnormal uterine bleeding with anticoagulant and hormone therapy use. *Blood* 2016;127:1417-25.
- 13) Koo BK, Kang J, Park KW, et al. Aspirin versus clopidogrel for chronic maintenance monotherapy after percutaneous coronary intervention (HOST-EXAM): an investigator-initiated, prospective, randomised, open-label, multicentre trial. *Lancet* 2021;397:2487-96.
- 14) Diener HC, Bogousslavsky J, Brass LM, et al. Aspirin and clopidogrel compared with clopidogrel alone after recent ischaemic stroke or transient ischaemic attack in high-risk patients (MATCH): randomised, double-blind, placebo-controlled trial. *Lancet* 2004;364:331-7.
- 15) Toyoda K, Uchiyama S, Yamaguchi T, et al. Dual antiplatelet therapy using cilostazol for secondary prevention in patients with high-risk ischaemic stroke in Japan: a multicentre, open-label, randomised controlled trial. *Lancet Neurol* 2019;18:539-48.
- 16) Ikeda Y, Shimada K, Teramoto T, et al. Low-dose aspirin for primary prevention of cardiovascular events in Japanese patients 60 years or older with atherosclerotic risk factors: a randomized clinical trial. *JAMA* 2014;312:2510-20.
- 17) Saito Y, Okada S, Ogawa H, et al. Low-dose aspirin for primary prevention of cardiovascular events in patients with type 2 diabetes mellitus: 10-year follow-up of a randomized controlled trial. *Circulation* 2017;135:659-70.
- 18) Sundström J, Hedberg J, Thuresson M, Aarskog P, Johannesen KM, Oldgren J. Low-dose aspirin discontinuation and risk of cardiovascular events: a Swedish nationwide, population-based cohort study. *Circulation* 2017;136:1183-92.
- 19) 日本有病者歯科医療学会, 日本口腔外科学会, 日本老年歯科医学会. 抗血栓療法患者の抜歯に関するガイドライン2020年版. 学術社; 2020.
- 20) 日本循環器学会, 日本不整脈心電学会. 2020年改訂版 不整脈薬物治療ガイドライン. 2020 (2023年10月更新).
- 21) Donzé J, Clair C, Hug B, et al. Risk of falls and major bleeds in patients on oral anticoagulation therapy. *Am J Med* 2012;125:773-8.
- 22) Man-Son-Hing M, Nichol G, Lau A, Laupacis A. Choosing antithrombotic therapy for elderly patients with atrial fibrillation who are at risk for falls. *Arch Intern Med* 1999;159:677-85.
- 23) Lau WCY, Torre CO, Man KKC, et al. Comparative effectiveness and safety between apixaban, dabigatran, edoxaban, and rivaroxaban among patients with atrial fibrillation: a multinational population-based cohort study. *Ann Intern Med* 2022;175:1515-24.
- 24) Ray WA, Chung CP, Stein CM, et al. Association of rivaroxaban vs apixaban with major ischemic or hemorrhagic events in patients with atrial fibrillation. *JAMA* 2021;326:2395-404.

3-10 慢性心不全の外来管理

- 1) 日本循環器学会, 日本心不全学会. 2025年改訂版 心不全診療ガイドライン. 2025.
- 2) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. エンレスト錠、アーチスト錠、セララ錠、フォシーガ錠、ケレンディア錠、ロケルマ懸濁用散、プレタールOD錠、アクトス錠、コララン錠、アルダクトンA、ラシックス錠、サムスカOD錠) .

- 3) Vaduganathan M, Claggett BL, Jhund PS, et al. Estimating lifetime benefits of comprehensive disease-modifying pharmacological therapies in patients with heart failure with reduced ejection fraction: a comparative analysis of three randomised controlled trials. *Lancet* 2020;396:121-8.
- 4) Halliday BP, Wassall R, Lota AS, et al. Withdrawal of pharmacological treatment for heart failure in patients with recovered dilated cardiomyopathy (TRED-HF): an open-label, pilot, randomised trial. *Lancet* 2019;393:61-73.
- 5) Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med* 1999;341:709-17.
- 6) McMurray JJV, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014;371:993-1004.
- 7) CIBIS-II Investigators and Committees. The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomised trial. *Lancet* 1999;353:9-13.
- 8) Zannad F, McMurray JJV, Krum H, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med* 2011;364:11-21.
- 9) McMurray JJV, Solomon SD, Inzucchi SE, et al. Dapagliflozin in patients with heart failure and reduced ejection fraction. *N Engl J Med* 2019;381:1995-2008.
- 10) Anker SD, Butler J, Filippatos G, et al. Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med* 2021;385:1451-61.
- 11) Solomon SD, McMurray JJV, Claggett B, et al. Dapagliflozin in heart failure with mildly reduced or preserved ejection fraction. *N Engl J Med* 2022;387:1089-98.
- 12) Solomon SD, McMurray JJV, Vaduganathan M, et al. Finerenone in heart failure with mildly reduced or preserved ejection fraction. *N Engl J Med* 2024;391:1475-85.
- 13) González-López E, Gallego-Delgado M, Guzzo-Merello G, et al. Wild-type transthyretin amyloidosis as a cause of heart failure with preserved ejection fraction. *Eur Heart J* 2015;36:2585-94.
- 14) Clark AL, Kalra PR, Petrie MC, Mark PB, Tomlinson LA, Tomson CR. Change in renal function associated with drug treatment in heart failure: national guidance. *Heart* 2019;105:904-10.
- 15) Adamson C, Docherty KF, Heerspink HJL, et al. Initial decline (dip) in estimated glomerular filtration rate after initiation of dapagliflozin in patients with heart failure and reduced ejection fraction: insights from DAPA-HF. *Circulation* 2022;146:438-49.
- 16) Kosiborod MN, Cherney DZI, Desai AS, et al. Sodium zirconium cyclosilicate for management of hyperkalemia during spironolactone optimization in patients with heart failure. *J Am Coll Cardiol* 2025;85:971-84.
- 17) Mentz RJ, Anstrom KJ, Eisenstein EL, et al. Effect of torsemide vs furosemide after discharge on all-cause mortality in patients hospitalized with heart failure: the TRANSFORM-HF randomized clinical trial. *JAMA* 2023;329:214-23.
- 18) Molloy C, Long L, Mordi IR, et al. Exercise-based cardiac rehabilitation for adults with heart failure. *Cochrane Database Syst Rev* 2024;3:CD003331.
- 19) Herrmann JJ, Brunner-La Rocca HP, Baltussen LEHJM, et al. Liberal fluid intake versus fluid restriction in chronic heart failure: a randomized clinical trial. *Nat Med* 2025;31:2062-8.
- 20) Rohde LE, Rover MM, Figueiredo Neto JA, et al. Short-term diuretic withdrawal in stable outpatients with mild heart failure and no fluid retention receiving optimal therapy: a double-blind, multicentre, randomized trial. *Eur Heart J* 2019;40:3605-12.

3-11 がんを疑ってから紹介まで

- 1) National Institute for Health and Care Excellence. Suspected cancer: recognition and referral (NICE guideline NG12). 2015 (2026年4月更新) .
- 2) Hamilton W, Peters TJ, Round A, Sharp D. What are the clinical features of lung cancer before the diagnosis is made? A population based case-control study. *Thorax* 2005;60:1059-65.
- 3) Stapley S, Peters TJ, Neal RD, Rose PW, Walter FM, Hamilton W. The risk of oesophago-gastric cancer in symptomatic patients in primary care: a large case-control study using electronic records. *Br J Cancer* 2013;108:25-31.
- 4) Hamilton W, Round A, Sharp D, Peters TJ. Clinical features of colorectal cancer before diagnosis: a population-based case-control study. *Br J Cancer* 2005;93:399-405.
- 5) Stapley S, Peters TJ, Neal RD, Rose PW, Walter FM, Hamilton W. The risk of pancreatic cancer in symptomatic patients in primary care: a large case-control study using electronic records. *Br J Cancer* 2012;106:1940-4.
- 6) Bailey SE, Ukoumunne OC, Shephard EA, Hamilton W. Clinical relevance of thrombocytosis in primary care: a prospective cohort study of cancer incidence using English electronic medical records and cancer registry data. *Br J Gen Pract* 2017;67:e405-13.
- 7) Hanna TP, King WD, Thibodeau S, et al. Mortality due to cancer treatment delay: systematic review and meta-analysis. *BMJ* 2020;371:m4087.
- 8) Fujimori M, Akechi T, Morita T, et al. Preferences of cancer patients regarding the disclosure of bad news. *Psychooncology* 2007;16:573-81.
- 9) Fujimori M, Shirai Y, Asai M, Kubota K, Katsumata N, Uchitomi Y. Effect of communication skills training program for oncologists based on patient preferences for communication when receiving bad news: a randomized controlled trial. *J Clin Oncol* 2014;32:2166-72.
- 10) 日本サイコオンコロジー学会, 日本がんサポーターブケア学会編. がん医療における患者-医療者間のコミュニケーションガイドライン 2022年版. 金原出版; 2022.
- 11) Yamauchi T, Inagaki M, Yonemoto N, et al. Death by suicide and other externally caused injuries following a cancer diagnosis: the Japan Public Health Center-based Prospective Study. *Psychooncology* 2014;23:1034-41.
- 12) 国立がん研究センターがん対策情報センター. 患者体験調査報告書 平成30年度調査. 2020.
- 13) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. ゼロダ、ティーエスワン) .
- 14) Musallam KM, Tamim HM, Richards T, et al. Preoperative anaemia and postoperative outcomes in non-cardiac surgery: a retrospective cohort study. *Lancet* 2011;378:1396-407.
- 15) Muscaritoli M, Arends J, Bachmann P, et al. ESPEN practical guideline: Clinical Nutrition in cancer. *Clin Nutr* 2021;40:2898-913.
- 16) Mills E, Eyawo O, Lockhart I, Kelly S, Wu P, Ebbert JO. Smoking cessation reduces postoperative complications: a systematic review and meta-analysis. *Am J Med* 2011;124:144-54.e8.
- 17) Temel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733-42.

第4部 腎臓と電解質の外來

4-1 健診の尿異常と腎機能の低下

- 1) 血尿診断ガイドライン改訂委員会編. 血尿診断ガイドライン2023. ライフサイエンス出版; 2023.
- 2) 日本腎臓学会編. エビデンスに基づくCKD診療ガイドライン2023. 東京医学社; 2023.

- 3) Konta T, Hao Z, Takasaki S, et al. Clinical utility of trace proteinuria for microalbuminuria screening in the general population. *Clin Exp Nephrol* 2007;11:51-5.
- 4) 日本腎臓学会編. CKD診療ガイド2024. 東京医学社；2024.
- 5) Iseki K, Ikemiya Y, Iseki C, Takishita S. Proteinuria and the risk of developing end-stage renal disease. *Kidney Int* 2003;63:1468-74.
- 6) Köhler H, Wandel E, Brunck B. Acanthocyturia—a characteristic marker for glomerular bleeding. *Kidney Int* 1991;40:115-20.
- 7) Vivante A, Afek A, Frenkel-Nir Y, et al. Persistent asymptomatic isolated microscopic hematuria in Israeli adolescents and young adults and risk for end-stage renal disease. *JAMA* 2011;306:729-36.
- 8) 厚生労働科学研究費補助金難治性疾患政策研究事業 難治性腎障害に関する調査研究班編. エビデンスに基づく多発性嚢胞腎 (PKD) 診療ガイドライン2020. 東京医学社；2020.
- 9) Imai E, Horio M, Yamagata K, et al. Slower decline of glomerular filtration rate in the Japanese general population: a longitudinal 10-year follow-up study. *Hypertens Res* 2008;31:433-41.
- 10) Corapi KM, Chen JL, Balk EM, Gordon CE. Bleeding complications of native kidney biopsy: a systematic review and meta-analysis. *Am J Kidney Dis* 2012;60:62-73.
- 11) Ginsberg JM, Chang BS, Matarese RA, Garella S. Use of single voided urine samples to estimate quantitative proteinuria. *N Engl J Med* 1983;309:1543-6.
- 12) Liu P, Quinn RR, Lam NN, et al. Accounting for age in the definition of chronic kidney disease. *JAMA Intern Med* 2021;181:1359-66.

4-2 CKDの継続診療

- 1) 日本腎臓学会編. CKD診療ガイド2024. 東京医学社；2024.
- 2) 日本腎臓学会編. エビデンスに基づくCKD診療ガイドライン2023. 東京医学社；2023.
- 3) 日本腎臓学会. CKD治療におけるSGLT2阻害薬の適正使用に関するrecommendation. *日腎会誌* 2023;65:1-10.
- 4) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。ニューロタン、フォシーガ、ジャディアンス、ケレンディア、バルトレックス、リリカ、クラビット、メトグルコ)。
- 5) Brenner BM, Cooper ME, de Zeeuw D, et al. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861-9.
- 6) Heerspink HJL, Stefánsson BV, Correa-Rotter R, et al. Dapagliflozin in patients with chronic kidney disease. *N Engl J Med* 2020;383:1436-46.
- 7) EMPA-KIDNEY Collaborative Group. Empagliflozin in patients with chronic kidney disease. *N Engl J Med* 2023;388:117-27.
- 8) Bakris GL, Agarwal R, Anker SD, et al. Effect of finerenone on chronic kidney disease outcomes in type 2 diabetes. *N Engl J Med* 2020;383:2219-29.
- 9) Perkovic V, Tuttle KR, Rossing P, et al. Effects of semaglutide on chronic kidney disease in patients with type 2 diabetes. *N Engl J Med* 2024;391:109-21.
- 10) Schmidt M, Mansfield KE, Bhaskaran K, et al. Serum creatinine elevation after renin-angiotensin system blockade and long term cardiorenal risks: cohort study. *BMJ* 2017;356:j791.
- 11) Keith DS, Nichols GA, Gullion CM, Brown JB, Smith DH. Longitudinal follow-up and outcomes among a population with chronic kidney disease in a large managed care organization. *Arch Intern Med* 2004;164:659-63.
- 12) 日本腎臓学会 CKD患者の血圧管理に関する検討小委員会. 慢性腎臓病 (CKD) 患者の血圧管理に関するステートメント. *日腎会誌* 2026;68:253-8.

- 13) Baigent C, Landray MJ, Reith C, et al. The effects of lowering LDL cholesterol with simvastatin plus ezetimibe in patients with chronic kidney disease (Study of Heart and Renal Protection): a randomised placebo-controlled trial. *Lancet* 2011;377:2181-92.
- 14) Lapi F, Azoulay L, Yin H, Nessim SJ, Suissa S. Concurrent use of diuretics, angiotensin converting enzyme inhibitors, and angiotensin receptor blockers with non-steroidal anti-inflammatory drugs and risk of acute kidney injury: nested case-control study. *BMJ* 2013;346:e8525.
- 15) Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int* 2024;105(4S):S117-314.
- 16) Cheung AK, Rahman M, Reboussin DM, et al. Effects of intensive BP control in CKD. *J Am Soc Nephrol* 2017;28:2812-23.
- 17) Jongs N, Chertow GM, Greene T, et al. Correlates and consequences of an acute change in eGFR in response to the SGLT2 inhibitor dapagliflozin in patients with CKD. *J Am Soc Nephrol* 2022;33:2094-107.
- 18) Cherney DZ, Perkins BA, Soleymanlou N, et al. Renal hemodynamic effect of sodium-glucose cotransporter 2 inhibition in patients with type 1 diabetes mellitus. *Circulation* 2014;129:587-97.
- 19) van Bommel EJM, Muskiet MHA, van Baar MJB, et al. The renal hemodynamic effects of the SGLT2 inhibitor dapagliflozin are caused by post-glomerular vasodilatation rather than pre-glomerular vasoconstriction in metformin-treated patients with type 2 diabetes in the randomized, double-blind RED trial. *Kidney Int* 2020;97:202-12.

4-3 進行したCKDと腎代替療法の準備

- 1) Smart NA, Dieberg G, Ladhani M, Titus T. Early referral to specialist nephrology services for preventing the progression to end-stage kidney disease. *Cochrane Database Syst Rev* 2014; (6):CD007333.
- 2) 日本腎臓学会編. エビデンスに基づくCKD診療ガイドライン2023. 東京医学社; 2023.
- 3) 日本透析医学会. 慢性腎臓病に伴う骨・ミネラル代謝異常の診療ガイドライン（2025年改訂版）. 透析会誌 2026;59:127-224.
- 4) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ネスブ、ダーブロック、リオナ、ホスレノール、カルタン、アルファロール、炭酸水素ナトリウム、ラシックス）.
- 5) Di Iorio BR, Bellasi A, Raphael KL, et al. Treatment of metabolic acidosis with sodium bicarbonate delays progression of chronic kidney disease: the UBI Study. *J Nephrol* 2019;32:989-1001.
- 6) Bhandari S, Mehta S, Khwaja A, et al. Renin-angiotensin system inhibition in advanced chronic kidney disease. *N Engl J Med* 2022;387:2021-32.
- 7) Tanaka T, Okamura T, Miura K, et al. A simple method to estimate populational 24-h urinary sodium and potassium excretion using a casual urine specimen. *J Hum Hypertens* 2002;16:97-103.
- 8) 日本腎臓学会, 日本透析医学会, 日本腹膜透析医学会, 日本臨床腎移植学会, 日本小児腎臓病学会編. 腎代替療法選択ガイド2020. ライフサイエンス出版; 2020.
- 9) 日本透析医学会. 透析の開始と継続に関する意思決定プロセスについての提言. 透析会誌 2020;53:173-217.
- 10) 日本透析医学会. 維持血液透析ガイドライン：血液透析導入. 透析会誌 2013;46:1107-55.
- 11) 日本透析医学会. 2011年版 慢性血液透析用バスキュラーアクセスの作製および修復に関するガイドライン. 透析会誌 2011;44:855-937.

- 12) 日本臓器移植ネットワーク. 臓器移植解説集 移植希望者の待機年数. <https://www.jotnw.or.jp/explanation/07/05/> (2026年10月閲覧) .
- 13) 全国健康保険協会. 特定疾病に係る高額療養費の特例 (特定疾病療養受療証) . <https://www.kyoukaikenpo.or.jp/> (2026年10月閲覧) .
- 14) 厚生労働省社会・援護局障害保健福祉部企画課長通知. 「身体障害認定基準の取扱い (身体障害認定要領) について」の一部改正について (障企発0226第2号, 平成30年2月26日) .
- 15) Tangri N, Grams ME, Levey AS, et al. Multinational assessment of accuracy of equations for predicting risk of kidney failure: a meta-analysis. *JAMA* 2016;315:164-74.
- 16) Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group. KDIGO 2024 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int* 2024;105(4S):S117-S314.
- 17) Cooper BA, Branley P, Bulfone L, et al. A randomized, controlled trial of early versus late initiation of dialysis. *N Engl J Med* 2010;363:609-19.
- 18) Verberne WR, Geers AB, Jellema WT, Vincent HH, van Delden JJ, Bos WJ. Comparative survival among older adults with advanced kidney disease managed conservatively versus with dialysis. *Clin J Am Soc Nephrol* 2016;11:633-40.
- 19) 日本透析医学会. 血液透析患者の糖尿病治療ガイド2012. 透析会誌 2013;46:311-57.
- 20) El Ters M, Scheers GJ, Taler SJ, et al. Association between prior peripherally inserted central catheters and lack of functioning arteriovenous fistulas: a case-control study in hemodialysis patients. *Am J Kidney Dis* 2012;60:601-8.

4-4 外来で見つかる電解質異常

- 1) UK Kidney Association. Clinical practice guideline: management of hyperkalaemia in adults. Updated July 2026.
- 2) Sevastos N, Theodossiades G, Efstathiou S, Papatheodoridis GV, Manesis E, Archimandritis AJ. Pseudohyperkalemia in serum: the phenomenon and its clinical magnitude. *J Lab Clin Med* 2006;147:139-44.
- 3) Bnaya A, Ganzel C, Shavit L. Pseudohyperkalemia in chronic lymphocytic leukemia: prevalence, impact, and management challenges. *Am J Med Sci* 2023;366:167-75.
- 4) Antoniou T, Gomes T, Juurlink DN, Loutfy MR, Glazier RH, Mamdani MM. Trimethoprim-sulfamethoxazole-induced hyperkalemia in patients receiving inhibitors of the renin-angiotensin system: a population-based study. *Arch Intern Med* 2010;170:1045-9.
- 5) Fralick M, Macdonald EM, Gomes T, et al. Co-trimoxazole and sudden death in patients receiving inhibitors of renin-angiotensin system: population based study. *BMJ* 2014;349:g6196.
- 6) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. ロケルマ、カリメート、塩化カリウム徐放錠、エディロール、マグミット) .
- 7) 厚生労働省. 重篤副作用疾患別対応マニュアル 偽アルドステロン症. 2006 (2022年2月改定) .
- 8) Spasovski G, Vanholder R, Allolio B, et al. Clinical practice guideline on diagnosis and treatment of hyponatraemia. *Eur J Endocrinol* 2014;170:G1-47.
- 9) Andersson NW, Wohlfahrt J, Feenstra B, Hviid A, Melbye M, Lund M. Cumulative incidence of thiazide-induced hyponatremia: a population-based cohort study. *Ann Intern Med* 2024;177:1-11.
- 10) Leth-Møller KB, Hansen AH, Torstensson M, et al. Antidepressants and the risk of hyponatremia: a Danish register-based population study. *BMJ Open* 2016;6:e011200.
- 11) Renneboog B, Musch W, Vandemergel X, Manto MU, Decaux G. Mild chronic hyponatremia is associated with falls, unsteadiness, and attention deficits. *Am J Med* 2006;119:71.e1-8.

- 12) Upala S, Sanguankeo A. Association between hyponatremia, osteoporosis, and fracture: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2016;101:1880-6.
- 13) Walker MD, Shane E. Hypercalcemia: a review. *JAMA* 2022;328:1624-36.
- 14) 骨粗鬆症の予防と治療ガイドライン作成委員会編. 骨粗鬆症の予防と治療ガイドライン2025年版. ライフサイエンス出版; 2025.
- 15) Cheungpasitporn W, Thongprayoon C, Kittanamongkolchai W, et al. Proton pump inhibitors linked to hypomagnesemia: a systematic review and meta-analysis of observational studies. *Ren Fail* 2015;37:1237-41.
- 16) Neal B, Wu Y, Feng X, et al. Effect of salt substitution on cardiovascular events and death. *N Engl J Med* 2021;385:1067-77.
- 17) Garrahy A, Galloway I, Hannon AM, et al. Fluid restriction therapy for chronic SIAD; results of a prospective randomized controlled trial. *J Clin Endocrinol Metab* 2020;105:dga619.
- 18) Krisanapan P, Vongsanin S, Pin-On P, Ruengorn C, Noppakun K. Efficacy of furosemide, oral sodium chloride, and fluid restriction for treatment of syndrome of inappropriate antidiuresis (SIAD): an open-label randomized controlled study (the EFFUSE-FLUID trial). *Am J Kidney Dis* 2020;76:203-12.

第5部 予防・健診と在宅医療

5-1 成人のワクチン

- 1) 予防接種法（昭和23年法律第68号）、予防接種法施行令（昭和23年政令第197号。令和8年9月11日政令第274号による改正まで）。
- 2) 日本呼吸器学会 感染症・結核学術部会ワクチンWG, 日本感染症学会ワクチン委員会, 日本ワクチン学会 合同委員会. 65歳以上の成人に対する肺炎球菌ワクチン接種に関する考え方（第8版）. 2026年4月1日（2026年6月15日一部修正）。
- 3) Kroger A, Bahta L, Long S, Sanchez P. General best practice guidelines for immunization: best practices guidance of the Advisory Committee on Immunization Practices (ACIP). Centers for Disease Control and Prevention（Altered immunocompetence、Special situations、Vaccine administration の章。2026年10月に閲覧）。
- 4) 予防接種法施行規則（昭和23年厚生省令第36号）、予防接種実施規則（昭和33年厚生省令第27号）。
- 5) Demicheli V, Jefferson T, Di Pietrantonj C, et al. Vaccines for preventing influenza in the elderly. *Cochrane Database Syst Rev* 2018;2:CD004876.
- 6) 厚生労働省. 第66回厚生科学審議会予防接種・ワクチン分科会 資料1 高用量インフルエンザワクチンについて. 2026年7月23日.
- 7) Bonten MJ, Huijts SM, Bolkenbaas M, et al. Polysaccharide conjugate vaccine against pneumococcal pneumonia in adults. *N Engl J Med* 2015;372:1114-25.
- 8) Lal H, Cunningham AL, Godeaux O, et al. Efficacy of an adjuvanted herpes zoster subunit vaccine in older adults. *N Engl J Med* 2015;372:2087-96.
- 9) Oxman MN, Levin MJ, Johnson GR, et al. A vaccine to prevent herpes zoster and postherpetic neuralgia in older adults. *N Engl J Med* 2005;352:2271-84.
- 10) 各ワクチンの電子添文（KEGG MEDICUSで2026年10月に確認）。
- 11) Papi A, Ison MG, Langley JM, et al. Respiratory syncytial virus prefusion F protein vaccine in older adults. *N Engl J Med* 2023;388:595-608.
- 12) 日本感染症学会, 日本呼吸器学会, 日本ワクチン学会. 成人のRSウイルスワクチンに関する見解. 2025年12月9日.

- 13) 日本環境感染学会ワクチン委員会. 医療関係者のためのワクチンガイドライン 第5版. 環境感染誌 2026;41(Suppl II):S1-54.
- 14) 厚生労働省検疫所 (FORTH). 黄熱について. <https://www.forth.go.jp/useful/yellowfever.html> (2026年10月閲覧).
- 15) 日本透析医学会「透析施設における標準的な透析操作と感染予防に関するガイドライン」改訂に向けたワーキンググループ. 透析施設における標準的な透析操作と感染予防に関するガイドライン (六訂版). 2023.
- 16) Fröbert O, Götberg M, Erlinge D, et al. Influenza vaccination after myocardial infarction: a randomized, double-blind, placebo-controlled, multicenter trial. *Circulation* 2021;144:1476-84.
- 17) Rubin LG, Levin MJ, Ljungman P, et al. 2013 IDSA clinical practice guideline for vaccination of the immunocompromised host. *Clin Infect Dis* 2014;58:e44-100.
- 18) 厚生労働省. 定期接種実施要領 (「予防接種法第5条第1項の規定による予防接種の実施について」平成25年3月30日健発0330第2号の別添. 令和8年3月31日感発0331第5号による改正まで).
- 19) McNeil MM, Weintraub ES, Duffy J, et al. Risk of anaphylaxis after vaccination in children and adults. *J Allergy Clin Immunol* 2016;137:868-78.
- 20) Morrison VA, Johnson GR, Schmader KE, et al. Long-term persistence of zoster vaccine efficacy. *Clin Infect Dis* 2015;60:900-9.
- 21) Strezova A, Diez-Domingo J, Al Shawafi K, et al. Long-term protection against herpes zoster by the adjuvanted recombinant zoster vaccine: interim efficacy, immunogenicity, and safety results up to 10 years after initial vaccination. *Open Forum Infect Dis* 2022;9:ofac485.
- 22) DiazGranados CA, Dunning AJ, Kimmel M, et al. Efficacy of high-dose versus standard-dose influenza vaccine in older adults. *N Engl J Med* 2014;371:635-45.
- 23) Johansen ND, Modin D, Pardo-Seco J, et al. Effectiveness of high-dose influenza vaccine against hospitalisations in older adults (FLUNITY-HD): an individual-level pooled analysis. *Lancet* 2025;406:2425-34.

5-2 がん検診と健診の結果への対応

- 1) Lee JK, Liles EG, Bent S, Levin TR, Corley DA. Accuracy of fecal immunochemical tests for colorectal cancer: systematic review and meta-analysis. *Ann Intern Med* 2014;160:171.
- 2) 厚生労働省. がん予防重点健康教育及びがん検診実施のための指針 (平成20年3月31日健発第0331058号 厚生労働省健康局長通知別添. 令和7年12月24日一部改正).
- 3) Hewitson P, Glasziou P, Irwig L, Towler B, Watson E. Screening for colorectal cancer using the faecal occult blood test, Hemoccult. *Cochrane Database Syst Rev* 2007;(1):CD001216.
- 4) Independent UK Panel on Breast Cancer Screening. The benefits and harms of breast cancer screening: an independent review. *Lancet* 2012;380:1778-86.
- 5) Hamashima C, Ogoshi K, Okamoto M, Shabana M, Kishimoto T, Fukao A. A community-based, case-control study evaluating mortality reduction from gastric cancer by endoscopic screening in Japan. *PLoS One* 2013;8:e79088.
- 6) Jun JK, Choi KS, Lee HY, et al. Effectiveness of the Korean National Cancer Screening Program in reducing gastric cancer mortality. *Gastroenterology* 2017;152:1319-28.e7.
- 7) 国立がん研究センターがん対策研究所. 有効性評価に基づく肺がん検診ガイドライン2025年度版. 2025.
- 8) Hugosson J, Roobol MJ, Månsson M, et al. A 16-yr follow-up of the European Randomized study of Screening for Prostate Cancer. *Eur Urol* 2019;76:43-51.
- 9) 日本泌尿器科学会 編. 前立腺がん検診ガイドライン 2018年版. メディカルレビュー社, 2018.

- 10) Ohuchi N, Suzuki A, Sobue T, et al. Sensitivity and specificity of mammography and adjunctive ultrasonography to screen for breast cancer in the Japan Strategic Anti-cancer Randomized Trial (J-START): a randomised controlled trial. *Lancet* 2016;387:341-8.
- 11) Kim S, Park BK, Seo JH, et al. Carbohydrate antigen 19-9 elevation without evidence of malignant or pancreatobiliary diseases. *Sci Rep* 2020;10:8820.
- 12) Corley DA, Jensen CD, Quinn VP, et al. Association between time to colonoscopy after a positive fecal test result and risk of colorectal cancer and cancer stage at diagnosis. *JAMA* 2017;317:1631-41.
- 13) 厚生労働省 がん検診のあり方に関する検討会. がん検診事業のあり方について. 2023.
- 14) 厚生労働省. 標準的な健診・保健指導プログラム（令和6年度版）. 第2編第3章 保健指導対象者の選定と階層化、別紙5 健診検査項目の保健指導判定値及び受診勧奨判定値、フィードバック文例集. 2024.
- 15) メタボリックシンドローム診断基準検討委員会. メタボリックシンドロームの定義と診断基準. 日内会誌 2005;94:794-809.
- 16) Knowler WC, Barrett-Connor E, Fowler SE, et al; Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002;346:393-403.
- 17) Saito T, Watanabe M, Nishida J, et al; Zensharen Study for Prevention of Lifestyle Diseases Group. Lifestyle modification and prevention of type 2 diabetes in overweight Japanese with impaired fasting glucose levels: a randomized controlled trial. *Arch Intern Med* 2011;171:1352-60.
- 18) Riley LK, Rupert J. Evaluation of patients with leukocytosis. *Am Fam Physician* 2015;92:1004-11.
- 19) 日本血液学会 編. 造血器腫瘍診療ガイドライン 第3.1版（2024年版）. 第1章 白血病 5 慢性リンパ性白血病／小リンパ球性リンパ腫. 2024.
- 20) Rothwell PM, Eliasziw M, Gutnikov SA, Warlow CP, Barnett HJ; Carotid Endarterectomy Trialists Collaboration. Endarterectomy for symptomatic carotid stenosis in relation to clinical subgroups and timing of surgery. *Lancet* 2004;363:915-24.
- 21) 日本超音波医学会用語・診断基準委員会, 頸動脈超音波診断ガイドライン小委員会. 超音波による頸動脈病変の標準的評価法2017. 2017.
- 22) 日本脳卒中学会 脳卒中ガイドライン委員会. 脳卒中治療ガイドライン2021〔改訂2025〕（改訂項目の公開版）. 2025.
- 23) Brott TG, Howard G, Lal BK, et al. Medical management and revascularization for asymptomatic carotid stenosis. *N Engl J Med* 2026;394:219-31.
- 24) US Preventive Services Task Force. Screening for asymptomatic carotid artery stenosis: US Preventive Services Task Force recommendation statement. *JAMA* 2021;325:476-81.
- 25) Lee SJ, Boscardin WJ, Stijacic-Cenzer I, Conell-Price J, O'Brien S, Walter LC. Time lag to benefit after screening for breast and colorectal cancer: meta-analysis of survival data from the United States, Sweden, United Kingdom, and Denmark. *BMJ* 2013;346:e8441.
- 26) Castells A, Quintero E, Bujanda L, et al. Effect of invitation to colonoscopy versus faecal immunochemical test screening on colorectal cancer mortality (COLONPREV): a pragmatic, randomised, controlled, non-inferiority trial. *Lancet* 2025;405:1231-9.
- 27) Bretthauer M, Løberg M, Wieszczky P, et al. Effect of colonoscopy screening on risks of colorectal cancer and related death. *N Engl J Med* 2022;387:1547-56.

5-3 健診の心電図異常・胸部X線異常

- 1) Yagi R, Mori Y, Goto S, Iwami T, Inoue K. Routine electrocardiogram screening and cardiovascular disease events in adults. *JAMA Intern Med* 2024;184:1035-44.
- 2) 日本人間ドック学会. 標準12誘導心電図検診判定マニュアル（2023年度版）. 2023.

- 3) 日本人間ドック学会. 胸部エックス線健診判定マニュアル.
- 4) 日本循環器学会, 日本不整脈心電学会. 2022年改訂版 不整脈の診断とリスク評価に関するガイドライン. 2022.
- 5) Pewsner D, Jüni P, Egger M, Battaglia M, Sundström J, Bachmann LM. Accuracy of electrocardiography in diagnosis of left ventricular hypertrophy in arterial hypertension: systematic review. *BMJ* 2007;335:711.
- 6) Sharma S, Drezner JA, Baggish A, et al. International recommendations for electrocardiographic interpretation in athletes. *J Am Coll Cardiol* 2017;69:1057-75.
- 7) US Preventive Services Task Force. Screening for cardiovascular disease risk with electrocardiography: US Preventive Services Task Force recommendation statement. *JAMA* 2018;319:2308-14.
- 8) Priori SG, Wilde AA, Horie M, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes. *Europace* 2013;15:1389-406.
- 9) Adler A, Topaz G, Heller K, et al. Fever-induced Brugada pattern: how common is it and what does it mean? *Heart Rhythm* 2013;10:1375-82.
- 10) Mizusawa Y, Morita H, Adler A, et al. Prognostic significance of fever-induced Brugada syndrome. *Heart Rhythm* 2016;13:1515-20.
- 11) Michowitz Y, Milman A, Sarquella-Brugada G, et al. Fever-related arrhythmic events in the multicenter Survey on Arrhythmic Events in Brugada Syndrome. *Heart Rhythm* 2018;15:1394-401.
- 12) Postema PG, Wolpert C, Amin AS, et al. Drugs and Brugada syndrome patients: review of the literature, recommendations, and an up-to-date website (www.brugadadrugs.org). *Heart Rhythm* 2009;6:1335-41 (薬の一覧は同サイトで2026年10月に確認) .
- 13) Obeyesekere MN, Leong-Sit P, Massel D, et al. Risk of arrhythmia and sudden death in patients with asymptomatic preexcitation: a meta-analysis. *Circulation* 2012;125:2308-15.
- 14) 日本肺癌学会肺がん検診委員会編. 肺がん検診の手引き 2020年改訂版.
- 15) National Lung Screening Trial Research Team. Results of initial low-dose computed tomographic screening for lung cancer. *N Engl J Med* 2013;368:1980-91.
- 16) Denning DW, Cadranel J, Beigelman-Aubry C, et al. Chronic pulmonary aspergillosis: rationale and clinical guidelines for diagnosis and management. *Eur Respir J* 2016;47:45-68.
- 17) Badgett RG, Mulrow CD, Otto PM, Ramírez G. How well can the chest radiograph diagnose left ventricular dysfunction? *J Gen Intern Med* 1996;11:625-34.
- 18) Imazio M, Adler Y. Management of pericardial effusion. *Eur Heart J* 2013;34:1186-97.
- 19) Oken MM, Hocking WG, Kvale PA, et al. Screening by chest radiograph and lung cancer mortality: the Prostate, Lung, Colorectal, and Ovarian (PLCO) randomized trial. *JAMA* 2011;306:1865-73.
- 20) Sennberg E, Friberg L, Frykman V, Al-Khalili F, Engdahl J, Rosenqvist M. Clinical outcomes in systematic screening for atrial fibrillation (STROKESTOP): a multicentre, parallel group, unmasked, randomised controlled trial. *Lancet* 2021;398:1498-506.
- 21) Svendsen JH, Diederichsen SZ, Højberg S, et al. Implantable loop recorder detection of atrial fibrillation to prevent stroke (The LOOP Study): a randomised controlled trial. *Lancet* 2021;398:1507-16.
- 22) US Preventive Services Task Force. Screening for atrial fibrillation: US Preventive Services Task Force recommendation statement. *JAMA* 2022;327:360-7.

5-4 禁煙と節酒

- 1) Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ* 2004;328:1519.
- 2) Stead LF, Buitrago D, Preciado N, Sanchez G, Hartmann-Boyce J, Lancaster T. Physician advice for smoking cessation. *Cochrane Database Syst Rev* 2013;(5):CD000165.
- 3) Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: a systematic review. *JAMA* 2003;290:86-97.
- 4) 日本循環器学会, 日本肺癌学会, 日本癌学会, 日本呼吸器学会. 禁煙治療のための標準手順書 第8.1版. 2021年9月.
- 5) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. チャンピックス錠、ニコチネルTTS、セリシクロ錠、レグテクト錠、メトグルコ錠) .
- 6) Livingstone-Banks J, Fanshawe TR, Thomas KH, et al. Nicotine receptor partial agonists for smoking cessation. *Cochrane Database Syst Rev* 2023;5:CD006103.
- 7) Hartmann-Boyce J, Chepkin SC, Ye W, Bullen C, Lancaster T. Nicotine replacement therapy versus control for smoking cessation. *Cochrane Database Syst Rev* 2018;5:CD000146.
- 8) ファイザー株式会社. チャンピックス錠 通常出荷再開のご案内. 2025年10月.
- 9) Anthenelli RM, Benowitz NL, West R, et al. Neuropsychiatric safety and efficacy of varenicline, bupropion, and nicotine patch in smokers with and without psychiatric disorders (EAGLES): a double-blind, randomised, placebo-controlled clinical trial. *Lancet* 2016;387:2507-20.
- 10) Hughes JR. Effects of abstinence from tobacco: valid symptoms and time course. *Nicotine Tob Res* 2007;9:315-27.
- 11) Aubin HJ, Farley A, Lycett D, Lahmek P, Aveyard P. Weight gain in smokers after quitting cigarettes: meta-analysis. *BMJ* 2012;345:e4439.
- 12) 厚生労働省. 健康に配慮した飲酒に関するガイドライン. 2024年2月.
- 13) 厚生労働省健康局. 標準的な健診・保健指導プログラム (令和6年度版). 2024年4月.
- 14) Kaner EF, Beyer FR, Muirhead C, et al. Effectiveness of brief alcohol interventions in primary care populations. *Cochrane Database Syst Rev* 2018;2:CD004148.
- 15) Miyata H, Takahashi M, Murai Y, et al. Nalmefene in alcohol-dependent patients with a high drinking risk: randomized controlled trial. *Psychiatry Clin Neurosci* 2019;73:697-706.
- 16) 松本俊彦, ほか. 処方薬・市販薬依存症患者の実態と通院治療プログラムの開発に関する研究. 令和7年度厚生労働科学研究費補助金 (障害者政策総合研究事業) 分担研究報告書 (全国精神科医療施設における薬物関連精神疾患の実態調査 2024年の二次解析) .
- 17) 厚生労働省. 濫用等のおそれのある医薬品 (医薬品医療機器等法施行規則第15条の2の規定に基づき厚生労働大臣が指定する医薬品. 令和5年厚生労働省告示第5号, 令和5年4月1日適用) .
- 18) Smith PC, Schmidt SM, Allensworth-Davies D, Saitz R. A single-question screening test for drug use in primary care. *Arch Intern Med* 2010;170:1155-60.
- 19) Johnson EE, Hamer R, Nora RM, Tan B, Eisenstein N, Engelhart C. The Lie/Bet Questionnaire for screening pathological gamblers. *Psychol Rep* 1997;80:83-8.
- 20) Johnson EE, Hamer RM, Nora RM. The Lie/Bet Questionnaire for screening pathological gamblers: a follow-up study. *Psychol Rep* 1998;83:1219-24.
- 21) 国立病院機構久里浜医療センター. 令和5年度 依存症に関する調査研究事業「ギャンブル障害およびギャンブル関連問題の実態調査」報告書. 2024年10月.
- 22) 厚生労働省. 依存症対策. <https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/0000070789.html> (2026年10月閲覧) .

- 23) 厚生労働省. 特掲診療料の施設基準等及びその届出に関する手続きの取扱いについて. 別添2 様式8の2 「ニコチン依存症管理料に係る報告書」記載上の注意 (四国厚生支局のページに載る令和7年8月報告用の様式. 2026年10月閲覧).
- 24) Zhao J, Stockwell T, Naimi T, Churchill S, Clay J, Sherk A. Association between daily alcohol intake and risk of all-cause mortality: a systematic review and meta-analyses. *JAMA Netw Open* 2023;6:e236185.
- 25) Hackshaw A, Morris JK, Boniface S, Tang JL, Milenkovic D. Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports. *BMJ* 2018;360:j5855.
- 26) Lindson-Hawley N, Banting M, West R, Michie S, Shinkins B, Aveyard P. Gradual versus abrupt smoking cessation: a randomized, controlled noninferiority trial. *Ann Intern Med* 2016;164:585-92.

5-5 訪問診療と在宅医療

- 1) 厚生労働省. 人生の最終段階における医療・ケアに関する意識調査 報告書 (令和4年度調査). 2023.
- 2) 厚生労働省. 令和5年 (2023) 人口動態統計月報年計 (概数) の概況. 2024.
- 3) 厚生労働省保険局医療課長. 診療報酬の算定方法の一部改正に伴う実施上の留意事項について (令和8年3月5日 保医発0305第6号). 別添1 第2章第2部 在宅医療 (C000 往診料、C001 在宅患者訪問診療料 (I)).
- 4) Gomes B, Calanzani N, Curiale V, McCrone P, Higginson IJ. Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers. *Cochrane Database Syst Rev* 2013;(6):CD007760.
- 5) Stall N, Nowaczynski M, Sinha SK. Systematic review of outcomes from home-based primary care programs for homebound older adults. *J Am Geriatr Soc* 2014;62:2243-51.
- 6) Hamano J, Yamaguchi T, Maeda I, et al. Multicenter cohort study on the survival time of cancer patients dying at home or in a hospital: does place matter? *Cancer* 2016;122:1453-60.
- 7) Schulz R, Beach SR. Caregiving as a risk factor for mortality: the Caregiver Health Effects Study. *JAMA* 1999;282:2215-9.
- 8) 厚生労働省保険局医療課長. 特別養護老人ホーム等における療養の給付の取扱いについて (平成18年3月31日 保医発第0331002号. その後の改正を含む).
- 9) 介護保険法 (平成9年法律第123号). 第8条第6項、第27条.
- 10) Kuroda K, Miura T, Kuroiwa S, Kuroda M, Kobayashi N, Kita K. What are the factors that cause emergency home visit in home medical care in Japan? *J Gen Fam Med* 2021;22:81-6.
- 11) Yokobayashi K, Matsushima M, Watanabe T, Fujinuma Y, Tazuma S. Prospective cohort study of fever incidence and risk in elderly persons living at home. *BMJ Open* 2014;4:e004998.
- 12) ロセフィン静注用 電子添文 (KEGG MEDICUSで2026年10月に確認).
- 13) 日本臨床救急医学会. 人生の最終段階にある傷病者の意思に沿った救急現場での心肺蘇生等のあり方に関する提言. 2017.
- 14) 医師法 (昭和23年法律第201号). 第20条.
- 15) 厚生労働省医政局医事課長. 医師法第20条ただし書の適切な運用について (医政医発0831第1号). 2012.
- 16) Downar J, Goldman R, Pinto R, Englesakis M, Adhikari NK. The "surprise question" for predicting death in seriously ill patients: a systematic review and meta-analysis. *CMAJ* 2017;189:E484-93.

5-6 海外渡航の前と後

- 1) Leder K, Torresi J, Libman MD, et al. GeoSentinel surveillance of illness in returned travelers, 2007-2011. *Ann Intern Med* 2013;158:456-68.
- 2) 厚生労働省検疫所（FORTH）．海外渡航のためのワクチン（予防接種）．<https://www.forth.go.jp/useful/vaccination.html>（2026年10月閲覧）．
- 3) 各ワクチン・薬剤の電子添文（KEGG MEDICUSで2026年10月に確認．エイムゲン、ラビビュール筋注用、ジェービックV、タイフィム ブイアイ、メンクアッドフィ筋注、黄熱ワクチン、マラロン配合錠、メファキン「ヒサミツ」錠275、ピブラマイシン錠、ロペミン、ジスロマック錠、クラビット錠、ダイアモックス、メラトベル、フラジール内服錠、ストロメクトール錠）．
- 4) 厚生労働省検疫所（FORTH）．黄熱について．<https://www.forth.go.jp/useful/yellowfever.html>（2026年10月閲覧）．
- 5) Centers for Disease Control and Prevention. CDC Yellow Book 2026: health information for international travel (Typhoid and paratyphoid fever, Meningococcal disease, Malaria, Mosquitoes, ticks, and other arthropods, Travelers' diarrhea, High-altitude travel and altitude illness, Deep vein thrombosis and pulmonary embolism, Jet lag disorder, Post-travel evaluation of the ill traveler, Leptospirosis, Post-travel dermatologic conditions) の章. 2026年10月に閲覧）．
- 6) 日本医療研究開発機構「わが国における輸入熱帯病・まれな寄生虫症の診断治療体制の整備」研究班. マラリア 診断・治療・予防の手引き 第6版. 2026.
- 7) 厚生労働省医薬・生活衛生局審査管理課長. 防除用医薬品及び防除用医薬部外品の製造販売承認申請に係る手続きについて（平成28年6月15日 薬生審査発0615第1号）．
- 8) Riddle MS, Connor BA, Beeching NJ, et al. Guidelines for the prevention and treatment of travelers' diarrhea: a graded expert panel report. *J Travel Med* 2017;24(suppl 1):S57-74.
- 9) Ritchie ND, Baggott AV, Andrew Todd WT. Acetazolamide for the prevention of acute mountain sickness--a systematic review and meta-analysis. *J Travel Med* 2012;19:298-307.
- 10) Kuipers S, Cannegieter SC, Middeldorp S, Robyn L, Büller HR, Rosendaal FR. The absolute risk of venous thrombosis after air travel: a cohort study of 8,755 employees of international organisations. *PLoS Med* 2007;4:e290.
- 11) Schünemann HJ, Cushman M, Burnett AE, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: prophylaxis for hospitalized and nonhospitalized medical patients. *Blood Adv* 2018;2:3198-225.
- 12) Coker RK, Armstrong A, Church AC, et al. BTS Clinical Statement on air travel for passengers with respiratory disease. *Thorax* 2022;77:329-50.
- 13) 厚生労働省 地方厚生局麻薬取締部. 麻薬・向精神薬を携帯して出入国する方へ（関東信越厚生局のページ. 2026年10月閲覧）．
- 14) Pinsker JE, Becker E, Mahnke CB, Ching M, Larson NS, Roy D. Extensive clinical experience: a simple guide to basal insulin adjustments for long-distance travel. *J Diabetes Metab Disord* 2013;12:59.
- 15) ACOG Committee Opinion No. 746: air travel during pregnancy. *Obstet Gynecol* 2018;132:e64-6.
- 16) Kantele A, Lääveri T, Mero S, et al. Antimicrobials increase travelers' risk of colonization by extended-spectrum betalactamase-producing Enterobacteriaceae. *Clin Infect Dis* 2015;60:837-46.
- 17) Clarke MJ, Broderick C, Hopewell S, Juszczak E, Eisinga A. Compression stockings for preventing deep vein thrombosis in airline passengers. *Cochrane Database Syst Rev* 2021;4:CD004002.

内科大全 外来編

よくある訴えと慢性疾患

2026年10月07日23:19 下書き（入稿用ではない）

本書に記載した薬剤の用量・用法、診断・治療の基準は、発行時点の電子添文とガイドラインに基づく。実際の使用にあたっては、最新の電子添文とガイドラインを確認すること。

本書の内容の無断での複製・転載を禁じる。