

救急外来と急変

内科大全救急編 文献

下書き 2026-10-07 23:03

2026年9月

文献

本文の右肩の番号 (1)2) …) は、章ごとの文献番号である。章の番号と題は本文と同じにしてある。

第1部 初期評価と蘇生

1-1 初期評価

- 1) Fieselmann JF, Hendryx MS, Helms CM, et al. Respiratory rate predicts cardiopulmonary arrest for internal medicine inpatients. *J Gen Intern Med* 1993;8:354-60.
- 2) Royal College of Physicians. National Early Warning Score (NEWS) 2: Standardising the assessment of acute-illness severity in the NHS. London: RCP; 2017.
- 3) 日本版敗血症診療ガイドライン2024特別委員会. 日本版敗血症診療ガイドライン2024. 日集中医誌 2024;31:S1165-313.
- 4) Seymour CW, Liu VX, Iwashyna TJ, et al. Assessment of clinical criteria for sepsis: for the Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* 2016;315:762-74.
- 5) Kristensen AK, Holler JG, Hallas J, et al. Is shock index a valid predictor of mortality in emergency department patients with hypertension, diabetes, high age, or receipt of β - or calcium channel blockers? *Ann Emerg Med* 2016;67:106-13.
- 6) Karjalainen J, Viitasalo M. Fever and cardiac rhythm. *Arch Intern Med* 1986;146:1169-71.
- 7) Deakin CD, Low JL. Accuracy of the advanced trauma life support guidelines for predicting systolic blood pressure using carotid, femoral, and radial pulses: observational study. *BMJ* 2000;321:673-4.
- 8) Lehman LW, Saeed M, Talmor D, et al. Methods of blood pressure measurement in the ICU. *Crit Care Med* 2013;41:34-40.
- 9) Niven DJ, Gaudet JE, Laupland KB, et al. Accuracy of peripheral thermometers for estimating temperature: a systematic review and meta-analysis. *Ann Intern Med* 2015;163:768-77.
- 10) 日本救急医学会, 日本救急看護学会, 日本小児救急医学会, 日本臨床救急医学会, 日本在宅救急医学会 監修. 緊急度判定支援システム JTAS2023ガイドブック. へるす出版; 2023.
- 11) 日本循環器学会. 急性冠症候群ガイドライン (2018年改訂版). 2019.
- 12) 厚生労働省 政策科学推進研究事業「外国人患者の受入環境整備に関する研究」研究班. 外国人患者の受入れのための医療機関向けマニュアル (第4.0版). 2021.
- 13) Gabayan GZ, Gould MK, Weiss RE, et al. Emergency department vital signs and outcomes after discharge. *Acad Emerg Med* 2017;24:846-54.
- 14) Obermeyer Z, Cohn B, Wilson M, et al. Early death after discharge from emergency departments: analysis of national US insurance claims data. *BMJ* 2017;356:j239.
- 15) Engel KG, Heisler M, Smith DM, et al. Patient comprehension of emergency department care and instructions: are patients aware of when they do not understand? *Ann Emerg Med* 2009;53:454-61.
- 16) Newman-Toker DE, Peterson SM, Badihian S, et al. Diagnostic errors in the emergency department: a systematic review. Comparative Effectiveness Review No. 258. Rockville (MD): Agency for Healthcare Research and Quality; 2022.

- 17) Westbrook JI, Raban MZ, Walter SR, et al. Task errors by emergency physicians are associated with interruptions, multitasking, fatigue and working memory capacity: a prospective, direct observation study. *BMJ Qual Saf* 2018;27:655-63.
- 18) Westbrook JI, Woods A, Rob MI, et al. Association of interruptions with an increased risk and severity of medication administration errors. *Arch Intern Med* 2010;170:683-90.

1-2 心停止と心拍再開後の管理

- 1) Perkins GD, Ji C, Deakin CD, et al. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med* 2018;379:711-21.
- 2) Kudenchuk PJ, Brown SP, Daya M, et al. Amiodarone, lidocaine, or placebo in out-of-hospital cardiac arrest. *N Engl J Med* 2016;374:1711-22.
- 3) 日本蘇生協議会監修. JRC蘇生ガイドライン2025. 医学書院；2026.
- 4) 日本蘇生協議会監修. JRC蘇生ガイドライン2020. 医学書院；2021.
- 5) Cheskes S, Verbeek PR, Drennan IR, et al. Defibrillation strategies for refractory ventricular fibrillation. *N Engl J Med* 2022;387:1947-56.
- 6) Resuscitation Council UK. 2025 Resuscitation Guidelines: Adult advanced life support guidelines. 2025.
- 7) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）.
- 8) 社会保険診療報酬支払基金. 審査情報提供事例（薬剤）49 アドレナリン①（循環器科2）. 2007（2014年更新）.
- 9) Donnino MW, Saliccioli JD, Howell MD, et al. Time to administration of epinephrine and outcome after in-hospital cardiac arrest with non-shockable rhythms: retrospective analysis of large in-hospital data registry. *BMJ* 2014;348:g3028.
- 10) Soar J, Böttiger BW, Carli P, et al. European Resuscitation Council Guidelines 2025: adult advanced life support. *Resuscitation* 2025;215:110769.
- 11) Andersen LW, Kurth T, Chase M, et al. Early administration of epinephrine (adrenaline) in patients with cardiac arrest with initial shockable rhythm in hospital: propensity score matched analysis. *BMJ* 2016;353:i1577.
- 12) Vallentin MF, Granfeldt A, Meilandt C, et al. Effect of intravenous or intraosseous calcium vs saline on return of spontaneous circulation in adults with out-of-hospital cardiac arrest: a randomized clinical trial. *JAMA* 2021;326:2268-76.
- 13) Couper K, Ji C, Deakin CD, et al. A randomized trial of drug route in out-of-hospital cardiac arrest. *N Engl J Med* 2025;392:336-48.
- 14) Andersen LW, Granfeldt A, Callaway CW, et al. Association between tracheal intubation during adult in-hospital cardiac arrest and survival. *JAMA* 2017;317:494-506.
- 15) Huis In 't Veld MA, Allison MG, Bostick DS, et al. Ultrasound use during cardiopulmonary resuscitation is associated with delays in chest compressions. *Resuscitation* 2017;119:95-8.
- 16) Inoue A, Hifumi T, Sakamoto T, et al. Extracorporeal cardiopulmonary resuscitation in adult patients with out-of-hospital cardiac arrest: a retrospective large cohort multicenter study in Japan. *Crit Care* 2022;26:129.
- 17) Inamasu J, Miyatake S, Tomioka H, et al. Subarachnoid haemorrhage as a cause of out-of-hospital cardiac arrest: a prospective computed tomography study. *Resuscitation* 2009;80:977-80.
- 18) Baldi E, Schnaubelt S, Caputo ML, et al. Association of timing of electrocardiogram acquisition after return of spontaneous circulation with coronary angiography findings in patients with out-of-hospital cardiac arrest. *JAMA Netw Open* 2021;4:e2032875.

- 19) Schmidt H, Kjaergaard J, Hassager C, et al. Oxygen targets in comatose survivors of cardiac arrest. *N Engl J Med* 2022;387:1467-76.
- 20) Eastwood G, Nichol AD, Hodgson C, et al. Mild hypercapnia or normocapnia after out-of-hospital cardiac arrest. *N Engl J Med* 2023;389:45-57.
- 21) Kjaergaard J, Møller JE, Schmidt H, et al. Blood-pressure targets in comatose survivors of cardiac arrest. *N Engl J Med* 2022;387:1456-66.
- 22) Dankiewicz J, Cronberg T, Lilja G, et al. Hypothermia versus normothermia after out-of-hospital cardiac arrest. *N Engl J Med* 2021;384:2283-94.
- 23) Lemkes JS, Janssens GN, van der Hoeven NW, et al. Coronary angiography after cardiac arrest without ST-segment elevation. *N Engl J Med* 2019;380:1397-407.
- 24) Desch S, Freund A, Akin I, et al. Angiography after out-of-hospital cardiac arrest without ST-segment elevation. *N Engl J Med* 2021;385:2544-53.
- 25) Nolan JP, Sandroni C, Böttiger BW, et al. European Resuscitation Council and European Society of Intensive Care Medicine guidelines 2021: post-resuscitation care. *Intensive Care Med* 2021;47:369-421.
- 26) Levine RL, Wayne MA, Miller CC. End-tidal carbon dioxide and outcome of out-of-hospital cardiac arrest. *N Engl J Med* 1997;337:301-6.
- 27) Goto Y, Maeda T, Goto YN. Termination-of-resuscitation rule for emergency department physicians treating out-of-hospital cardiac arrest patients: an observational cohort study. *Crit Care* 2013;17:R235.
- 28) 厚生労働省医政局, 厚生労働省政策統括官(統計・情報システム管理、労使関係担当). 死亡診断書(死体検案書)記入マニュアル 令和8年度版. 2026.
- 29) 医師法(昭和23年法律第201号)第21条.
- 30) Jabre P, Belpomme V, Azoulay E, et al. Family presence during cardiopulmonary resuscitation. *N Engl J Med* 2013;368:1008-18.
- 31) Goldberger ZD, Chan PS, Berg RA, et al. Duration of resuscitation efforts and survival after in-hospital cardiac arrest: an observational study. *Lancet* 2012;380:1473-81.
- 32) Okubo M, Komukai S, Andersen LW, et al. Duration of cardiopulmonary resuscitation and outcomes for adults with in-hospital cardiac arrest: retrospective cohort study. *BMJ* 2024;384:e076019.

1-3 ショック・血圧低下

- 1) Vincent JL, De Backer D. Circulatory shock. *N Engl J Med* 2013;369:1726-34.
- 2) De Backer D, Biston P, Devriendt J, et al. Comparison of dopamine and norepinephrine in the treatment of shock. *N Engl J Med* 2010;362:779-89.
- 3) 日本版敗血症診療ガイドライン2024特別委員会. 日本版敗血症診療ガイドライン2024. 日集中医誌 2024;31:S1165-313.
- 4) 日本循環器学会, 日本心不全学会. 2025年改訂版 心不全診療ガイドライン. 2025.
- 5) ANDROMEDA-SHOCK-2 Investigators. Personalized hemodynamic resuscitation targeting capillary refill time in early septic shock: the ANDROMEDA-SHOCK-2 randomized clinical trial. *JAMA* 2025;334:1988-99.
- 6) McGee S, Abernethy WB 3rd, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA* 1999;281:1022-9.
- 7) Stevenson LW, Perloff JK. The limited reliability of physical signs for estimating hemodynamics in chronic heart failure. *JAMA* 1989;261:884-8.
- 8) Mizuguchi Y, Mouri H, Jo T, et al. Clinical features and outcomes of Shoshin beriberi. *Int Heart J* 2024;65:271-8.

- 9) Lei Y, Zheng MH, Huang W, Zhang J, Lu Y. Wet beriberi with multiple organ failure remarkably reversed by thiamine administration: a case report and literature review. *Medicine (Baltimore)* 2018;97:e0010.
- 10) Stickles SP, Carpenter CR, Gekle R, et al. The diagnostic accuracy of a point-of-care ultrasound protocol for shock etiology: a systematic review and meta-analysis. *CJEM* 2019;21:406-17.
- 11) Atkinson PR, Milne J, Diegelmann L, et al. Does point-of-care ultrasonography improve clinical outcomes in emergency department patients with undifferentiated hypotension? An international randomized controlled trial from the SHoC-ED investigators. *Ann Emerg Med* 2018;72:478-89.
- 12) Roy CL, Minor MA, Brookhart MA, Choudhry NK. Does this patient with a pericardial effusion have cardiac tamponade? *JAMA* 2007;297:1810-8.
- 13) Shapiro NI, Douglas IS, Brower RG, et al. Early restrictive or liberal fluid management for sepsis-induced hypotension. *N Engl J Med* 2023;388:499-510.
- 14) Semler MW, Self WH, Wanderer JP, et al. Balanced crystalloids versus saline in critically ill adults. *N Engl J Med* 2018;378:829-39.
- 15) Asfar P, Meziani F, Hamel JF, et al. High versus low blood-pressure target in patients with septic shock. *N Engl J Med* 2014;370:1583-93.
- 16) Lamontagne F, Richards-Belle A, Thomas K, et al. Effect of reduced exposure to vasopressors on 90-day mortality in older critically ill patients with vasodilatory hypotension: a randomized clinical trial. *JAMA* 2020;323:938-49.
- 17) Permpikul C, Tongyoo S, Viarasilpa T, et al. Early use of norepinephrine in septic shock resuscitation (CENSER): a randomized trial. *Am J Respir Crit Care Med* 2019;199:1097-105.
- 18) Evans L, Rhodes A, Alhazzani W, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021. *Intensive Care Med* 2021;47:1181-247.
- 19) Muller G, Contou D, Ehrmann S, et al. Deferring arterial catheterization in critically ill patients with shock. *N Engl J Med* 2025;393:1875-88.
- 20) Prescott HC, Antonelli M, Alhazzani W, et al. Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock 2026. *Crit Care Med* 2026;54:725-812.
- 21) Russell JA, Walley KR, Singer J, et al. Vasopressin versus norepinephrine infusion in patients with septic shock. *N Engl J Med* 2008;358:877-87.
- 22) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）。
- 23) Hochman JS, Sleeper LA, Webb JG, et al. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. *N Engl J Med* 1999;341:625-34.
- 24) Levy B, Clere-Jehl R, Legras A, et al. Epinephrine versus norepinephrine for cardiogenic shock after acute myocardial infarction. *J Am Coll Cardiol* 2018;72:173-82.
- 25) Thiele H, Zeymer U, Neumann FJ, et al. Intraaortic balloon support for myocardial infarction with cardiogenic shock. *N Engl J Med* 2012;367:1287-96.
- 26) Thiele H, Zeymer U, Akin I, et al. Extracorporeal life support in infarct-related cardiogenic shock. *N Engl J Med* 2023;389:1286-97.
- 27) Möller JE, Engstrøm T, Jensen LO, et al. Microaxial flow pump or standard care in infarct-related cardiogenic shock. *N Engl J Med* 2024;390:1382-93.
- 28) Jones AE, Yiannibas V, Johnson C, et al. Emergency department hypotension predicts sudden unexpected in-hospital mortality: a prospective cohort study. *Chest* 2006;130:941-6.
- 29) Rivers E, Nguyen B, Havstad S, et al. Early goal-directed therapy in the treatment of severe sepsis and septic shock. *N Engl J Med* 2001;345:1368-77.
- 30) PRISM Investigators. Early, goal-directed therapy for septic shock: a patient-level meta-analysis. *N Engl J Med* 2017;376:2223-34.

- 31) Jansen TC, van Bommel J, Schoonderbeek FJ, et al. Early lactate-guided therapy in intensive care unit patients: a multicenter, open-label, randomized controlled trial. *Am J Respir Crit Care Med* 2010;182:752-61.
- 32) Jones AE, Shapiro NI, Trzeciak S, et al. Lactate clearance vs central venous oxygen saturation as goals of early sepsis therapy: a randomized clinical trial. *JAMA* 2010;303:739-46.
- 33) Hernández G, Ospina-Tascón GA, Damiani LP, et al. Effect of a resuscitation strategy targeting peripheral perfusion status vs serum lactate levels on 28-day mortality among patients with septic shock: the ANDROMEDA-SHOCK randomized clinical trial. *JAMA* 2019;321:654-64.
- 34) Levy B, Gibot S, Franck P, et al. Relation between muscle Na⁺K⁺ ATPase activity and raised lactate concentrations in septic shock: a prospective study. *Lancet* 2005;365:871-5.

1-4 院内急変（RRSと院内心停止）

- 1) Schein RM, Hazday N, Pena M, et al. Clinical antecedents to in-hospital cardiopulmonary arrest. *Chest* 1990;98:1388-92.
- 2) Yokoyama H, Yonemoto N, Yonezawa K, et al. Report from the Japanese registry of CPR for in-hospital cardiac arrest (J-RCPR). *Circ J* 2011;75:815-22.
- 3) Royal College of Physicians. National Early Warning Score (NEWS) 2: standardising the assessment of acute-illness severity in the NHS. RCP; 2017.
- 4) Chan PS, Jain R, Nallmothu BK, et al. Rapid response teams: a systematic review and meta-analysis. *Arch Intern Med* 2010;170:18-26.
- 5) Barwise A, Thongprayoon C, Gajic O, et al. Delayed rapid response team activation is associated with increased hospital mortality, morbidity, and length of stay in a tertiary care institution. *Crit Care Med* 2016;44:54-63.
- 6) Hillman K, Chen J, Cretikos M, et al. Introduction of the medical emergency team (MET) system: a cluster-randomised controlled trial. *Lancet* 2005;365:2091-7.
- 7) Naito T, Fujiwara S, Kawasaki T, et al. First report based on the online registry of a Japanese multicenter rapid response system: a descriptive study of 35 institutions in Japan. *Acute Med Surg* 2020;7:e454.
- 8) 日本蘇生協議会監修. JRC蘇生ガイドライン2025. 医学書院；2026.
- 9) Chan PS, Krumholz HM, Nichol G, et al. Delayed time to defibrillation after in-hospital cardiac arrest. *N Engl J Med* 2008;358:9-17.
- 10) Fu ES, Downs JB, Schweiger JW, et al. Supplemental oxygen impairs detection of hypoventilation by pulse oximetry. *Chest* 2004;126:1552-8.
- 11) Joffe AM, Hetzel S, Liew EC. A two-handed jaw-thrust technique is superior to the one-handed "EC-clamp" technique for mask ventilation in the apneic unconscious person. *Anesthesiology* 2010;113:873-9.
- 12) 日本麻酔科学会. 日本麻酔科学会気道管理ガイドライン2014（日本語訳）：より安全な麻酔導入のために. 2015.
- 13) Habrat D. How to insert an oropharyngeal airway. MSD Manual Professional Version. Merck & Co.（2026年10月に閲覧）.
- 14) Habrat D. How to insert a nasopharyngeal airway. MSD Manual Professional Version. Merck & Co.（2026年10月に閲覧）.
- 15) Langeron O, Masso E, Huraux C, et al. Prediction of difficult mask ventilation. *Anesthesiology* 2000;92:1229-36.
- 16) Cook TM, Woodall N, Harper J, et al. Major complications of airway management in the UK: results of the Fourth National Audit Project of the Royal College of Anaesthetists and the

- Difficult Airway Society. Part 2: intensive care and emergency departments. *Br J Anaesth* 2011;106:632-42.
- 17) 西村匡司, 丸藤哲, 日本集中治療医学会倫理委員会. Do Not Attempt Resuscitation (DNAR) 指示のあり方についての勧告. *日集中医誌* 2017;24:208-9.
 - 18) Fernando SM, Tran A, Cheng W, et al. Pre-arrest and intra-arrest prognostic factors associated with survival after in-hospital cardiac arrest: systematic review and meta-analysis. *BMJ* 2019;367:l6373.
 - 19) Jones DA, Bagshaw SM, Barrett J, et al. The role of the medical emergency team in end-of-life care: a multicenter, prospective, observational study. *Crit Care Med* 2012;40:98-103.
 - 20) 日本蘇生協議会監修. JRC蘇生ガイドライン2020. 医学書院; 2021.
 - 21) Edelson DP, Litzinger B, Arora V, et al. Improving in-hospital cardiac arrest process and outcomes with performance debriefing. *Arch Intern Med* 2008;168:1063-9.
 - 22) Peberdy MA, Ornato JP, Larkin GL, et al. Survival from in-hospital cardiac arrest during nights and weekends. *JAMA* 2008;299:785-92.

1-5 人工呼吸の基礎

- 1) Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000;342:1301-8.
- 2) 日本集中治療医学会, 日本呼吸器学会, 日本呼吸療法医学会. ARDS診療ガイドライン2021. *日集中医誌* 2022;29:295-332.
- 3) Amato MBP, Meade MO, Slutsky AS, et al. Driving pressure and survival in the acute respiratory distress syndrome. *N Engl J Med* 2015;372:747-55.
- 4) ARDS Definition Task Force. Acute respiratory distress syndrome: the Berlin Definition. *JAMA* 2012;307:2526-33.
- 5) Guérin C, Reignier J, Richard JC, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159-68.
- 6) National Heart, Lung, and Blood Institute PETAL Clinical Trials Network. Early neuromuscular blockade in the acute respiratory distress syndrome. *N Engl J Med* 2019;380:1997-2008.
- 7) Combes A, Hajage D, Capellier G, et al. Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med* 2018;378:1965-75.
- 8) 日本集中治療医学会J-PADガイドライン作成委員会. 日本版・集中治療室における成人重症患者に対する痛み・不穏・せん妄管理のための臨床ガイドライン. *日集中医誌* 2014;21:539-79.
- 9) Devlin JW, Skrobik Y, Gélinas C, et al. Clinical practice guidelines for the prevention and management of pain, agitation/sedation, delirium, immobility, and sleep disruption in adults in the ICU. *Crit Care Med* 2018;46:e825-73.
- 10) Hughes CG, Mailloux PT, Devlin JW, et al. Dexmedetomidine or propofol for sedation in mechanically ventilated adults with sepsis. *N Engl J Med* 2021;384:1424-36.
- 11) 各薬剤の電子添文（フェンタニル注射液、ディプリバン注、プレセデックス静注液。2026年10月に確認）。
- 12) Fuller BM, Pappal RD, Mohr NM, et al. Awareness with paralysis among critically ill emergency department patients: a prospective cohort study. *Crit Care Med* 2022;50:1449-60.
- 13) Girard TD, Kress JP, Fuchs BD, et al. Efficacy and safety of a paired sedation and ventilator weaning protocol for mechanically ventilated patients in intensive care (Awakening and Breathing Controlled trial): a randomised controlled trial. *Lancet* 2008;371:126-34.
- 14) 日本集中治療医学会, 日本呼吸療法医学会, 日本クリティカルケア看護学会. 人工呼吸器離脱に関する3学会合同プロトコル. 2015.

- 15) Yang KL, Tobin MJ. A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation. *N Engl J Med* 1991;324:1445-50.
- 16) Subirà C, Hernández G, Vázquez A, et al. Effect of pressure support vs T-piece ventilation strategies during spontaneous breathing trials on successful extubation among patients receiving mechanical ventilation: a randomized clinical trial. *JAMA* 2019;321:2175-82.
- 17) Thille AW, Harrois A, Schortgen F, et al. Outcomes of extubation failure in medical intensive care unit patients. *Crit Care Med* 2011;39:2612-8.
- 18) François B, Bellissant E, Gissot V, et al. 12-h pretreatment with methylprednisolone versus placebo for prevention of postextubation laryngeal oedema: a randomised double-blind trial. *Lancet* 2007;369:1083-9.
- 19) Hernández G, Vaquero C, González P, et al. Effect of postextubation high-flow nasal cannula vs conventional oxygen therapy on reintubation in low-risk patients: a randomized clinical trial. *JAMA* 2016;315:1354-61.
- 20) Thille AW, Muller G, Gacouin A, et al. Effect of postextubation high-flow nasal oxygen with noninvasive ventilation vs high-flow nasal oxygen alone on reintubation among patients at high risk of extubation failure: a randomized clinical trial. *JAMA* 2019;322:1465-75.
- 21) Oczkowski S, Ergon B, Bos L, et al. ERS clinical practice guidelines: high-flow nasal cannula in acute respiratory failure. *Eur Respir J* 2022;59:2101574.
- 22) Young D, Harrison DA, Cuthbertson BH, et al. Effect of early vs late tracheostomy placement on survival in patients receiving mechanical ventilation: the TracMan randomized trial. *JAMA* 2013;309:2121-9.
- 23) 各薬剤の電子添文（ケタラル静注用、ディプリバン注、ドルミカム注射液、エスラックス静注、スキサメトニウム注、プリディオ静注。KEGG MEDICUSで2026年10月に確認）。
- 24) Loughnan D, Walker HGM, Woodland N, et al. Ketamine versus alternate agents to prevent postinduction haemodynamic instability during tracheal intubation of critically ill adults: a systematic review and meta-analysis. *Br J Anaesth* 2026. doi:10.1016/j.bja.2026.02.023.
- 25) Guihard B, Chollet-Xémard C, Lakhnati P, et al. Effect of rocuronium vs succinylcholine on endotracheal intubation success rate among patients undergoing out-of-hospital rapid sequence intubation: a randomized clinical trial. *JAMA* 2019;322:2303-12.
- 26) Prekker ME, Driver BE, Trent SA, et al. Video versus direct laryngoscopy for tracheal intubation of critically ill adults. *N Engl J Med* 2023;389:418-29.
- 27) Russotto V, Myatra SN, Laffey JG, et al. Intubation practices and adverse peri-intubation events in critically ill patients from 29 countries. *JAMA* 2021;325:1164-72.
- 28) Russell DW, Casey JD, Gibbs KW, et al. Effect of fluid bolus administration on cardiovascular collapse among critically ill patients undergoing tracheal intubation: a randomized clinical trial. *JAMA* 2022;328:270-9.
- 29) Gattinoni L, Pesenti A. The concept of "baby lung". *Intensive Care Med* 2005;31:776-84.
- 30) Serpa Neto A, Cardoso SO, Manetta JA, et al. Association between use of lung-protective ventilation with lower tidal volumes and clinical outcomes among patients without acute respiratory distress syndrome: a meta-analysis. *JAMA* 2012;308:1651-9.
- 31) Writing Group for the PREVENT Investigators. Effect of a low vs intermediate tidal volume strategy on ventilator-free days in intensive care unit patients without ARDS: a randomized clinical trial. *JAMA* 2018;320:1872-80.

1-6 頭部・胸部・腹部CTの見方

- 1) Alfaro D, Levitt MA, English DK, et al. Accuracy of interpretation of cranial computed tomography scans in an emergency medicine residency program. *Ann Emerg Med* 1995;25:169-74.

- 2) Shaish H, Ream J, Huang C, et al. Diagnostic accuracy of unenhanced computed tomography for evaluation of acute abdominal pain in the emergency department. *JAMA Surg* 2023;158:e231112.
- 3) イオパミロン注 電子添文（バイエル薬品）. 2. 禁忌、8. 重要な基本的注意、9.1 合併症・既往歴等のある患者、14. 適用上の注意.
- 4) Katayama H, Yamaguchi K, Kozuka T, et al. Adverse reactions to ionic and nonionic contrast media. A report from the Japanese Committee on the Safety of Contrast Media. *Radiology* 1990;175:621-8.
- 5) Bullock MR, Chesnut R, Ghajar J, et al. Surgical management of acute subdural hematomas. *Neurosurgery* 2006;58(3 Suppl):S16-24.
- 6) Chalela JA, Kidwell CS, Nentwich LM, et al. Magnetic resonance imaging and computed tomography in emergency assessment of patients with suspected acute stroke: a prospective comparison. *Lancet* 2007;369:293-8.
- 7) Mair G, Boyd EV, Chappell FM, et al. Sensitivity and specificity of the hyperdense artery sign for arterial obstruction in acute ischemic stroke. *Stroke* 2015;46:102-7.
- 8) Lev MH, Farkas J, Gemmete JJ, et al. Acute stroke: improved nonenhanced CT detection—benefits of soft-copy interpretation by using variable window width and center level settings. *Radiology* 1999;213:150-5.
- 9) Perry JJ, Stiell IG, Sivilotti ML, et al. Sensitivity of computed tomography performed within six hours of onset of headache for diagnosis of subarachnoid haemorrhage: prospective cohort study. *BMJ* 2011;343:d4277.
- 10) Dubosh NM, Bellolio MF, Rabinstein AA, Edlow JA. Sensitivity of early brain computed tomography to exclude aneurysmal subarachnoid hemorrhage: a systematic review and meta-analysis. *Stroke* 2016;47:750-5.
- 11) Hwang DY, Silva GS, Furie KL, Greer DM. Comparative sensitivity of computed tomography vs. magnetic resonance imaging for detecting acute posterior fossa infarct. *J Emerg Med* 2012;42:559-65.
- 12) Edlow BL, Hurwitz S, Edlow JA. Diagnosis of DWI-negative acute ischemic stroke: a meta-analysis. *Neurology* 2017;89:256-62.
- 13) Bankier AA, MacMahon H, Colby T, et al. Fleischner Society: glossary of terms for thoracic imaging. *Radiology* 2024;310:e232558.
- 14) Gadkowski LB, Stout JE. Cavitory pulmonary disease. *Clin Microbiol Rev* 2008;21:305-33.
- 15) Stein PD, Fowler SE, Goodman LR, et al. Multidetector computed tomography for acute pulmonary embolism. *N Engl J Med* 2006;354:2317-27.
- 16) Millet I, Taourel P, Ruyer A, Molinari N. Value of CT findings to predict surgical ischemia in small bowel obstruction: a systematic review and meta-analysis. *Eur Radiol* 2015;25:1823-35.
- 17) Menke J. Diagnostic accuracy of multidetector CT in acute mesenteric ischemia: systematic review and meta-analysis. *Radiology* 2010;256:93-101.
- 18) Rud B, Vejborg TS, Rappeport ED, et al. Computed tomography for diagnosis of acute appendicitis in adults. *Cochrane Database Syst Rev* 2019;2019(11):CD009977.
- 19) Laméris W, van Randen A, Bipat S, et al. Graded compression ultrasonography and computed tomography in acute colonic diverticulitis: meta-analysis of test accuracy. *Eur Radiol* 2008;18:2498-511.
- 20) Kiewiet JJ, Leeuwenburgh MM, Bipat S, et al. A systematic review and meta-analysis of diagnostic performance of imaging in acute cholecystitis. *Radiology* 2012;264:708-20.
- 21) Wertz JR, Lopez JM, Olson D, Thompson WM. Comparing the diagnostic accuracy of ultrasound and CT in evaluating acute cholecystitis. *AJR Am J Roentgenol* 2018;211:W92-7.

- 22) Ahn JH, Chung JJ, Yu JS, Kim JH, Cho ES, Kim DJ. Prognostic value of gallbladder wall thickening in patients with acute hepatitis A. *Ultrasonography* 2015;34:139-43.
- 23) 日本医療機能評価機構 医療事故情報収集等事業. 医療安全情報 No.138 画像診断報告書の確認不足 (第2報). 2018年5月.
- 24) Lasser EC, Berry CC, Talner LB, et al. Pretreatment with corticosteroids to alleviate reactions to intravenous contrast material. *N Engl J Med* 1987;317:845-9.
- 25) Lasser EC, Berry CC, Mishkin MM, et al. Pretreatment with corticosteroids to prevent adverse reactions to nonionic contrast media. *AJR Am J Roentgenol* 1994;162:523-6.
- 26) McDonald JS, Larson NB, Kolbe AB, et al. Prevention of allergic-like reactions at repeat CT: steroid pretreatment versus contrast material substitution. *Radiology* 2021;301:133-40.
- 27) Abe S, Fukuda H, Tobe K, Ibukuro K. Protective effect against repeat adverse reactions to iodinated contrast medium: premedication vs. changing the contrast medium. *Eur Radiol* 2016;26:2148-54.
- 28) 日本医学放射線学会 造影剤安全性委員会. ヨード造影剤ならびにガドリニウム造影剤の急性 (即時性) 副作用発症の危険性低減を目的としたステロイド前投薬に関する提言 (2022年12月改訂第3版). 2022.

1-7 穿刺の手法

- 1) Gordon CE, Feller-Kopman D, Balk EM, Smetana GW. Pneumothorax following thoracentesis: a systematic review and meta-analysis. *Arch Intern Med* 2010;170:332-9.
- 2) Mercaldi CJ, Lanes SF. Ultrasound guidance decreases complications and improves the cost of care among patients undergoing thoracentesis and paracentesis. *Chest* 2013;143:532-8.
- 3) Asciak R, Bedawi EO, Bhatnagar R, et al. British Thoracic Society Clinical Statement on pleural procedures. *Thorax* 2023;78(Suppl 3):s43-s68.
- 4) Coz Yataco A, Soghier I, Hébert PC, et al. Transfusion of fresh frozen plasma and platelets in critically ill adults: an American College of Chest Physicians clinical practice guideline. *Chest* 2025;168:661-76.
- 5) Ault MJ, Rosen BT, Scher J, Feinglass J, Barsuk JH. Thoracentesis outcomes: a 12-year experience. *Thorax* 2015;70:127-32.
- 6) Tan JL, Lokan T, Chinnaratha MA, Veysey M. Risk of bleeding after abdominal paracentesis in patients with chronic liver disease and coagulopathy: a systematic review and meta-analysis. *JGH Open* 2024;8:e70013.
- 7) Ahmed I, Gertner E. Safety of arthrocentesis and joint injection in patients receiving anticoagulation at therapeutic levels. *Am J Med* 2012;125:265-9.
- 8) キシロカイン注ポリアンブ1%の電子添文 (KEGG MEDICUSで2026年10月に確認). 6. 用法及び用量.
- 9) Engelborghs S, Niemantsverdriet E, Struyfs H, et al. Consensus guidelines for lumbar puncture in patients with neurological diseases. *Alzheimers Dement (Amst)* 2017;8:111-26.
- 10) 日本神経学会, 日本神経治療学会, 日本神経感染症学会監修. 細菌性髄膜炎診療ガイドライン2014. 南江堂; 2014.
- 11) Hasbun R, Abrahams J, Jekel J, Quagliarello VJ. Computed tomography of the head before lumbar puncture in adults with suspected meningitis. *N Engl J Med* 2001;345:1727-33.
- 12) Bodilsen J, Mariager T, Vestergaard HH, et al. Association of lumbar puncture with spinal hematoma in patients with and without coagulopathy. *JAMA* 2020;324:1419-28.
- 13) Nath S, Koziarz A, Badhiwala JH, et al. Atraumatic versus conventional lumbar puncture needles: a systematic review and meta-analysis. *Lancet* 2018;391:1197-204.

- 14) Whiteley W, Al-Shahi R, Warlow CP, Zeidler M, Lueck CJ. CSF opening pressure: reference interval and the effect of body mass index. *Neurology* 2006;67:1690-1.
- 15) Glantz MJ, Cole BF, Glantz LK, et al. Cerebrospinal fluid cytology in patients with cancer: minimizing false-negative results. *Cancer* 1998;82:733-9.
- 16) Wasserstrom WR, Glass JP, Posner JB. Diagnosis and treatment of leptomeningeal metastases from solid tumors: experience with 90 patients. *Cancer* 1982;49:759-72.
- 17) Strupp M, Brandt T, Müller A. Incidence of post-lumbar puncture syndrome reduced by reinserting the stylet: a randomized prospective study of 600 patients. *J Neurol* 1998;245:589-92.
- 18) Arevalo-Rodriguez I, Ciapponi A, Roqué i Figuls M, et al. Posture and fluids for preventing post-dural puncture headache. *Cochrane Database Syst Rev* 2016;3:CD009199.
- 19) Rowley MW, Agarwal S, Seetharam AB, Hirsch KS. Real-time ultrasound-guided paracentesis by radiologists: near zero risk of hemorrhage without correction of coagulopathy. *J Vasc Interv Radiol* 2019;30:259-64.
- 20) Sakai H, Sheer TA, Mendler MH, Runyon BA. Choosing the location for non-image guided abdominal paracentesis. *Liver Int* 2005;25:984-6.
- 21) 日本消化器病学会, 日本肝臓学会. 肝硬変診療ガイドライン2020 (改訂第3版). 2020.
- 22) Biggins SW, Angeli P, Garcia-Tsao G, et al. Diagnosis, evaluation, and management of ascites, spontaneous bacterial peritonitis and hepatorenal syndrome: 2021 practice guidance by the American Association for the Study of Liver Diseases. *Hepatology* 2021;74:1014-48.
- 23) De Gottardi A, Thévenot T, Spahr L, et al. Risk of complications after abdominal paracentesis in cirrhotic patients: a prospective study. *Clin Gastroenterol Hepatol* 2009;7:906-9.
- 24) Ravn C, Neyt J, Benito N, et al. Guideline for management of septic arthritis in native joints (SANJO). *J Bone Jt Infect* 2023;8:29-37.
- 25) Shiloh AL, Savel RH, Paulin LM, Eisen LA. Ultrasound-guided catheterization of the radial artery: a systematic review and meta-analysis of randomized controlled trials. *Chest* 2011;139:524-9.
- 26) Scheer B, Perel A, Pfeiffer UJ. Clinical review: complications and risk factors of peripheral arterial catheters used for haemodynamic monitoring in anaesthesia and intensive care medicine. *Crit Care* 2002;6:199-204.
- 27) Stolz LA, Stolz U, Howe C, Farrell IJ, Adhikari S. Ultrasound-guided peripheral venous access: a meta-analysis and systematic review. *J Vasc Access* 2015;16:321-6.
- 28) Gottlieb M, Holladay D, Peksa GD. Comparison of short- vs long-axis technique for ultrasound-guided peripheral line placement: a systematic review and meta-analysis. *Cureus* 2018;10:e2718.
- 29) Pandurangadu AV, Tucker J, Brackney AR, Bahl A. Ultrasound-guided intravenous catheter survival impacted by amount of catheter residing in the vein. *Emerg Med J* 2018;35:550-5.
- 30) Bahl A, Hang B, Brackney A, et al. Standard long IV catheters versus extended dwell catheters: a randomized comparison of ultrasound-guided catheter survival. *Am J Emerg Med* 2019;37:715-21.
- 31) 日本ペインクリニック学会, 日本麻酔科学会, 日本区域麻酔学会. 抗血栓療法中の区域麻酔・神経ブロックガイドライン. 2016.
- 32) Alemán C, Alegre J, Armadans L, et al. The value of chest roentgenography in the diagnosis of pneumothorax after thoracentesis. *Am J Med* 1999;107:340-3.
- 33) Petersen WG, Zimmerman R. Limited utility of chest radiograph after thoracentesis. *Chest* 2000;117:1038-42.
- 34) Jarvis MA, Jarvis CL, Jones PR, Spyt TJ. Reliability of Allen's test in selection of patients for radial artery harvest. *Ann Thorac Surg* 2000;70:1362-5.

- 35) Valgimigli M, Campo G, Penzo C, et al. Transradial coronary catheterization and intervention across the whole spectrum of Allen test results. *J Am Coll Cardiol* 2014;63:1833-41.

第2部 意識と神経の症候

2-1 意識障害

- 1) Leong LB, Jian KH, Vasu A, Seow E. Prospective study of patients with altered mental status: clinical features and outcome. *Int J Emerg Med* 2008;1:179-82.
- 2) 日本糖尿病学会編. 糖尿病診療ガイドライン2024. 南江堂；2024.
- 3) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）.
- 4) Ikeda M, Matsunaga T, Irabu N, Yoshida S. Using vital signs to diagnose impaired consciousness: cross sectional observational study. *BMJ* 2002;325:800.
- 5) Tokuda Y, Nakazato N, Stein GH. Pupillary evaluation for differential diagnosis of coma. *Postgrad Med J* 2003;79:49-51.
- 6) Graus F, Titulaer MJ, Balu R, et al. A clinical approach to diagnosis of autoimmune encephalitis. *Lancet Neurol* 2016;15:391-404.
- 7) Arena JE, Rabinstein AA. Transient global amnesia. *Mayo Clin Proc* 2015;90:264-72.
- 8) Hodges JR, Warlow CP. Syndromes of transient amnesia: towards a classification. A study of 153 cases. *J Neurol Neurosurg Psychiatry* 1990;53:834-43.
- 9) Arena JE, Brown RD, Mandrekar J, Rabinstein AA. Long-term outcome in patients with transient global amnesia: a population-based study. *Mayo Clin Proc* 2017;92:399-405.
- 10) 日本緩和医療学会ガイドライン統括委員会編. がん疼痛の薬物療法に関するガイドライン2020年版. 金原出版；2020.
- 11) Penninga EI, Graudal N, Ladekarl MB, Jürgens G. Adverse events associated with flumazenil treatment for the management of suspected benzodiazepine intoxication: a systematic review with meta-analyses of randomised trials. *Basic Clin Pharmacol Toxicol* 2016;118:37-44.
- 12) Payne LE, Gagnon DJ, Riker RR, et al. Cefepime-induced neurotoxicity: a systematic review. *Crit Care* 2017;21:276.
- 13) Hasbun R, Abrahams J, Jekel J, Quagliarello VJ. Computed tomography of the head before lumbar puncture in adults with suspected meningitis. *N Engl J Med* 2001;345:1727-33.
- 14) 日本神経学会, 日本神経治療学会, 日本神経感染症学会監修. 細菌性髄膜炎診療ガイドライン2014. 南江堂；2014.
- 15) Freund Y, Viglino D, Cachanado M, et al. Effect of noninvasive airway management of comatose patients with acute poisoning: a randomized clinical trial. *JAMA* 2023;330:2267-74.
- 16) Mercer-Smith GW, Appleton M, Hanon ÉA, Bowron A. Blood samples for ammonia analysis do not require transport to the laboratory on ice: a study of ammonia stability and cause of in vitro ammonia increase in samples from patients with hyperammonaemia. *Clin Chem Lab Med* 2025;63:1132-8.
- 17) Gaul C, Dietrich W, Friedrich I, et al. Neurological symptoms in type A aortic dissections. *Stroke* 2007;38:292-7.

2-2 失神

- 1) 日本循環器学会, 日本不整脈心電学会. 2022年改訂版 不整脈の診断とリスク評価に関するガイドライン. 2022.
- 2) Brignole M, Moya A, de Lange FJ, et al. 2018 ESC Guidelines for the diagnosis and management of syncope. *Eur Heart J* 2018;39:1883-948.

- 3) Soteriades ES, Evans JC, Larson MG, et al. Incidence and prognosis of syncope. *N Engl J Med* 2002;347:878-85.
- 4) Shen WK, Sheldon RS, Benditt DG, et al. 2017 ACC/AHA/HRS guideline for the evaluation and management of patients with syncope. *Circulation* 2017;136:e60-e122.
- 5) Benbadis SR, Wolgamuth BR, Goren H, et al. Value of tongue biting in the diagnosis of seizures. *Arch Intern Med* 1995;155:2346-9.
- 6) Sheldon R, Rose S, Ritchie D, et al. Historical criteria that distinguish syncope from seizures. *J Am Coll Cardiol* 2002;40:142-8.
- 7) Albassam OT, Redelmeier RJ, Shadowitz S, et al. Did this patient have cardiac syncope? The Rational Clinical Examination systematic review. *JAMA* 2019;321:2448-57.
- 8) Freeman R, Wieling W, Axelrod FB, et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clin Auton Res* 2011;21:69-72.
- 9) Prandoni P, Lensing AW, Prins MH, et al. Prevalence of pulmonary embolism among patients hospitalized for syncope. *N Engl J Med* 2016;375:1524-31.
- 10) Costantino G, Ruwald MH, Quinn J, et al. Prevalence of pulmonary embolism in patients with syncope. *JAMA Intern Med* 2018;178:356-62.
- 11) Nallamoorthy BK, Mehta RH, Saint S, et al. Syncope in acute aortic dissection: diagnostic, prognostic, and clinical implications. *Am J Med* 2002;113:468-71.
- 12) 日本循環器学会, 日本心血管インターベンション治療学会, 日本心臓病学会. 2023年 JCS/CVIT/JCC ガイドライン フォーカスアップデート版 冠攣縮性狭心症と冠微小循環障害の診断と治療. 2023.
- 13) Igarashi Y, Tamura Y, Tanabe Y, et al. Angina-linked syncope and lack of calcium antagonist therapy predict cardiac arrest before definitive diagnosis of vasospastic angina. *Coron Artery Dis* 1994;5:881-7.
- 14) Thiruganasambandamoorthy V, Rowe BH, Sivilotti MLA, et al. Duration of electrocardiographic monitoring of emergency department patients with syncope. *Circulation* 2019;139:1396-406.
- 15) Thiruganasambandamoorthy V, Kwong K, Wells GA, et al. Development of the Canadian Syncope Risk Score to predict serious adverse events after emergency department assessment of syncope. *CMAJ* 2016;188:E289-98.
- 16) Thiruganasambandamoorthy V, Sivilotti MLA, Le Sage N, et al. Multicenter emergency department validation of the Canadian Syncope Risk Score. *JAMA Intern Med* 2020;180:737-44.
- 17) Zimmermann T, du Fay de Lavallaz J, Nestelberger T, et al. International validation of the Canadian Syncope Risk Score: a cohort study. *Ann Intern Med* 2022;175:783-94.
- 18) Saccilotto RT, Nickel CH, Bucher HC, et al. San Francisco Syncope Rule to predict short-term serious outcomes: a systematic review. *CMAJ* 2011;183:E1116-26.
- 19) van Dijk N, Quartieri F, Blanc JJ, et al. Effectiveness of physical counterpressure maneuvers in preventing vasovagal syncope: the Physical Counterpressure Manoeuvres Trial (PC-Trial). *J Am Coll Cardiol* 2006;48:1652-7.
- 20) Aziz HM, Zarzecki MP, Garcia-Zamora S, et al. Pathogenesis and management of Brugada syndrome: recent advances and protocol for umbrella reviews of meta-analyses in major arrhythmic events risk stratification. *J Clin Med* 2022;11:1912.
- 21) Michowitz Y, Milman A, Sarquella-Brugada G, et al. Fever-related arrhythmic events in the multicenter Survey on Arrhythmic Events in Brugada Syndrome. *Heart Rhythm* 2018;15:1394-401.
- 22) 道路交通法施行令 第33条の2の3.
- 23) 日本不整脈学会 植込み型デバイス関連社会問題対策委員会 失神患者の自動車運転に関するワーキンググループ. 再発性の失神患者における自動車運転制限のガイドラインとその運用指針. 2013.

- 24) Numé AK, Gislason G, Christiansen CB, et al. Syncope and motor vehicle crash risk: a Danish nationwide study. *JAMA Intern Med* 2016;176:503-10.
- 25) Viau JA, Chaudry H, Hannigan A, et al. The yield of computed tomography of the head among patients presenting with syncope: a systematic review. *Acad Emerg Med* 2019;26:479-90.
- 26) Kühne M, Schaer B, Moulay N, Sticherling C, Osswald S. Holter monitoring for syncope: diagnostic yield in different patient groups and impact on device implantation. *QJM* 2007;100:771-7.
- 27) Krahn AD, Klein GJ, Yee R, Skanes AC. Randomized assessment of syncope trial: conventional diagnostic testing versus a prolonged monitoring strategy. *Circulation* 2001;104:46-51.

2-3 けいれん・てんかん重積

- 1) 日本神経学会監修. てんかん診療ガイドライン2018 (追補版2022を含む). 医学書院; 2018.
- 2) Trinka E, Cock H, Hesdorffer D, et al. A definition and classification of status epilepticus--Report of the ILAE Task Force on Classification of Status Epilepticus. *Epilepsia* 2015;56:1515-23.
- 3) Kapur J, Elm J, Chamberlain JM, et al. Randomized trial of three anticonvulsant medications for status epilepticus. *N Engl J Med* 2019;381:2103-13.
- 4) 各薬剤の電子添文 (KEGG MEDICUSで2026年9月に確認).
- 5) Silbergleit R, Durkalski V, Lowenstein D, et al. Intramuscular versus intravenous therapy for prehospital status epilepticus. *N Engl J Med* 2012;366:591-600.
- 6) Alldredge BK, Gelb AM, Isaacs SM, et al. A comparison of lorazepam, diazepam, and placebo for the treatment of out-of-hospital status epilepticus. *N Engl J Med* 2001;345:631-7.
- 7) Sathe AG, Tillman H, Coles LD, et al. Underdosing of benzodiazepines in patients with status epilepticus enrolled in Established Status Epilepticus Treatment Trial. *Acad Emerg Med* 2019;26:940-3.
- 8) Chamberlain JM, Kapur J, Shinnar S, et al. Efficacy of levetiracetam, fosphenytoin, and valproate for established status epilepticus by age group (ESETT): a double-blind, responsive-adaptive, randomised controlled trial. *Lancet* 2020;395:1217-24.
- 9) DeLorenzo RJ, Waterhouse EJ, Towne AR, et al. Persistent nonconvulsive status epilepticus after the control of convulsive status epilepticus. *Epilepsia* 1998;39:833-40.
- 10) Towne AR, Waterhouse EJ, Boggs JG, et al. Prevalence of nonconvulsive status epilepticus in comatose patients. *Neurology* 2000;54:340-5.
- 11) Avbersek A, Sisodiya S. Does the primary literature provide support for clinical signs used to distinguish psychogenic nonepileptic seizures from epileptic seizures? *J Neurol Neurosurg Psychiatry* 2010;81:719-25.
- 12) Matz O, Zdebik C, Zechbauer S, et al. Lactate as a diagnostic marker in transient loss of consciousness. *Seizure* 2016;40:71-5.
- 13) Hung TY, Chen CC, Wang TL, Su CF, Wang RF. Transient hyperammonemia in seizures: a prospective study. *Epilepsia* 2011;52:2043-9.
- 14) Harden CL, Huff JS, Schwartz TH, et al. Reassessment: neuroimaging in the emergency patient presenting with seizure (an evidence-based review). *Neurology* 2007;69:1772-80.
- 15) Orringer CE, Eustace JC, Wunsch CD, Gardner LB. Natural history of lactic acidosis after grand-mal seizures. *N Engl J Med* 1977;297:796-9.
- 16) Beghi E, Carpio A, Forsgren L, et al. Recommendation for a definition of acute symptomatic seizure. *Epilepsia* 2010;51:671-5.
- 17) Tumani H, Jobs C, Bretschneider J, Hoppner AC, Kerling F, Fauser S. Effect of epileptic seizures on the cerebrospinal fluid--a systematic retrospective analysis. *Epilepsy Res* 2015;114:23-31.

- 18) Chen DK, So YT, Fisher RS. Use of serum prolactin in diagnosing epileptic seizures: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 2005;65:668-75.
- 19) Huff JS, Melnick ER, Tomaszewski CA, et al. Clinical policy: critical issues in the evaluation and management of adult patients presenting to the emergency department with seizures. *Ann Emerg Med* 2014;63:437-47.
- 20) Marson A, Jacoby A, Johnson A, et al. Immediate versus deferred antiepileptic drug treatment for early epilepsy and single seizures: a randomised controlled trial. *Lancet* 2005;365:2007-13.
- 21) 警察庁交通局運転免許課長. 一定の病氣等に係る運転免許関係事務に関する運用上の留意事項について (通達) (丁運発第73号) 別添「一定の病氣に係る免許の可否等の運用基準」. 2025.
- 22) 日本てんかん学会. てんかんと運転に関する提言. 2012.
- 23) 日本てんかん学会. てんかんに関する医師の届け出ガイドライン. 2014.
- 24) Krumholz A, Wiebe S, Gronseth GS, et al. Evidence-based guideline: management of an unprovoked first seizure in adults: report of the Guideline Development Subcommittee of the American Academy of Neurology and the American Epilepsy Society. *Neurology* 2015;84:1705-13.
- 25) Prasad M, Krishnan PR, Sequeira R, Al-Roomi K. Anticonvulsant therapy for status epilepticus. *Cochrane Database Syst Rev* 2014;(9):CD003723.
- 26) Chamberlain JM, Okada P, Holsti M, et al. Lorazepam vs diazepam for pediatric status epilepticus: a randomized clinical trial. *JAMA* 2014;311:1652-60.
- 27) Greenblatt DJ, Divoll M. Diazepam versus lorazepam: relationship of drug distribution to duration of clinical action. *Adv Neurol* 1983;34:487-91.

2-4 頭痛

- 1) 日本神経学会, 日本頭痛学会, 日本神経治療学会監修. 頭痛の診療ガイドライン2021. 医学書院; 2021.
- 2) Kowalski RG, Claassen J, Kreiter KT, et al. Initial misdiagnosis and outcome after subarachnoid hemorrhage. *JAMA* 2004;291:866-9.
- 3) Goldstein JN, Camargo CA Jr, Pelletier AJ, Edlow JA. Headache in United States emergency departments: demographics, work-up and frequency of pathological diagnoses. *Cephalalgia* 2006;26:684-90.
- 4) Polmear A. Sentinel headaches in aneurysmal subarachnoid haemorrhage: what is the true incidence? A systematic review. *Cephalalgia* 2003;23:935-41.
- 5) Landtblom AM, Fridriksson S, Boivie J, et al. Sudden onset headache: a prospective study of features, incidence and causes. *Cephalalgia* 2002;22:354-60.
- 6) Perry JJ, Stiell IG, Sivilotti ML, et al. Sensitivity of computed tomography performed within six hours of onset of headache for diagnosis of subarachnoid haemorrhage: prospective cohort study. *BMJ* 2011;343:d4277.
- 7) Dubosh NM, Bellolio MF, Rabinstein AA, Edlow JA. Sensitivity of early brain computed tomography to exclude aneurysmal subarachnoid hemorrhage: a systematic review and meta-analysis. *Stroke* 2016;47:750-5.
- 8) Perry JJ, Stiell IG, Sivilotti ML, et al. Clinical decision rules to rule out subarachnoid hemorrhage for acute headache. *JAMA* 2013;310:1248-55.
- 9) Perry JJ, Sivilotti ML, Sutherland J, et al. Validation of the Ottawa Subarachnoid Hemorrhage Rule in patients with acute headache. *CMAJ* 2017;189:E1379-85.
- 10) 日本脳卒中学会 脳卒中ガイドライン委員会編. 脳卒中治療ガイドライン2021〔改訂2025〕. 協和企画; 2025.

- 11) Cruickshank A, Auld P, Beetham R, et al. Revised national guidelines for analysis of cerebrospinal fluid for bilirubin in suspected subarachnoid haemorrhage. *Ann Clin Biochem* 2008;45:238-44.
- 12) Perry JJ, Alyahya B, Sivilotti ML, et al. Differentiation between traumatic tap and aneurysmal subarachnoid hemorrhage: prospective cohort study. *BMJ* 2015;350:h568.
- 13) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）。
- 14) Post R, Germans MR, Tjerkstra MA, et al. Ultra-early tranexamic acid after subarachnoid haemorrhage (ULTRA): a randomised controlled trial. *Lancet* 2021;397:112-8.
- 15) 日本神経学会, 日本神経治療学会, 日本神経感染症学会監修. 細菌性髄膜炎診療ガイドライン2014. 南江堂; 2014.
- 16) 日本循環器学会ほか. 2026年改訂版 大型血管炎診療ガイドライン. 2026.
- 17) van der Geest KSM, Sandovici M, Brouwer E, Mackie SL. Diagnostic accuracy of symptoms, physical signs, and laboratory tests for giant cell arteritis: a systematic review and meta-analysis. *JAMA Intern Med* 2020;180:1295-304.
- 18) Silbert PL, Mokri B, Schievink WI. Headache and neck pain in spontaneous internal carotid and vertebral artery dissections. *Neurology* 1995;45:1517-22.
- 19) Brouwer MC, Coutinho JM, van de Beek D. Clinical characteristics and outcome of brain abscess: systematic review and meta-analysis. *Neurology* 2014;82:806-13.
- 20) Cohen AS, Burns B, Goadsby PJ. High-flow oxygen for treatment of cluster headache: a randomized trial. *JAMA* 2009;302:2451-7.
- 21) Headache Classification Committee of the International Headache Society (IHS). The International Classification of Headache Disorders, 3rd edition. *Cephalalgia* 2018;38:1-211.
- 22) Parlak I, Erdur B, Parlak M, et al. Intravenous administration of metoclopramide by 2 min bolus vs 15 min infusion: does it affect the improvement of headache while reducing the side effects? *Postgrad Med J* 2007;83:664-8.
- 23) Egerton-Warburton D, Povey K. Administration of metoclopramide by infusion or bolus does not affect the incidence of drug-induced akathisia. *Emerg Med Australas* 2013;25:207-12.
- 24) Friedman BW, Irizarry E, Solorzano C, et al. Randomized study of IV prochlorperazine plus diphenhydramine vs IV hydromorphone for migraine. *Neurology* 2017;89:2075-82.
- 25) Colman I, Friedman BW, Brown MD, et al. Parenteral dexamethasone for acute severe migraine headache: meta-analysis of randomised controlled trials for preventing recurrence. *BMJ* 2008;336:1359-61.
- 26) Dentali F, Squizzato A, Marchesi C, Bonzini M, Ferro JM, Ageno W. D-dimer testing in the diagnosis of cerebral vein thrombosis: a systematic review and a meta-analysis of the literature. *J Thromb Haemost* 2012;10:582-9.
- 27) Heldner MR, Zuurbier SM, Li B, et al. Prediction of cerebral venous thrombosis with a new clinical score and D-dimer levels. *Neurology* 2020;95:e898-909.

2-5 めまい

- 1) Kerber KA, Brown DL, Lisabeth LD, et al. Stroke among patients with dizziness, vertigo, and imbalance in the emergency department: a population-based study. *Stroke* 2006;37:2484-7.
- 2) Newman-Toker DE, Cannon LM, Stofferahn ME, et al. Imprecision in patient reports of dizziness symptom quality: a cross-sectional study conducted in an acute care setting. *Mayo Clin Proc* 2007;82:1329-40.
- 3) Newman-Toker DE, Hsieh YH, Camargo CA Jr, et al. Spectrum of dizziness visits to US emergency departments: cross-sectional analysis from a nationally representative sample. *Mayo Clin Proc* 2008;83:765-75.

- 4) Edlow JA, Carpenter C, Akhter M, et al. Guidelines for reasonable and appropriate care in the emergency department 3 (GRACE-3): acute dizziness and vertigo in the emergency department. *Acad Emerg Med* 2023;30:442-86.
- 5) Lindell E, Finizia C, Johansson M, et al. Asking about dizziness when turning in bed predicts examination findings for benign paroxysmal positional vertigo. *J Vestib Res* 2018;28:339-47.
- 6) Paul NL, Simoni M, Rothwell PM; Oxford Vascular Study. Transient isolated brainstem symptoms preceding posterior circulation stroke: a population-based study. *Lancet Neurol* 2013;12:65-71.
- 7) Kattah JC, Talkad AV, Wang DZ, et al. HINTS to diagnose stroke in the acute vestibular syndrome: three-step bedside oculomotor examination more sensitive than early MRI diffusion-weighted imaging. *Stroke* 2009;40:3504-10.
- 8) Newman-Toker DE, Kerber KA, Hsieh YH, et al. HINTS outperforms ABCD2 to screen for stroke in acute continuous vertigo and dizziness. *Acad Emerg Med* 2013;20:986-96.
- 9) Shah VP, Oliveira J E Silva L, Farah W, et al. Diagnostic accuracy of the physical examination in emergency department patients with acute vertigo or dizziness: a systematic review and meta-analysis for GRACE-3. *Acad Emerg Med* 2023;30:552-78.
- 10) Ohle R, Montpellier RA, Marchadier V, et al. Can emergency physicians accurately rule out a central cause of vertigo using the HINTS examination? A systematic review and meta-analysis. *Acad Emerg Med* 2020;27:887-96.
- 11) Gottlieb M, Peksa GD, Carlson JN. Head impulse, nystagmus, and test of skew examination for diagnosing central causes of acute vestibular syndrome. *Cochrane Database Syst Rev* 2023;11:CD015089.
- 12) Ronchetti M, Bartalucci P, Pepe G, et al. Diagnostic accuracy of the STANDING algorithm in patients with isolated vertigo: a multicentre prospective study (STANDING-M). *Emerg Med J* 2025;42:791-7.
- 13) Lee H, Sohn SI, Cho YW, et al. Cerebellar infarction presenting isolated vertigo: frequency and vascular topographical patterns. *Neurology* 2006;67:1178-83.
- 14) Chalela JA, Kidwell CS, Nentwich LM, et al. Magnetic resonance imaging and computed tomography in emergency assessment of patients with suspected acute stroke: a prospective comparison. *Lancet* 2007;369:293-8.
- 15) Saber Tehrani AS, Kattah JC, Mantokoudis G, et al. Small strokes causing severe vertigo: frequency of false-negative MRIs and nonlacunar mechanisms. *Neurology* 2014;83:169-73.
- 16) Entwistle T, Perchyonok Y, Fitt G. Thin section magnetic resonance diffusion-weighted imaging in the detection of acute infratentorial stroke. *J Med Imaging Radiat Oncol* 2016;60:616-23.
- 17) Edlow JA, Newman-Toker DE, Savitz SI. Diagnosis and initial management of cerebellar infarction. *Lancet Neurol* 2008;7:951-64.
- 18) Bhattacharyya N, Gubbels SP, Schwartz SR, et al. Clinical practice guideline: benign paroxysmal positional vertigo (update). *Otolaryngol Head Neck Surg* 2017;156(3 Suppl):S1-S47.
- 19) von Brevern M, Bertholon P, Brandt T, et al. Benign paroxysmal positional vertigo: diagnostic criteria. *J Vestib Res* 2015;25:105-17.
- 20) Hilton MP, Pinder DK. The Epley (canalith repositioning) manoeuvre for benign paroxysmal positional vertigo. *Cochrane Database Syst Rev* 2014;(12):CD003162.
- 21) 日本めまい平衡医学会. 良性発作性頭めまい症 (BPPV) 診療ガイドライン2023年版. 2023.
- 22) Kim JS, Oh SY, Lee SH, et al. Randomized clinical trial for geotropic horizontal canal benign paroxysmal positional vertigo. *Neurology* 2012;79:700-7.
- 23) Girommetti G, Lapenna R, Panichi R, et al. Residual dizziness after successful repositioning maneuver for idiopathic benign paroxysmal positional vertigo: a review. *Audiol Res* 2017;7:178.
- 24) 日本めまい平衡医学会. 前庭神経炎診療ガイドライン2021年版. 2021.

- 25) Strupp M, Zingler VC, Arbusow V, et al. Methylprednisolone, valacyclovir, or the combination for vestibular neuritis. *N Engl J Med* 2004;351:354-61.
- 26) Fishman JM, Burgess C, Waddell A. Corticosteroids for the treatment of idiopathic acute vestibular dysfunction (vestibular neuritis). *Cochrane Database Syst Rev* 2011;(5):CD008607.
- 27) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）。
- 28) Atzema CL, Grewal K, Lu H, et al. Outcomes among patients discharged from the emergency department with a diagnosis of peripheral vertigo. *Ann Neurol* 2016;79:32-41.
- 29) Grewal K, Austin PC, Kapral MK, et al. Missed strokes using computed tomography imaging in patients with vertigo: population-based cohort study. *Stroke* 2015;46:108-13.
- 30) Hunter BR, Wang AZ, Bucca AW, et al. Efficacy of benzodiazepines or antihistamines for patients with acute vertigo: a systematic review and meta-analysis. *JAMA Neurol* 2022;79:846-55.
- 31) Chi CY, Chen YC, Cheng MT, et al. Diphenhydramine, sodium bicarbonate, or combination for acute peripheral vertigo: a randomized clinical trial. *JAMA Netw Open* 2025;8:e2541472.

2-6 脳卒中を疑う症状

- 1) The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995;333:1581-7.
- 2) Goldstein LB, Simel DL. Is this patient having a stroke? *JAMA* 2005;293:2391-402.
- 3) Amin HP, Madsen TE, Bravata DM, et al. Diagnosis, workup, risk reduction of transient ischemic attack in the emergency department setting: a scientific statement from the American Heart Association. *Stroke* 2023;54:e109-21.
- 4) Johnston SC, Rothwell PM, Nguyen-Huynh MN, et al. Validation and refinement of scores to predict very early stroke risk after transient ischaemic attack. *Lancet* 2007;369:283-92.
- 5) 日本脳卒中学会 脳卒中医療向上・社会保険委員会 静注血栓溶解療法指針改訂部会. 静注血栓溶解（rt-PA）療法 適正治療指針 第三版. 脳卒中 2019;41:205-46.
- 6) Martin-Schild S, Albright KC, Tanksley J, et al. Zero on the NIHSS does not equal the absence of stroke. *Ann Emerg Med* 2011;57:42-5.
- 7) Zhang X, Kedar S, Lynn MJ, Newman NJ, Biousse V. Homonymous hemianopias: clinical-anatomic correlations in 904 cases. *Neurology* 2006;66:906-10.
- 8) Suzuki K, Nakajima N, Kunitomo K, et al. Emergent large vessel occlusion screen is an ideal prehospital scale to avoid missing endovascular therapy in acute stroke. *Stroke* 2018;49:2096-101.
- 9) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）。
- 10) Emberson J, Lees KR, Lyden P, et al. Effect of treatment delay, age, and stroke severity on the effects of intravenous thrombolysis with alteplase for acute ischaemic stroke: a meta-analysis of individual patient data from randomised trials. *Lancet* 2014;384:1929-35.
- 11) Thomalla G, Simonsen CZ, Boutitie F, et al. MRI-guided thrombolysis for stroke with unknown time of onset. *N Engl J Med* 2018;379:611-22.
- 12) 日本脳卒中学会 脳卒中ガイドライン委員会編. 脳卒中治療ガイドライン2021〔改訂2025〕. 協和企画；2025.
- 13) Goyal M, Menon BK, van Zwam WH, et al. Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016;387:1723-31.
- 14) Nogueira RG, Jadhav AP, Haussen DC, et al. Thrombectomy 6 to 24 hours after stroke with a mismatch between deficit and infarct. *N Engl J Med* 2018;378:11-21.

- 15) 日本脳卒中学会, 日本脳神経外科学会, 日本脳神経血管内治療学会. 経皮経管の脳血栓回収用機器 適正使用指針 第5版. 脳卒中 2024;46:181-216.
- 16) Anderson CS, Heeley E, Huang Y, et al. Rapid blood-pressure lowering in patients with acute intracerebral hemorrhage. *N Engl J Med* 2013;368:2355-65.
- 17) Brott T, Broderick J, Kothari R, et al. Early hemorrhage growth in patients with intracerebral hemorrhage. *Stroke* 1997;28:1-5.
- 18) Ma L, Hu X, Song L, et al. The third Intensive Care Bundle with Blood Pressure Reduction in Acute Cerebral Haemorrhage Trial (INTERACT3): an international, stepped wedge cluster randomised controlled trial. *Lancet* 2023;402:27-40.
- 19) Sprigg N, Flaherty K, Appleton JP, et al. Tranexamic acid for hyperacute primary IntraCerebral Haemorrhage (TICH-2): an international randomised, placebo-controlled, phase 3 superiority trial. *Lancet* 2018;391:2107-15.
- 20) Szeto CLC, Hui KF. Spontaneous spinal epidural hematoma mimicking stroke and its outcome post intravenous thrombolysis. *Cerebrovasc Dis* 2022;51:265-9.
- 21) Zinkstok SM, Engelter ST, Gensicke H, et al. Safety of thrombolysis in stroke mimics: results from a multicenter cohort study. *Stroke* 2013;44:1080-4.
- 22) Tarnutzer AA, Lee SH, Robinson KA, Wang Z, Edlow JA, Newman-Toker DE. ED misdiagnosis of cerebrovascular events in the era of modern neuroimaging: a meta-analysis. *Neurology* 2017;88:1468-77.
- 23) Chen HS, Cui Y, Zhou ZH, et al. Dual antiplatelet therapy vs alteplase for patients with minor nondisabling acute ischemic stroke: the ARAMIS randomized clinical trial. *JAMA* 2023;329:2135-44.
- 24) Fahimi J, Navi BB, Kamel H. Potential misdiagnoses of Bell's palsy in the emergency department. *Ann Emerg Med* 2014;63:428-34.
- 25) Baugh RF, Basura GJ, Ishii LE, et al. Clinical practice guideline: Bell's palsy. *Otolaryngol Head Neck Surg* 2013;149(3 Suppl):S1-27.
- 26) Park H, Kim YJ, Kim WY. Need for routine brain magnetic resonance imaging for unilateral facial palsy in emergency department. *Diagnostics (Basel)* 2025;15:2135.

2-7 脱力・運動麻痺

-
- 1) 日本神経学会監修, ギラン・バレー症候群, フィッシャー症候群診療ガイドライン作成委員会編. ギラン・バレー症候群, フィッシャー症候群診療ガイドライン2024. 南江堂; 2024.
 - 2) Leonhard SE, Mandarakas MR, Gondim FAA, et al. Diagnosis and management of Guillain-Barré syndrome in ten steps. *Nat Rev Neurol* 2019;15:671-83.
 - 3) Ahn UM, Ahn NU, Buchowski JM, et al. Cauda equina syndrome secondary to lumbar disc herniation: a meta-analysis of surgical outcomes. *Spine* 2000;25:1515-22.
 - 4) Miyakoshi N, Suda K, Kudo D, et al. A nationwide survey on the incidence and characteristics of traumatic spinal cord injury in Japan in 2018. *Spinal Cord* 2021;59:626-34.
 - 5) Ruts L, Drenthen J, Jongen JLM, et al. Pain in Guillain-Barré syndrome: a long-term follow-up study. *Neurology* 2010;75:1439-47.
 - 6) Walgaard C, Lingsma HF, Ruts L, et al. Prediction of respiratory insufficiency in Guillain-Barré syndrome. *Ann Neurol* 2010;67:781-7.
 - 7) 献血ヴェノグロブリンIH 5%静注の電子添文 (KEGG MEDICUSで2026年10月に確認) .
 - 8) Lin SH, Lin YF, Chen DT, Chu P, Hsu CW, Halperin ML. Laboratory tests to determine the cause of hypokalemia and paralysis. *Arch Intern Med* 2004;164:1561-6.
 - 9) 日本神経学会監修, 重症筋無力症／ランバート・イートン筋無力症候群診療ガイドライン作成委員会編. 重症筋無力症／ランバート・イートン筋無力症候群診療ガイドライン2022. 南江堂; 2022.

- 10) Giannoccaro MP, Paolucci M, Zenesini C, et al. Comparison of ice pack test and single-fiber EMG diagnostic accuracy in patients referred for myasthenic ptosis. *Neurology* 2020;95:e1800-6.
- 11) メスチノン錠60 mgの電子添文（2025年10月改訂 第2版。KEGG MEDICUSで2026年10月に確認）。
- 12) スキサメトニウム注、エスラックス静注の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 13) Hughes RA, Brassington R, Gunn AA, van Doorn PA. Corticosteroids for Guillain-Barré syndrome. *Cochrane Database Syst Rev* 2016;10:CD001446.
- 14) Hughes RA, Swan AV, van Doorn PA. Intravenous immunoglobulin for Guillain-Barré syndrome. *Cochrane Database Syst Rev* 2014;(9):CD002063.

第3部 胸部の症候と循環

3-1 胸痛（背部痛を含む）

- 1) 日本循環器学会, ほか. 急性冠症候群ガイドライン（2018年改訂版）. 2019.
- 2) Welch RD, Zalski RJ, Frederick PD, et al. Prognostic value of a normal or nonspecific initial electrocardiogram in acute myocardial infarction. *JAMA* 2001;286:1977-84.
- 3) Hofmann R, James SK, Jernberg T, et al. Oxygen therapy in suspected acute myocardial infarction. *N Engl J Med* 2017;377:1240-9.
- 4) Thygesen K, Alpert JS, Jaffe AS, et al. Fourth universal definition of myocardial infarction (2018). *Circulation* 2018;138:e618-51.
- 5) Canto JG, Shlipak MG, Rogers WJ, et al. Prevalence, clinical characteristics, and mortality among patients with myocardial infarction presenting without chest pain. *JAMA* 2000;283:3223-9.
- 6) Chew DP, Lambrakis K, Blyth A, et al. A randomized trial of a 1-hour troponin T protocol in suspected acute coronary syndromes: the Rapid Assessment of Possible Acute Coronary Syndrome in the Emergency Department With High-Sensitivity Troponin T Study (RAPID-TnT). *Circulation* 2019;140:1543-56.
- 7) Byrne RA, Rossello X, Coughlan JJ, et al. 2023 ESC Guidelines for the management of acute coronary syndromes. *Eur Heart J* 2023;44:3720-826.
- 8) Backus BE, Six AJ, Kelder JC, et al. A prospective validation of the HEART score for chest pain patients at the emergency department. *Int J Cardiol* 2013;168:2153-8.
- 9) Poldervaart JM, Reitsma JB, Backus BE, et al. Effect of using the HEART score in patients with chest pain in the emergency department: a stepped-wedge, cluster randomized trial. *Ann Intern Med* 2017;166:689-97.
- 10) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）。
- 11) ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988;2:349-60.
- 12) Amsterdam EA, Wenger NK, Brindis RG, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: executive summary. *Circulation* 2014;130:2354-94.
- 13) Henrikson CA, Howell EE, Bush DE, et al. Chest pain relief by nitroglycerin does not predict active coronary artery disease. *Ann Intern Med* 2003;139:979-86.
- 14) Meine TJ, Roe MT, Chen AY, et al. Association of intravenous morphine use and outcomes in acute coronary syndromes: results from the CRUSADE Quality Improvement Initiative. *Am Heart J* 2005;149:1043-9.

- 15) 日本循環器学会, 日本心臓血管外科学会, 日本胸部外科学会, 日本血管外科学会. 2020年改訂版 大動脈瘤・大動脈解離診療ガイドライン. 2020.
- 16) Nazerian P, Mueller C, Soeiro AM, et al. Diagnostic accuracy of the aortic dissection detection risk score plus D-dimer for acute aortic syndromes: the ADvISED prospective multicenter study. *Circulation* 2018;137:250-8.
- 17) Kurabayashi M, Miwa N, Ueshima D, et al. Factors leading to failure to diagnose acute aortic dissection in the emergency room. *J Cardiol* 2011;58:287-93.
- 18) Klompas M. Does this patient have an acute thoracic aortic dissection? *JAMA* 2002;287:2262-72.
- 19) Hagan PG, Nienaber CA, Isselbacher EM, et al. The International Registry of Acute Aortic Dissection (IRAD): new insights into an old disease. *JAMA* 2000;283:897-903.
- 20) 日本循環器学会, 日本肺高血圧・肺循環学会. 2025年改訂版 肺血栓症・深部静脈血栓症および肺高血圧症に関するガイドライン. 2025.
- 21) Stein PD, Beemath A, Matta F, et al. Clinical characteristics of patients with acute pulmonary embolism: data from PIOPED II. *Am J Med* 2007;120:871-9.
- 22) Wells PS, Anderson DR, Rodger M, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000;83:416-20.
- 23) Freund Y, Cachanado M, Aubry A, et al. Effect of the Pulmonary Embolism Rule-Out Criteria on subsequent thromboembolic events among low-risk emergency department patients: the PROPER randomized clinical trial. *JAMA* 2018;319:559-66.
- 24) Kouritas VK, Papagiannopoulos K, Lazaridis G, et al. Pneumomediastinum. *J Thorac Dis* 2015;7(Suppl 1):S44-9.
- 25) Imazio M, Gaita F, LeWinter M. Evaluation and treatment of pericarditis: a systematic review. *JAMA* 2015;314:1498-506.
- 26) Adler Y, Charron P, Imazio M, et al. 2015 ESC Guidelines for the diagnosis and management of pericardial diseases. *Eur Heart J* 2015;36:2921-64.
- 27) Imazio M, Brucato A, Cemin R, et al. A randomized trial of colchicine for acute pericarditis. *N Engl J Med* 2013;369:1522-8.
- 28) 日本整形外科学会, 日本腰痛学会監修. 腰痛診療ガイドライン2019 改訂第2版. 南江堂; 2019.
- 29) Pope JH, Aufderheide TP, Ruthazer R, et al. Missed diagnoses of acute cardiac ischemia in the emergency department. *N Engl J Med* 2000;342:1163-70.
- 30) Fanaroff AC, Rymer JA, Goldstein SA, et al. Does this patient with chest pain have acute coronary syndrome? The Rational Clinical Examination systematic review. *JAMA* 2015;314:1955-65.
- 31) Kubica J, Adamski P, Ostrowska M, et al. Morphine delays and attenuates ticagrelor exposure and action in patients with myocardial infarction: the randomized, double-blind, placebo-controlled IMPRESSION trial. *Eur Heart J* 2016;37:245-52.
- 32) Duarte GS, Nunes-Ferreira A, Rodrigues FB, et al. Morphine in acute coronary syndrome: systematic review and meta-analysis. *BMJ Open* 2019;9:e025232.
- 33) Furtado RHM, Nicolau JC, Guo J, et al. Morphine and cardiovascular outcomes among patients with non-ST-segment elevation acute coronary syndromes undergoing coronary angiography. *J Am Coll Cardiol* 2020;75:289-300.

3-2 呼吸困難・SpO₂低下

- 1) Maisel AS, Krishnaswamy P, Nowak RM, et al. Rapid measurement of B-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med* 2002;347:161-7.

- 2) O'Driscoll BR, Howard LS, Earis J, et al. BTS guideline for oxygen use in adults in healthcare and emergency settings. *Thorax* 2017;72(Suppl 1):ii1-ii90.
- 3) Cortelli P, Calandra-Buonaura G, Benarroch EE, et al. Stridor in multiple system atrophy: consensus statement on diagnosis, prognosis, and treatment. *Neurology* 2019;93:630-9.
- 4) 日本アレルギー学会. アナフィラキシーガイドライン2022. 2022.
- 5) Davidson K, Shojaaee S. Managing massive hemoptysis. *Chest* 2020;157:77-88.
- 6) Radchenko C, Alraiyes AH, Shojaaee S. A systematic approach to the management of massive hemoptysis. *J Thorac Dis* 2017;9(Suppl 10):S1069-86.
- 7) 日本呼吸器学会COPDガイドライン第6版作成委員会編. COPD（慢性閉塞性肺疾患）診断と治療のためのガイドライン2022 [第6版]. メディカルレビュー社; 2022.
- 8) McCanny P, Bennett K, Staunton P, et al. Venous vs arterial blood gases in the assessment of patients presenting with an exacerbation of chronic obstructive pulmonary disease. *Am J Emerg Med* 2012;30:896-900.
- 9) Alrajhi K, Woo MY, Vaillancourt C. Test characteristics of ultrasonography for the detection of pneumothorax: a systematic review and meta-analysis. *Chest* 2012;141:703-8.
- 10) Self WH, Courtney DM, McNaughton CD, et al. High discordance of chest x-ray and computed tomography for detection of pulmonary opacities in ED patients: implications for diagnosing pneumonia. *Am J Emerg Med* 2013;31:401-5.
- 11) Maw AM, Hassanin A, Ho PM, et al. Diagnostic accuracy of point-of-care lung ultrasonography and chest radiography in adults with symptoms suggestive of acute decompensated heart failure: a systematic review and meta-analysis. *JAMA Netw Open* 2019;2:e190703.
- 12) 日本循環器学会, 日本心不全学会. 2025年改訂版 心不全診療ガイドライン. 2025.
- 13) Januzzi JL, van Kimmenade R, Lainchbury J, et al. NT-proBNP testing for diagnosis and short-term prognosis in acute destabilized heart failure: an international pooled analysis of 1256 patients. *Eur Heart J* 2006;27:330-7.
- 14) Wang CS, FitzGerald JM, Schulzer M, et al. Does this dyspneic patient in the emergency department have congestive heart failure? *JAMA* 2005;294:1944-56.
- 15) 日本アレルギー学会喘息ガイドライン専門部会監修. 喘息予防・管理ガイドライン2024. 協和企画; 2024.
- 16) Callahan M. Hypoxic hazards of traditional paper bag rebreathing in hyperventilating patients. *Ann Emerg Med* 1989;18:622-8.
- 17) Lawn ND, Fletcher DD, Henderson RD, et al. Anticipating mechanical ventilation in Guillain-Barré syndrome. *Arch Neurol* 2001;58:893-8.
- 18) Uchiyama H, Suda T, Nakamura Y, et al. Alterations in smoking habits are associated with acute eosinophilic pneumonia. *Chest* 2008;133:1174-80.
- 19) De Giacomo F, Vassallo R, Yi ES, Ryu JH. Acute eosinophilic pneumonia. Causes, diagnosis, and management. *Am J Respir Crit Care Med* 2018;197:728-36.
- 20) Lichtenstein DA, Mezière GA. Relevance of lung ultrasound in the diagnosis of acute respiratory failure: the BLUE protocol. *Chest* 2008;134:117-25.
- 21) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）.
- 22) Leuppi JD, Schuetz P, Bingisser R, et al. Short-term vs conventional glucocorticoid therapy in acute exacerbations of chronic obstructive pulmonary disease: the REDUCE randomized clinical trial. *JAMA* 2013;309:2223-31.
- 23) Couturaud F, Bertoletti L, Pastre J, et al. Prevalence of pulmonary embolism among patients with COPD hospitalized with acutely worsening respiratory symptoms. *JAMA* 2021;325:59-68.
- 24) Gibbs KW, Semler MW, Driver BE, et al. Noninvasive ventilation for preoxygenation during emergency intubation. *N Engl J Med* 2024;390:2165-77.

- 25) Fartoukh M, Khoshnood B, Parrot A, et al. Early prediction of in-hospital mortality of patients with hemoptysis: an approach to defining severe hemoptysis. *Respiration* 2012;83:106-14.
- 26) Kinoshita T, Ohbe H, Matsui H, et al. Effect of tranexamic acid on mortality in patients with haemoptysis: a nationwide study. *Crit Care* 2019;23:347.
- 27) Hampson NB. Pulse oximetry in severe carbon monoxide poisoning. *Chest* 1998;114:1036-41.
- 28) Barker SJ, Tremper KK, Hyatt J. Effects of methemoglobinemia on pulse oximetry and mixed venous oximetry. *Anesthesiology* 1989;70:112-7.
- 29) Global Initiative for Asthma. Global strategy for asthma management and prevention, 2026 update. 2026.
- 30) Ratto D, Alfaro C, Sipsey J, Glovsky MM, Sharma OP. Are intravenous corticosteroids required in status asthmaticus? *JAMA* 1988;260:527-9.
- 31) Harrison BD, Stokes TC, Hart GJ, Vaughan DA, Ali NJ, Robinson AA. Need for intravenous hydrocortisone in addition to oral prednisolone in patients admitted to hospital with severe asthma without ventilatory failure. *Lancet* 1986;1:181-4.
- 32) Rowe BH, Spooner C, Ducharme FM, Bretzlaff JA, Bota GW. Early emergency department treatment of acute asthma with systemic corticosteroids. *Cochrane Database Syst Rev* 2001; (1):CD002178.

3-3 頻脈・徐脈

- 1) Soar J, Böttiger BW, Carli P, et al. European Resuscitation Council Guidelines 2025: adult advanced life support. *Resuscitation* 2025;215:110769.
- 2) 日本循環器学会, 日本不整脈心電学会. 2020年改訂版 不整脈薬物治療ガイドライン. 2020.
- 3) 日本麻酔科学会 安全委員会. 安全な鎮静のためのプラクティカルガイド. 2021年11月26日制定. https://anesth.or.jp/files/pdf/practical_guide_for_safe_sedation_20220111.pdf (2026年10月7日閲覧)
- 4) Appelboom A, Reuben A, Mann C, et al. Postural modification to the standard Valsalva manoeuvre for emergency treatment of supraventricular tachycardias (REVERT): a randomised controlled trial. *Lancet* 2015;386:1747-53.
- 5) 各薬剤の電子添文 (KEGG MEDICUSで2026年9月に確認) .
- 6) 日本循環器学会, 日本不整脈心電学会. 2024年JCS/JHRSガイドライン フォーカスアップデート版 不整脈治療. 2024.
- 7) Akhtar M, Shenasa M, Jazayeri M, et al. Wide QRS complex tachycardia: reappraisal of a common clinical problem. *Ann Intern Med* 1988;109:905-12.
- 8) Baerman JM, Morady F, DiCarlo LA Jr, et al. Differentiation of ventricular tachycardia from supraventricular tachycardia with aberration: value of the clinical history. *Ann Emerg Med* 1987;16:40-3.
- 9) Brugada P, Brugada J, Mont L, et al. A new approach to the differential diagnosis of a regular tachycardia with a wide QRS complex. *Circulation* 1991;83:1649-59.
- 10) 日本循環器学会, 日本不整脈心電学会. 不整脈非薬物治療ガイドライン (2018年改訂版) . 2018.
- 11) 日本循環器学会, ほか. 2016年版 心臓サルコイドーシスの診療ガイドライン (JCS 2016) .
- 12) 日本循環器学会, 日本血管外科学会. 2022年改訂版 末梢動脈疾患ガイドライン. 2022.
- 13) Kusumoto FM, Schoenfeld MH, Barrett C, et al. 2018 ACC/AHA/HR guideline on the evaluation and management of patients with bradycardia and cardiac conduction delay. *Circulation* 2019;140:e382-482.
- 14) Alabed S, Sabouni A, Providencia R, Atallah E, Qintar M, Chico TJ. Adenosine versus intravenous calcium channel antagonists for supraventricular tachycardia. *Cochrane Database Syst Rev* 2017;(10):CD005154.

- 15) DiMarco JP, Miles W, Akhtar M, et al. Adenosine for paroxysmal supraventricular tachycardia: dose ranging and comparison with verapamil. *Ann Intern Med* 1990;113:104-10.

3-4 高血圧緊急症

- 1) 日本高血圧学会. 高血圧管理・治療ガイドライン2025. 2025. (英語版: *Hypertens Res* 2026;49:9-235)
- 2) Astarita A, Covella M, Vallelonga F, et al. Hypertensive emergencies and urgencies in emergency departments: a systematic review and meta-analysis. *J Hypertens* 2020;38:1203-10.
- 3) Zampaglione B, Pascale C, Marchisio M, Cavallo-Perin P. Hypertensive urgencies and emergencies: prevalence and clinical presentation. *Hypertension* 1996;27:144-7.
- 4) van den Born BH, Lip GYH, Brguljan-Hitij J, et al. ESC Council on hypertension position document on the management of hypertensive emergencies. *Eur Heart J Cardiovasc Pharmacother* 2019;5:37-46.
- 5) van den Born BJ, Honnebieer UP, Koopmans RP, et al. Microangiopathic hemolysis and renal failure in malignant hypertension. *Hypertension* 2005;45:246-51.
- 6) Shibagaki Y, Fujita T. Thrombotic microangiopathy in malignant hypertension and hemolytic uremic syndrome (HUS)/thrombotic thrombocytopenic purpura (TTP): can we differentiate one from the other? *Hypertens Res* 2005;28:89-95.
- 7) Siddiqi TJ, Usman MS, Rashid AM, et al. Clinical outcomes in hypertensive emergency: a systematic review and meta-analysis. *J Am Heart Assoc* 2023;12:e029355.
- 8) Patel KK, Young L, Howell EH, et al. Characteristics and outcomes of patients presenting with hypertensive urgency in the office setting. *JAMA Intern Med* 2016;176:981-8.
- 9) Grassi D, O'Flaherty M, Pellizzari M, et al. Hypertensive urgencies in the emergency department: evaluating blood pressure response to rest and to antihypertensive drugs with different profiles. *J Clin Hypertens (Greenwich)* 2008;10:662-7.
- 10) Park SK, Lee DY, Kim WJ, et al. Comparing the clinical efficacy of resting and antihypertensive medication in patients of hypertensive urgency: a randomized, control trial. *J Hypertens* 2017;35:1474-80.
- 11) 各薬剤の電子添文 (KEGG MEDICUSで2026年9月に確認)。
- 12) 日本循環器学会, 日本心臓血管外科学会, 日本胸外科学会, 日本血管外科学会. 2020年改訂版 大動脈瘤・大動脈解離診療ガイドライン. 2020.
- 13) 日本高血圧学会高血圧治療ガイドライン作成委員会編. 高血圧治療ガイドライン2019. ライフサイエンス出版; 2019.
- 14) Cotter G, Metzkor E, Kaluski E, et al. Randomised trial of high-dose isosorbide dinitrate plus low-dose furosemide versus high-dose furosemide plus low-dose isosorbide dinitrate in severe pulmonary oedema. *Lancet* 1998;351:389-93.
- 15) Berbenetz N, Wang Y, Brown J, et al. Non-invasive positive pressure ventilation (CPAP or bilevel NPPV) for cardiogenic pulmonary oedema. *Cochrane Database Syst Rev* 2019;4:CD005351.
- 16) Gray A, Goodacre S, Newby DE, et al. Noninvasive ventilation in acute cardiogenic pulmonary edema. *N Engl J Med* 2008;359:142-51.

第4部 腹部と消化器の症候

4-1 腹痛

- 1) 急性腹症診療ガイドライン出版委員会編. 急性腹症診療ガイドライン2015. 医学書院; 2015.

- 2) Lewis LM, Banet GA, Blanda M, Hustey FM, Meldon SW, Gerson LW. Etiology and clinical course of abdominal pain in senior patients: a prospective, multicenter study. *J Gerontol A Biol Sci Med Sci* 2005;60:1071-6.
- 3) 日本循環器学会, 日本心臓血管外科学会, 日本胸部外科学会, 日本血管外科学会. 2020年改訂版 大動脈瘤・大動脈解離診療ガイドライン. 2020.
- 4) Bala M, Catena F, Kashuk J, et al. Acute mesenteric ischemia: updated guidelines of the World Society of Emergency Surgery. *World J Emerg Surg* 2022;17:54.
- 5) 日本泌尿器科学会編. 急性陰嚢症診療ガイドライン 2014年版. 金原出版; 2014.
- 6) Ho WK, Wang YF, Wu HH, et al. Ruptured corpus luteum with hemoperitoneum: case characteristics and demographic changes over time. *Taiwan J Obstet Gynecol* 2009;48:108-12.
- 7) Bottomley C, Bourne T. Diagnosis and management of ovarian cyst accidents. *Best Pract Res Clin Obstet Gynaecol* 2009;23:711-24.
- 8) Umpierrez G, Freire AX. Abdominal pain in patients with hyperglycemic crises. *J Crit Care* 2002;17:63-7.
- 9) Yoshida H, Mamada Y, Taniai N, Uchida E. Spontaneous ruptured hepatocellular carcinoma. *Hepatol Res* 2016;46:13-21.
- 10) Wagner JM, McKinney WP, Carpenter JL. Does this patient have appendicitis? *JAMA* 1996;276:1589-94.
- 11) Ohle R, O'Reilly P, O'Brien KK, Fahey T, Dimitrov BD. The Alvarado score for predicting acute appendicitis: a systematic review. *BMC Med* 2011;9:139.
- 12) Cardall T, Glasser J, Guss DA. Clinical value of the total white blood cell count and temperature in the evaluation of patients with suspected appendicitis. *Acad Emerg Med* 2004;11:1021-7.
- 13) Andersson RE. Meta-analysis of the clinical and laboratory diagnosis of appendicitis. *Br J Surg* 2004;91:28-37.
- 14) Terasawa T, Blackmore CC, Bent S, Kohlwes RJ. Systematic review: computed tomography and ultrasonography to detect acute appendicitis in adults and adolescents. *Ann Intern Med* 2004;141:537-46.
- 15) Taylor MR, Lalani N. Adult small bowel obstruction. *Acad Emerg Med* 2013;20:528-44.
- 16) Ten Broek RPG, Krielen P, Di Saverio S, et al. Bologna guidelines for diagnosis and management of adhesive small bowel obstruction (ASBO): 2017 update of the evidence-based guidelines from the world society of emergency surgery ASBO working group. *World J Emerg Surg* 2018;13:24.
- 17) 日本消化管学会. 大腸憩室症（憩室出血・憩室炎）ガイドライン. 日本消化管学会雑誌 2017;1(Suppl).
- 18) Daniels L, Ünlü Ç, de Korte N, et al. Randomized clinical trial of observational versus antibiotic treatment for a first episode of CT-proven uncomplicated acute diverticulitis. *Br J Surg* 2017;104:52-61.
- 19) Singh AK, Gervais DA, Hahn PF, Sagar P, Mueller PR, Novelline RA. Acute epiploic appendagitis and its mimics. *Radiographics* 2005;25:1521-34.
- 20) Schnedl WJ, Krause R, Tafelt E, Tillich M, Lipp RW, Wallner-Liebmann SJ. Insights into epiploic appendagitis. *Nat Rev Gastroenterol Hepatol* 2011;8:45-9.
- 21) Pawlak M, East B, de Beaux AC. Algorithm for management of an incarcerated inguinal hernia in the emergency settings with manual reduction. Taxis, the technique and its safety. *Hernia* 2021;25:1253-8.
- 22) Gruber PJ, Silverman RA, Gottesfeld S, Flaster E. Presence of fever and leukocytosis in acute cholecystitis. *Ann Emerg Med* 1996;28:273-7.
- 23) 急性腹症診療ガイドライン出版委員会編. 急性腹症診療ガイドライン2025 第2版. 医学書院; 2025.

- 24) Manterola C, Vial M, Moraga J, Astudillo P. Analgesia in patients with acute abdominal pain. *Cochrane Database Syst Rev* 2011;(1):CD005660.
- 25) Ranji SR, Goldman LE, Simel DL, Shojania KG. Do opiates affect the clinical evaluation of patients with acute abdominal pain? *JAMA* 2006;296:1764-74.
- 26) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認。グリセリン浣腸はJAPICで2026年10月に確認）。
- 27) Toorenvliet BR, Bakker RF, Flu HC, Merkus JW, Hamming JF, Breslau PJ. Standard outpatient re-evaluation for patients not admitted to the hospital after emergency department evaluation for acute abdominal pain. *World J Surg* 2010;34:480-6.
- 28) Thomson H, Francis DM. Abdominal-wall tenderness: a useful sign in the acute abdomen. *Lancet* 1977;2:1053-4.
- 29) Azhar B, Patel SR, Holt PJ, Hinchliffe RJ, Thompson MM, Karthikesalingam A. Misdiagnosis of ruptured abdominal aortic aneurysm: systematic review and meta-analysis. *J Endovasc Ther* 2014;21:568-75.
- 30) CODA Collaborative, Flum DR, Davidson GH, et al. A randomized trial comparing antibiotics with appendectomy for appendicitis. *N Engl J Med* 2020;383:1907-19.
- 31) Salminen P, Tuominen R, Paaanen H, et al. Five-year follow-up of antibiotic therapy for uncomplicated acute appendicitis in the APPAC randomized clinical trial. *JAMA* 2018;320:1259-65.

4-2 吐血・下血

- 1) de Franchis R, Bosch J, Garcia-Tsao G, et al. Baveno VII - Renewing consensus in portal hypertension. *J Hepatol* 2022;76:959-74.
- 2) McGee S, Abernethy WB 3rd, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA* 1999;281:1022-9.
- 3) Demetriades D, Chan LS, Bhasin P, et al. Relative bradycardia in patients with traumatic hypotension. *J Trauma* 1998;45:534-9.
- 4) 日本輸血・細胞治療学会ガイドライン委員会. 輸血療法実践ガイド. 日本輸血細胞治療学会誌 2026;72:521-70.
- 5) 日本麻酔科学会, 日本輸血・細胞治療学会. 危機的出血への対応ガイドライン（2025年4月改訂）. 2025.
- 6) Srygley FD, Gerardo CJ, Tran T, Fisher DA. Does this patient have a severe upper gastrointestinal bleed? *JAMA* 2012;307:1072-9.
- 7) Oakland K, Chadwick G, East JE, et al. Diagnosis and management of acute lower gastrointestinal bleeding: guidelines from the British Society of Gastroenterology. *Gut* 2019;68:776-89.
- 8) 藤城光弘, 井口幹崇, 小野敏嗣, ほか. 非静脈瘤性上部消化管出血における内視鏡診療ガイドライン（第2版）. 日本消化器内視鏡学会雑誌 2024;66:1515-38.
- 9) Umezawa S, Nagata N, Arimoto J, et al. Contrast-enhanced CT for colonic diverticular bleeding before colonoscopy: a prospective multicenter study. *Radiology* 2018;288:755-61.
- 10) オメプラゾール注射用20 mg 電子添文（KEGG MEDICUSで2026年10月に確認）。
- 11) Kanno T, Yuan Y, Tse F, et al. Proton pump inhibitor treatment initiated prior to endoscopic diagnosis in upper gastrointestinal bleeding. *Cochrane Database Syst Rev* 2022;(1):CD005415.
- 12) Barkun AN, Almadi M, Kuipers EJ, et al. Management of nonvariceal upper gastrointestinal bleeding: guideline recommendations from the International Consensus Group. *Ann Intern Med* 2019;171:805-22.

- 13) Sung JJ, Lau JY, Ching JY, et al. Continuation of low-dose aspirin therapy in peptic ulcer bleeding: a randomized trial. *Ann Intern Med* 2010;152:1-9.
- 14) Blatchford O, Murray WR, Blatchford M. A risk score to predict need for treatment for upper-gastrointestinal haemorrhage. *Lancet* 2000;356:1318-21.
- 15) Stanley AJ, Laine L, Dalton HR, et al. Comparison of risk scoring systems for patients presenting with upper gastrointestinal bleeding: international multicentre prospective study. *BMJ* 2017;356:i6432.
- 16) Stanley AJ, Ashley D, Dalton HR, et al. Outpatient management of patients with low-risk upper-gastrointestinal haemorrhage: multicentre validation and prospective evaluation. *Lancet* 2009;373:42-7.
- 17) Laine L, Barkun AN, Saltzman JR, et al. ACG clinical guideline: upper gastrointestinal and ulcer bleeding. *Am J Gastroenterol* 2021;116:899-917.
- 18) Oakland K, Kothiwale S, Forehand T, et al. External validation of the Oakland score to assess safe hospital discharge among adult patients with acute lower gastrointestinal bleeding in the US. *JAMA Netw Open* 2020;3:e209630.
- 19) Oakland K, Jairath V, Uberoi R, et al. Derivation and validation of a novel risk score for safe discharge after acute lower gastrointestinal bleeding: a modelling study. *Lancet Gastroenterol Hepatol* 2017;2:635-43.
- 20) Kinjo K, Aoki T, Kobayashi K, et al. Validation of British Society of Gastroenterology guidelines for acute lower GI bleeding from 8956 cases in Japan. *Gastrointest Endosc* 2025;101:1131-44.
- 21) Alshamsi F, Jaeschke R, Baw B, Alhazzani W. Prophylactic endotracheal intubation in patients with upper gastrointestinal bleeding undergoing endoscopy: a systematic review and meta-analysis. *Saudi J Med Med Sci* 2017;5:201-9.

4-3 嘔吐・下痢

- 1) Mahajan P, Basu T, Pai CW, et al. Factors associated with potentially missed diagnosis of appendicitis in the emergency department. *JAMA Netw Open* 2020;3:e200612.
- 2) 日本循環器学会, ほか. 2023年改訂版 心筋炎の診断・治療に関するガイドライン. 2023.
- 3) Kiriya S, Takada T, Strasberg SM, et al. New diagnostic criteria and severity assessment of acute cholangitis in revised Tokyo Guidelines. *J Hepatobiliary Pancreat Sci* 2012;19:548-56.
- 4) 食品安全委員会. ファクトシート (ブドウ球菌食中毒 令和3年3月30日更新、ウエルシュ菌食中毒 令和7年4月11日更新、カンピロバクター 平成28年9月30日更新、セレウス菌食中毒 令和3年3月30日更新、ヒスタミン 令和3年3月30日更新) .
- 5) 厚生労働省. 腸管出血性大腸菌Q&A (厚生労働省のページ、2026年10月に閲覧) .
- 6) 厚生労働省. アニサキスによる食中毒を予防しましょう (厚生労働省のページ、2026年10月に閲覧) .
- 7) 厚生労働省. ノロウイルスに関するQ&A (最終改定 令和3年11月19日) .
- 8) 厚生労働省. 自然毒のリスクプロファイル：魚類：シガテラ毒 (厚生労働省のページ、2026年10月に閲覧) .
- 9) McGee S, Abernethy WB 3rd, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA* 1999;281:1022-9.
- 10) 厚生労働省健康・生活衛生局感染症対策部感染症対策課. 抗微生物薬適正使用の手引き 第四版 医科・外来編. 2026.
- 11) Shane AL, Mody RK, Crump JA, et al. 2017 Infectious Diseases Society of America clinical practice guidelines for the diagnosis and management of infectious diarrhea. *Clin Infect Dis* 2017;65:e45-80.
- 12) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認。プリンペラン注射液、ナウゼリン錠、ピオフェルミンR錠、ミヤBM錠) .

- 13) Furyk JS, Meek RA, Egerton-Warburton D. Drugs for the treatment of nausea and vomiting in adults in the emergency department setting. *Cochrane Database Syst Rev* 2015;(9):CD010106.
- 14) アキネトン注射液5 mgの電子添文 (KEGG MEDICUSで2026年10月に確認) .
- 15) Collinson S, Deans A, Padua-Zamora A, et al. Probiotics for treating acute infectious diarrhoea. *Cochrane Database Syst Rev* 2020;(12):CD003048.

4-4 側腹部痛・腰背部痛

- 1) 日本泌尿器科学会, 日本尿路結石症学会, 日本泌尿器内視鏡・ロボティクス学会編. 尿路結石症診療ガイドライン 第3版. 2023.
- 2) Kobayashi T, Nishizawa K, Mitsumori K, Ogura K. Impact of date of onset on the absence of hematuria in patients with acute renal colic. *J Urol* 2003;170:1093-6.
- 3) Rubano E, Mehta N, Caputo W, Paladino L, Sinert R. Systematic review: emergency department bedside ultrasonography for diagnosing suspected abdominal aortic aneurysm. *Acad Emerg Med* 2013;20:128-38.
- 4) Bourgault M, Grimbert P, Verret C, et al. Acute renal infarction: a case series. *Clin J Am Soc Nephrol* 2013;8:392-8.
- 5) 石川勲. 運動後の急性腎障害. 日本内科学会雑誌 2014;103:1101-7.
- 6) Moore CL, Bomann S, Daniels B, et al. Derivation and validation of a clinical prediction rule for uncomplicated ureteral stone—the STONE score: retrospective and prospective observational cohort studies. *BMJ* 2014;348:g2191.
- 7) Wang RC, Rodriguez RM, Moghadassi M, et al. External validation of the STONE score, a clinical prediction rule for ureteral stone: an observational multi-institutional study. *Ann Emerg Med* 2016;67:423-32.
- 8) Smith-Bindman R, Aubin C, Bailitz J, et al. Ultrasonography versus computed tomography for suspected nephrolithiasis. *N Engl J Med* 2014;371:1100-10.
- 9) Pathan SA, Mitra B, Straney LD, et al. Delivering safe and effective analgesia for management of renal colic in the emergency department: a double-blind, multigroup, randomised controlled trial. *Lancet* 2016;387:1999-2007.
- 10) 各薬剤の電子添文 (KEGG MEDICUSで2026年10月に確認. ボルタレンサボ、アセリオ静注液、ソセゴン注射液15 mg) .
- 11) Abrahamian FM, Krishnadasan A, Mower WR, Moran GJ, Talan DA. Association of pyuria and clinical characteristics with the presence of urinary tract infection among patients with acute nephrolithiasis. *Ann Emerg Med* 2013;62:526-33.
- 12) 日本感染症学会, 日本化学療法学会. JAID/JSC感染症治療ガイドライン2015—尿路感染症・男性性器感染症—. 日化療誌 2016;64:1-30.
- 13) van Nieuwkoop C, Hoppe BP, Bonten TN, et al. Predicting the need for radiologic imaging in adults with febrile urinary tract infection. *Clin Infect Dis* 2010;51:1266-72.
- 14) Berbari EF, Kanj SS, Kowalski TJ, et al. 2015 Infectious Diseases Society of America (IDSA) clinical practice guidelines for the diagnosis and treatment of native vertebral osteomyelitis in adults. *Clin Infect Dis* 2015;61:e26-46.
- 15) 厚生労働科学研究費補助金難治性疾患政策研究事業 難治性腎障害に関する調査研究班編. エビデンスに基づく急速進行性腎炎症候群 (RPGN) 診療ガイドライン2020. 東京医学社 ; 2020.
- 16) Coll DM, Varanelli MJ, Smith RC. Relationship of spontaneous passage of ureteral calculi to stone size and location as revealed by unenhanced helical CT. *AJR Am J Roentgenol* 2002;178:101-3.
- 17) Miller OF, Kane CJ. Time to stone passage for observed ureteral calculi: a guide for patient education. *J Urol* 1999;162:688-90.

- 18) Katzouraki G, Zubairi AJ, Hershkovich O, Grevitt MP. A prospective study of the role of bladder scanning and post-void residual volume measurement in improving diagnostic accuracy of cauda equina syndrome. *Bone Joint J* 2020;102-B:677-82.
- 19) Todd NV. Bladder ultrasonography in the assessment of cauda equina syndrome in the emergency department: a literature review. *Ann R Coll Surg Engl* 2024;106:9-12.
- 20) Tabrah J, Wilson N, Phillips D, Böhning D. Can digital rectal examination be used to detect cauda equina compression in people presenting with acute cauda equina syndrome? A systematic review and meta-analysis of diagnostic test accuracy studies. *Musculoskelet Sci Pract* 2022;58:102523.
- 21) Davis DP, Wold RM, Patel RJ, et al. The clinical presentation and impact of diagnostic delays on emergency department patients with spinal epidural abscess. *J Emerg Med* 2004;26:285-91.
- 22) Davis DP, Salazar A, Chan TC, Vilke GM. Prospective evaluation of a clinical decision guideline to diagnose spinal epidural abscess in patients who present to the emergency department with spine pain. *J Neurosurg Spine* 2011;14:765-70.
- 23) Ito Z, Harada A, Matsui Y, et al. Can you diagnose for vertebral fracture correctly by plain X-ray? *Osteoporos Int* 2006;17:1584-91.
- 24) Hicks D, Li CY. Management of macroscopic haematuria in the emergency department. *Emerg Med J* 2007;24:385-90.
- 25) 血尿診断ガイドライン改訂委員会編. 血尿診断ガイドライン2023. ライフサイエンス出版; 2023.
- 26) 日本泌尿器科学会編. 男性下部尿路症状・前立腺肥大症診療ガイドライン. リッチヒルメディカル; 2017.
- 27) Mefford JM, Tungate RM, Amini L, et al. A comparison of urolithiasis in the presence and absence of microscopic hematuria in the emergency department. *West J Emerg Med* 2017;18:775-9.
- 28) Worster AS, Bhanich Supapol W. Fluids and diuretics for acute ureteric colic. *Cochrane Database Syst Rev* 2012;(2):CD004926.
- 29) Campschroer T, Zhu X, Vernooij RW, Lock MT. Alpha-blockers as medical expulsive therapy for ureteral stones. *Cochrane Database Syst Rev* 2018;(4):CD008509.
- 30) Meltzer AC, Burrows PK, Wolfson AB, et al. Effect of tamsulosin on passage of symptomatic ureteral stones: a randomized clinical trial. *JAMA Intern Med* 2018;178:1051-7.

第5部 全身の症候と内分泌・代謝

5-1 発熱と敗血症の拾い上げ

- 1) Churpek MM, Snyder A, Han X, et al. Quick Sepsis-related Organ Failure Assessment, Systemic Inflammatory Response Syndrome, and Early Warning Scores for detecting clinical deterioration in infected patients outside the intensive care unit. *Am J Respir Crit Care Med* 2017;195:906-11.
- 2) Tokuda Y, Miyasato H, Stein GH, Kishaba T. The degree of chills for risk of bacteremia in acute febrile illness. *Am J Med* 2005;118:1417.
- 3) Mikkelsen ME, Miltiades AN, Gaieski DF, et al. Serum lactate is associated with mortality in severe sepsis independent of organ failure and shock. *Crit Care Med* 2009;37:1670-7.
- 4) Prescott HC, Antonelli M, Alhazzani W, et al. Surviving Sepsis Campaign: international guidelines for management of sepsis and septic shock 2026. *Crit Care Med* 2026;54:725-812.
- 5) 日本版敗血症診療ガイドライン2024特別委員会. 日本版敗血症診療ガイドライン2024. 日集中医誌 2024;31:S1165-313.

- 6) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ロセフィン静注用、ジスロマック、ミノマイシン）。
- 7) Coburn B, Morris AM, Tomlinson G, Detsky AS. Does this adult patient with suspected bacteremia require blood cultures? JAMA 2012;308:502-11.
- 8) Shapiro NI, Wolfe RE, Wright SB, Moore R, Bates DW. Who needs a blood culture? A prospectively derived and validated prediction rule. J Emerg Med 2008;35:255-64.
- 9) High KP, Bradley SF, Gravenstein S, et al. Clinical practice guideline for the evaluation of fever and infection in older adult residents of long-term care facilities: 2008 update by the Infectious Diseases Society of America. Clin Infect Dis 2009;48:149-71.
- 10) Marco CA, Schoenfeld CN, Hansen KN, et al. Fever in geriatric emergency patients: clinical features associated with serious illness. Ann Emerg Med 1995;26:18-24.
- 11) 日本臨床腫瘍学会. 発熱性好中球減少症（FN）診療ガイドライン 改訂第3版. 2024.
- 12) Taplitz RA, Kennedy EB, Bow EJ, et al. Outpatient management of fever and neutropenia in adults treated for malignancy: American Society of Clinical Oncology and Infectious Diseases Society of America clinical practice guideline update. J Clin Oncol 2018;36:1443-53.
- 13) Theilacker C, Ludewig K, Serr A, et al. Overwhelming postsplenectomy infection: a prospective multicenter cohort study. Clin Infect Dis 2016;62:871-8.
- 14) Wilson ME, Weld LH, Boggild A, et al. Fever in returned travelers: results from the GeoSentinel Surveillance Network. Clin Infect Dis 2007;44:1560-8.
- 15) 日本医療研究開発機構「わが国における輸入熱帯病・まれな寄生虫症の診断治療体制の整備」研究班. マラリア 診断・治療・予防の手引き 第6版. 2026.
- 16) 厚生労働省. デング熱に関するQ&A（第4版、令和8年6月29日）。
- 17) 東京都. 東京都感染症マニュアル2018. 2018.
- 18) Matono T, Kutsuna S, Kato Y, et al. Role of classic signs as diagnostic predictors for enteric fever among returned travellers: relative bradycardia and eosinopenia. PLoS One 2017;12:e0179814.
- 19) Chatham-Stephens K, Medalla F, Hughes M, et al. Emergence of extensively drug-resistant Salmonella Typhi infections among travelers to or from Pakistan — United States, 2016–2018. MMWR Morb Mortal Wkly Rep 2019;68:11-3.
- 20) 国立感染症研究所. レプトスピラ症とは（国立感染症研究所のページ、2026年10月に閲覧）。
- 21) 厚生労働行政推進調査事業「一類感染症等の患者発生時に備えた臨床対応及び行政との連携体制の構築のための研究」班. 重症熱性血小板減少症候群（SFTS）診療の手引き 2025年版. 2025.
- 22) Huang DT, Yealy DM, Filbin MR, et al. Procalcitonin-guided use of antibiotics for lower respiratory tract infection. N Engl J Med 2018;379:236-49.
- 23) Usman OA, Usman AA, Ward MA. Comparison of SIRS, qSOFA, and NEWS for the early identification of sepsis in the emergency department. Am J Emerg Med 2019;37:1490-7.

5-2 アナフィラキシーと危険な皮疹

- 1) 日本アレルギー学会Anaphylaxis対策委員会編. アナフィラキシーガイドライン2022. 日本アレルギー学会；2022.
- 2) Campbell RL, Hagan JB, Manivannan V, et al. Evaluation of National Institute of Allergy and Infectious Diseases/Food Allergy and Anaphylaxis Network criteria for the diagnosis of anaphylaxis in emergency department patients. J Allergy Clin Immunol 2012;129:748-52.
- 3) Inomata N, Chin K, Nagashima M, Ikezawa Z. Late-onset anaphylaxis due to poly (γ-glutamic acid) in the soup of commercial cold Chinese noodles in a patient with allergy to fermented soybeans (natto). Allergol Int 2011;60:393-6.
- 4) Inomata N, Miyakawa M, Aihara M. Surfing as a risk factor for sensitization to poly(γ-glutamic acid) in fermented soybeans, natto, allergy. Allergol Int 2018;67:341-6.

- 5) Commins SP, Satinover SM, Hosen J, et al. Delayed anaphylaxis, angioedema, or urticaria after consumption of red meat in patients with IgE antibodies specific for galactose-alpha-1,3-galactose. *J Allergy Clin Immunol* 2009;123:426-33.
- 6) Chinuki Y, Ishiwata K, Yamaji K, Takahashi H, Morita E. Haemaphysalis longicornis tick bites are a possible cause of red meat allergy in Japan. *Allergy* 2016;71:421-5.
- 7) Tejedor-Alonso MA, Farias-Aquino E, Pérez-Fernández E, et al. Relationship between anaphylaxis and use of beta-blockers and angiotensin-converting enzyme inhibitors: a systematic review and meta-analysis of observational studies. *J Allergy Clin Immunol Pract* 2019;7:879-97.
- 8) Simons FE, Gu X, Simons KJ. Epinephrine absorption in adults: intramuscular versus subcutaneous injection. *J Allergy Clin Immunol* 2001;108:871-3.
- 9) ボスミン注1 mgの電子添文（2026年3月改訂 第3版。KEGG MEDICUSで2026年10月に確認）。
- 10) Kim TH, Yoon SH, Hong H, Kang HR, Cho SH, Lee SY. Duration of observation for detecting a biphasic reaction in anaphylaxis: a meta-analysis. *Int Arch Allergy Immunol* 2019;179:31-6.
- 11) Shaker MS, Wallace DV, Golden DBK, et al. Anaphylaxis—a 2020 practice parameter update, systematic review, and Grading of Recommendations, Assessment, Development and Evaluation (GRADE) analysis. *J Allergy Clin Immunol* 2020;145:1082-123.
- 12) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。エピペン注射液、ネフィー点鼻液、ボラミン注、フィラジル皮下注、ペリナートP静注用）。
- 13) 日本小児アレルギー学会. 食物アレルギー診療ガイドライン2021. 協和企画；2021.
- 14) 厚生労働省. ヒスタミンによる食中毒について（厚生労働省のページ、2026年10月に閲覧）。
- 15) 厚生労働省. 重篤副作用疾患別対応マニュアル 血管性浮腫（非ステロイド性抗炎症薬によらないもの）。令和8年2月改定。
- 16) Sinert R, Levy P, Bernstein JA, et al. Randomized trial of icatibant for angiotensin-converting enzyme inhibitor-induced upper airway angioedema. *J Allergy Clin Immunol Pract* 2017;5:1402-9.
- 17) 日本皮膚科学会. 重症多形滲出性紅斑 スティーヴンス・ジョンソン症候群・中毒性表皮壊死症診療ガイドライン補遺2025. 日皮会誌 2025;135:701-14.
- 18) 日本皮膚科学会. 薬剤性過敏症症候群診療ガイドライン2023. 日皮会誌 2024;134:559-80.
- 19) Campbell RL, Bellolio MF, Knutson BD, et al. Epinephrine in anaphylaxis: higher risk of cardiovascular complications and overdose after administration of intravenous bolus epinephrine compared with intramuscular epinephrine. *J Allergy Clin Immunol Pract* 2015;3:76-80.

5-3 血糖と電解質の緊急

- 1) 日本糖尿病学会編. 糖尿病診療ガイドライン2024. 南江堂；2024.
- 2) ビソプロロールフマル酸塩（メインテート錠）の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 3) Namba M, Iwakura T, Nishimura R, et al. The current status of treatment-related severe hypoglycemia in Japanese patients with diabetes mellitus: a report from the committee on a survey of severe hypoglycemia in the Japan Diabetes Society. *Diabetol Int* 2018;9:84-99.
- 4) グリメピリド（アマリール錠）の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 5) Fitzpatrick D, Duncan EA. Improving post-hypoglycaemic patient safety in the prehospital environment: a systematic review. *Emerg Med J* 2009;26:472-8.
- 6) Betten DP, Castle DJ, Hughes MJ, Henney JN. Frequency of return visits to the emergency department in patients discharged following hypoglycemia episodes. *Int J Emerg Med* 2018;11:28.

32 文献

- 7) Umpierrez GE, Davis GM, ElSayed NA, et al. Hyperglycemic crises in adults with diabetes: a consensus report. *Diabetes Care* 2024;47:1257-75.
- 8) Imagawa A, Hanafusa T, Uchigata Y, et al. Fulminant type 1 diabetes: a nationwide survey in Japan. *Diabetes Care* 2003;26:2345-52.
- 9) Umpierrez G, Freire AX. Abdominal pain in patients with hyperglycemic crises. *J Crit Care* 2002;17:63-7.
- 10) メトホルミン（メトグルコ錠）の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 11) Calello DP, Liu KD, Wiegand TJ, et al. Extracorporeal treatment for metformin poisoning: systematic review and recommendations from the Extracorporeal Treatments in Poisoning Workgroup. *Crit Care Med* 2015;43:1716-30.
- 12) Driver BE, Olives TD, Bischof JE, Salmen MR, Miner JR. Discharge glucose is not associated with short-term adverse outcomes in emergency department patients with moderate to severe hyperglycemia. *Ann Emerg Med* 2016;68:697-705.
- 13) UK Kidney Association. Clinical practice guideline: management of hyperkalaemia in adults. Updated July 2026.
- 14) Montague BT, Ouellette JR, Buller GK. Retrospective review of the frequency of ECG changes in hyperkalemia. *Clin J Am Soc Nephrol* 2008;3:324-30.
- 15) Durfey N, Lehnhof B, Bergeson A, et al. Severe hyperkalemia: can the electrocardiogram risk stratify for short-term adverse events? *West J Emerg Med* 2017;18:963-71.
- 16) Spasovski G, Vanholder R, Allolio B, et al. Clinical practice guideline on diagnosis and treatment of hyponatraemia. *Eur J Endocrinol* 2014;170:G1-47.
- 17) Musch W, Decaux G. Treating the syndrome of inappropriate ADH secretion with isotonic saline. *QJM* 1998;91:749-53.
- 18) El-Hajj Fuleihan G, Clines GA, Hu MI, et al. Treatment of hypercalcemia of malignancy in adults: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2023;108:507-28.
- 19) 酸化マグネシウム（マグミット錠）の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 20) Levine M, Nikkanen H, Pallin DJ. The effects of intravenous calcium in patients with digoxin toxicity. *J Emerg Med* 2011;40:41-6.
- 21) グルコン酸カルシウム（カルチコール注射液8.5%）の電子添文（KEGG MEDICUSで2026年10月に確認）。

5-4 甲状腺クリーゼ・粘液水腫性昏迷

- 1) Akamizu T, Satoh T, Isozaki O, et al. Diagnostic criteria, clinical features, and incidence of thyroid storm based on nationwide surveys. *Thyroid* 2012;22:661-79.
- 2) Ono Y, Ono S, Yasunaga H, et al. Clinical characteristics and outcomes of myxedema coma: analysis of a national inpatient database in Japan. *J Epidemiol* 2017;27:117-22.
- 3) 日本甲状腺学会, 日本内分泌学会. 甲状腺クリーゼ診療ガイドライン2017. 2017. (英語版: Satoh T, Isozaki O, Suzuki A, et al. 2016 Guidelines for the management of thyroid storm from The Japan Thyroid Association and Japan Endocrine Society (first edition). *Endocr J* 2016;63:1025-64.)
- 4) Taylor P, Gibb F, Medici M, et al. Management of endocrine emergencies: joint consensus statement for management of myxoedema coma. *Eur Thyroid J* 2026;15:ETJ260044.
- 5) Isozaki O, Satoh T, Wakino S, et al. Treatment and management of thyroid storm: analysis of the nationwide surveys. *Clin Endocrinol (Oxf)* 2016;84:912-8.
- 6) Ono Y, Ono S, Yasunaga H, Matsui H, Fushimi K, Tanaka Y. Factors associated with mortality of thyroid storm: analysis using a national inpatient database in Japan. *Medicine (Baltimore)* 2016;95:e2848.

- 7) Furukawa Y, Tanaka K, Isozaki O, et al. Prospective multicenter registry-based study on thyroid storm: the guidelines for management from Japan are useful. *J Clin Endocrinol Metab* 2024;110:e87-96.
- 8) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 9) 日本甲状腺学会. パセドウ病治療ガイドライン2019. 2019.
- 10) Matsuo Y, Jo T, Watanabe H, et al. Clinical efficacy of beta-1 selective beta-blockers versus propranolol in patients with thyroid storm: a retrospective cohort study. *Crit Care Med* 2024;52:1077-86.
- 11) 日本甲状腺学会 臨床重要課題「粘液水腫性昏睡の診断基準と治療指針」作成委員会. 粘液水腫性昏睡の診断基準（3次案）. 2010年12月.（小西美絵乃，盛田幸司，田中祐司. 粘液水腫性昏睡. 日内会誌 2010;99:769-75 に掲載）
- 12) Garber JR, Cobin RH, Gharib H, et al. Clinical practice guidelines for hypothyroidism in adults: cosponsored by the American Association of Clinical Endocrinologists and the American Thyroid Association. *Endocr Pract* 2012;18:988-1028.
- 13) Yamamoto T, Fukuyama J, Fujiyoshi A. Factors associated with mortality of myxedema coma: report of eight cases and literature survey. *Thyroid* 1999;9:1167-74.

5-5 高齢者の「元気がない・動けない」

- 1) 総務省消防庁. 令和7年版 消防白書. 第2章第5節 救急体制. 2026.
- 2) Karakoumis J, Nickel CH, Kirsch M, et al. Emergency presentations with nonspecific complaints—the burden of morbidity and the spectrum of underlying disease. *Medicine (Baltimore)* 2015;94:e840.
- 3) Lamantia MA, Stewart PW, Platts-Mills TF, et al. Predictive value of initial triage vital signs for critically ill older adults. *West J Emerg Med* 2013;14:453-60.
- 4) Fleming J, Brayne C; Cambridge City over-75s Cohort (CC75C) study collaboration. Inability to get up after falling, subsequent time on floor, and summoning help: prospective cohort study in people over 90. *BMJ* 2008;337:a2227.
- 5) 日本循環器学会, 日本TDM学会. 2015年版 循環器薬の薬物血中濃度モニタリングに関するガイドライン. 2015.
- 6) Norman DC. Fever in the elderly. *Clin Infect Dis* 2000;31:148-51.
- 7) Nicolle LE, Gupta K, Bradley SF, et al. Clinical practice guideline for the management of asymptomatic bacteriuria: 2019 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2019;68:e83-110.
- 8) Canto JG, Shlipak MG, Rogers WJ, et al. Prevalence, clinical characteristics, and mortality among patients with myocardial infarction presenting without chest pain. *JAMA* 2000;283:3223-9.
- 9) Budnitz DS, Lovegrove MC, Shehab N, Richards CL. Emergency hospitalizations for adverse drug events in older Americans. *N Engl J Med* 2011;365:2002-12.
- 10) McGee S, Abernethy WB 3rd, Simel DL. The rational clinical examination. Is this patient hypovolemic? *JAMA* 1999;281:1022-9.
- 11) Uchino S, Bellomo R, Goldsmith D. The meaning of the blood urea nitrogen/creatinine ratio in acute kidney injury. *Clin Kidney J* 2012;5:187-91.
- 12) Han JH, Zimmerman EE, Cutler N, et al. Delirium in older emergency department patients: recognition, risk factors, and psychomotor subtypes. *Acad Emerg Med* 2009;16:193-200.
- 13) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。ジゴシン錠、アリセプト錠、メネシット配合錠）。

- 14) Rockwood K, Theou O. Using the Clinical Frailty Scale in allocating scarce health care resources. *Can Geriatr J* 2020;23:210-5.
- 15) Kaeppli T, Rueegg M, Dreher-Hummel T, et al. Validation of the Clinical Frailty Scale for prediction of thirty-day mortality in the emergency department. *Ann Emerg Med* 2020;76:291-300.
- 16) 高齢者虐待の防止、高齢者の養護者に対する支援等に関する法律（平成17年法律第124号）。第7条。
- 17) 厚生労働省老健局高齢者支援課 認知症・虐待防止対策推進室長通知。市町村や地域包括支援センターにおける高齢者の「セルフ・ネグレクト」及び消費者被害への対応について（平成27年7月10日 老推発0710第2号）。
- 18) Carpenter CR, Shelton E, Fowler S, et al. Risk factors and screening instruments to predict adverse outcomes for undifferentiated older emergency department patients: a systematic review and meta-analysis. *Acad Emerg Med* 2015;22:1-21.
- 19) バラシクロビル（バルトレックス錠）の電子添文（PMDAで2026年10月に確認）。
- 20) Rutschmann OT, Chevalley T, Zumwald C, Luthy C, Vermeulen B, Sarasin FP. Pitfalls in the emergency department triage of frail elderly patients without specific complaints. *Swiss Med Wkly* 2005;135:145-50.
- 21) Sundvall PD, Elm M, Ulleryd P, et al. Interleukin-6 concentrations in the urine and dipstick analyses were related to bacteriuria but not symptoms in the elderly: a cross sectional study of 421 nursing home residents. *BMC Geriatr* 2014;14:88.
- 22) Dasgupta M, Brymer C, Elsayed S. Treatment of asymptomatic UTI in older delirious medical in-patients: a prospective cohort study. *Arch Gerontol Geriatr* 2017;72:127-34.

第6部 中毒・環境・外傷とその他の救急

6-1 急性中毒

- 1) Boehnert MT, Lovejoy FH. Value of the QRS duration versus the serum drug level in predicting seizures and ventricular arrhythmias after an acute overdose of tricyclic antidepressants. *N Engl J Med* 1985;313:474-9.
- 2) Saitman A, Park HD, Fitzgerald RL. False-positive interferences of common urine drug screen immunoassays: a review. *J Anal Toxicol* 2014;38:387-96.
- 3) 日本中毒情報センター。中毒110番・電話サービス（医療機関専用）。<https://www.j-poison-ic.jp/110service/service-guide-medical/>（2026年9月に確認）
- 4) Hoegberg LCG, Gosselin S, Buckley NA, et al. Recommendations from the Clinical Toxicology Recommendations Collaborative on the administration of activated charcoal in acute oral overdose. *Clin Toxicol (Phila)* 2026;64:419-75.
- 5) Chyka PA, Seger D, Krenzelok EP, Vale JA. Position paper: single-dose activated charcoal. *Clin Toxicol (Phila)* 2005;43:61-87.
- 6) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）。
- 7) Benson BE, Hoppu K, Troutman WG, et al. Position paper update: gastric lavage for gastrointestinal decontamination. *Clin Toxicol (Phila)* 2013;51:140-6.
- 8) Thanacoody R, Caravati EM, Troutman B, et al. Position paper update: whole bowel irrigation for gastrointestinal decontamination of overdose patients. *Clin Toxicol (Phila)* 2015;53:5-12.
- 9) Eddleston M, Buckley NA, Eyer P, Dawson AH. Management of acute organophosphorus pesticide poisoning. *Lancet* 2008;371:597-607.
- 10) Proudfoot AT, Krenzelok EP, Vale JA. Position paper on urine alkalinization. *J Toxicol Clin Toxicol* 2004;42:1-26.

- 11) St-Onge M, Anseeuw K, Cantrell FL, et al. Experts consensus recommendations for the management of calcium channel blocker poisoning in adults. *Crit Care Med* 2017;45:e306-15.
- 12) Smilkstein MJ, Knapp GL, Kulig KW, Rumack BH. Efficacy of oral N-acetylcysteine in the treatment of acetaminophen overdose: analysis of the national multicenter study (1976 to 1985). *N Engl J Med* 1988;319:1557-62.
- 13) Lavonas EJ, Akpunonu PD, Arens AM, et al. 2023 American Heart Association focused update on the management of patients with cardiac arrest or life-threatening toxicity due to poisoning. *Circulation* 2023;148:e149-84.
- 14) Isbister GK, Bowe SJ, Dawson A, Whyte IM. Relative toxicity of selective serotonin reuptake inhibitors (SSRIs) in overdose. *J Toxicol Clin Toxicol* 2004;42:277-85.
- 15) Ghannoum M, Yates C, Galvao TF, et al. Extracorporeal treatment for carbamazepine poisoning: systematic review and recommendations from the EXTRIP workgroup. *Clin Toxicol (Phila)* 2014;52:993-1004.
- 16) Ghannoum M, Laliberté M, Nolin TD, et al. Extracorporeal treatment for valproic acid poisoning: systematic review and recommendations from the EXTRIP workgroup. *Clin Toxicol (Phila)* 2015;53:454-65.
- 17) Gawarammana IB, Buckley NA. Medical management of paraquat ingestion. *Br J Clin Pharmacol* 2011;72:745-57.
- 18) Weaver LK, Howe S, Hopkins R, Chan KJ. Carboxyhemoglobin half-life in carbon monoxide-poisoned patients treated with 100% oxygen at atmospheric pressure. *Chest* 2000;117:801-8.
- 19) Weaver LK, Hopkins RO, Chan KJ, et al. Hyperbaric oxygen for acute carbon monoxide poisoning. *N Engl J Med* 2002;347:1057-67.
- 20) Buckley NA, Juurlink DN, Isbister G, et al. Hyperbaric oxygen for carbon monoxide poisoning. *Cochrane Database Syst Rev* 2011;(4):CD002041.
- 21) Choi IS. Delayed neurologic sequelae in carbon monoxide intoxication. *Arch Neurol* 1983;40:433-5.
- 22) Henry CR, Satran D, Lindgren B, Adkinson C, Nicholson CI, Henry TD. Myocardial injury and long-term mortality following moderate to severe carbon monoxide poisoning. *JAMA* 2006;295:398-402.
- 23) Wolf SJ, Maloney GE, Shih RD, Shy BD, Brown MD; American College of Emergency Physicians. Clinical policy: critical issues in the evaluation and management of adult patients presenting to the emergency department with acute carbon monoxide poisoning. *Ann Emerg Med* 2017;69:98-107.e6.
- 24) Rose JJ, Wang L, Xu Q, et al. Carbon monoxide poisoning: pathogenesis, management, and future directions of therapy. *Am J Respir Crit Care Med* 2017;195:596-606.
- 25) Baud FJ, Barriot P, Toffis V, et al. Elevated blood cyanide concentrations in victims of smoke inhalation. *N Engl J Med* 1991;325:1761-6.
- 26) Kamijo Y, Takai M, Fujita Y, Usui K. A retrospective study on the epidemiological and clinical features of emergency patients with large or massive consumption of caffeinated supplements or energy drinks in Japan. *Intern Med* 2018;57:2141-6.
- 27) 日本麻酔科学会. 局所麻酔薬中毒への対応プラクティカルガイド. 2017年6月制定.
- 28) Brett AS, Rothschild N, Gray R, Perry M. Predicting the clinical course in intentional drug overdose: implications for use of the intensive care unit. *Arch Intern Med* 1987;147:133-7.
- 29) Decker BS, Goldfarb DS, Dargan PI, et al. Extracorporeal treatment for lithium poisoning: systematic review and recommendations from the EXTRIP workgroup. *Clin J Am Soc Nephrol* 2015;10:875-87.

- 30) Juurlink DN, Gosselin S, Kielstein JT, et al. Extracorporeal treatment for salicylate poisoning: systematic review and recommendations from the EXTRIP workgroup. *Ann Emerg Med* 2015;66:165-81.
- 31) Carroll R, Metcalfe C, Gunnell D. Hospital presenting self-harm and risk of fatal and non-fatal repetition: systematic review and meta-analysis. *PLoS One* 2014;9:e89944.
- 32) 日本臨床救急医学会. 自殺未遂患者への対応：救急外来（ER）・救急科・救命救急センターのスタッフのための手引き. 2009.
- 33) Kawanishi C, Aruga T, Ishizuka N, et al. Assertive case management versus enhanced usual care for people with mental health problems who had attempted suicide and were admitted to hospital emergency departments in Japan (ACTION-J): a multicentre, randomised controlled trial. *Lancet Psychiatry* 2014;1:193-201.
- 34) Woolf AD, Erdman AR, Nelson LS, et al. Tricyclic antidepressant poisoning: an evidence-based consensus guideline for out-of-hospital management. *Clin Toxicol (Phila)* 2007;45:203-33.
- 35) American College of Medical Toxicology. Guidance document: management priorities in salicylate toxicity. *J Med Toxicol* 2015;11:149-52.
- 36) American Academy of Clinical Toxicology; European Association of Poisons Centres and Clinical Toxicologists. Position paper: cathartics. *J Toxicol Clin Toxicol* 2004;42:243-53.
- 37) Eddleston M, Juszczak E, Buckley NA, et al. Multiple-dose activated charcoal in acute self-poisoning: a randomised controlled trial. *Lancet* 2008;371:579-87.

6-2 アルコール関連の救急

- 1) Jones AW. Evidence-based survey of the elimination rates of ethanol from blood with applications in forensic casework. *Forensic Sci Int* 2010;200:1-20.
- 2) Klein LR, Cole JB, Driver BE, Battista C, Jelinek R, Martel ML. Unsuspected critical illness among emergency department patients presenting for acute alcohol intoxication. *Ann Emerg Med* 2018;71:279-88.
- 3) Easter JS, Haukoos JS, Claud J, et al. Traumatic intracranial injury in intoxicated patients with minor head trauma. *Acad Emerg Med* 2013;20:753-60.
- 4) Pursell RA, Pudek M, Brubacher J, Abu-Laban RB. Derivation and validation of a formula to calculate the contribution of ethanol to the osmolal gap. *Ann Emerg Med* 2001;38:653-9.
- 5) Perez SR, Keijzers G, Steele M, Byrnes J, Scuffham PA. Intravenous 0.9% sodium chloride therapy does not reduce length of stay of alcohol-intoxicated patients in the emergency department: a randomised controlled trial. *Emerg Med Australas* 2013;25:527-34.
- 6) Sullivan JT, Sykora K, Schneiderman J, Naranjo CA, Sellers EM. Assessment of alcohol withdrawal: the revised clinical institute withdrawal assessment for alcohol scale (CIWA-Ar). *Br J Addict* 1989;84:1353-7.
- 7) The ASAM clinical practice guideline on alcohol withdrawal management. *J Addict Med* 2020;14(3S Suppl 1):1-72.
- 8) Mayo-Smith MF. Pharmacological management of alcohol withdrawal: a meta-analysis and evidence-based practice guideline. *JAMA* 1997;278:144-51.
- 9) 日本アルコール・アディクション医学会, 日本アルコール関連問題学会. 新アルコール・薬物使用障害の診断治療ガイドラインに基づいたアルコール依存症の診断治療の手引き【第1版】. 2018 (2019年小改訂).
- 10) Saitz R, Mayo-Smith MF, Roberts MS, Redmond HA, Bernard DR, Calkins DR. Individualized treatment for alcohol withdrawal: a randomized double-blind controlled trial. *JAMA* 1994;272:519-23.
- 11) 各薬剤の電子添文（KEGG MEDICUSで2026年9月に確認）.

- 12) D'Onofrio G, Rathlev NK, Ulrich AS, Fish SS, Freedland ES. Lorazepam for the prevention of recurrent seizures related to alcohol. *N Engl J Med* 1999;340:915-9.
- 13) Maldonado JR, Sher Y, Das S, et al. Prospective validation study of the Prediction of Alcohol Withdrawal Severity Scale (PAWSS) in medically ill inpatients: a new scale for the prediction of complicated alcohol withdrawal syndrome. *Alcohol Alcohol* 2015;50:509-18.
- 14) Wood E, Albarqouni L, Tkachuk S, et al. Will this hospitalized patient develop severe alcohol withdrawal syndrome? The Rational Clinical Examination systematic review. *JAMA* 2018;320:825-33.
- 15) Harper CG, Giles M, Finlay-Jones R. Clinical signs in the Wernicke-Korsakoff complex: a retrospective analysis of 131 cases diagnosed at necropsy. *J Neurol Neurosurg Psychiatry* 1986;49:341-5.
- 16) Galvin R, Bråthen G, Ivashynka A, Hillbom M, Tanasescu R, Leone MA. EFNS guidelines for diagnosis, therapy and prevention of Wernicke encephalopathy. *Eur J Neurol* 2010;17:1408-18.
- 17) National Institute for Health and Care Excellence. Alcohol-use disorders: diagnosis and management of physical complications (CG100). 2010 (updated 2017).
- 18) Wrenn KD, Slovis CM, Minion GE, Rutkowski R. The syndrome of alcoholic ketoacidosis. *Am J Med* 1991;91:119-28.
- 19) Thursz MR, Richardson P, Allison M, et al. Prednisolone or pentoxifylline for alcoholic hepatitis. *N Engl J Med* 2015;372:1619-28.
- 20) Crabb DW, Batailler R, Chalasani NP, et al. Standard definitions and common data elements for clinical trials in patients with alcoholic hepatitis: recommendation from the NIAAA Alcoholic Hepatitis Consortia. *Gastroenterology* 2016;150:785-90.
- 21) 厚生労働省健康局. 標準的な健診・保健指導プログラム（令和6年度版）. 2024.
- 22) 厚生労働省. 健康診断および保健指導におけるアルコール健康障害への早期介入に関するガイドライン. 2024.
- 23) US Preventive Services Task Force; Curry SJ, Krist AH, Owens DK, et al. Screening and behavioral counseling interventions to reduce unhealthy alcohol use in adolescents and adults: US Preventive Services Task Force recommendation statement. *JAMA* 2018;320:1899-909.
- 24) Day E, Bentham PW, Callaghan R, Kuruvilla T, George S. Thiamine for prevention and treatment of Wernicke-Korsakoff syndrome in people who abuse alcohol. *Cochrane Database Syst Rev* 2013;(7):CD004033.
- 25) Dingwall KM, Delima JF, Binks P, Batey R, Bowden SC. What is the optimum thiamine dose to treat or prevent Wernicke's encephalopathy or Wernicke-Korsakoff syndrome? Results of a randomized controlled trial. *Alcohol Clin Exp Res* 2022;46:1133-47.

6-3 興奮・自殺企図

- 1) Richmond JS, Berlin JS, Fishkind AB, et al. Verbal de-escalation of the agitated patient: consensus statement of the American Association for Emergency Psychiatry Project BETA De-escalation Workgroup. *West J Emerg Med* 2012;13:17-25.
- 2) 日本精神科救急学会監修. 精神科救急医療ガイドライン2022年版. 2022.
- 3) Nordstrom K, Zun LS, Wilson MP, et al. Medical evaluation and triage of the agitated patient: consensus statement of the American Association for Emergency Psychiatry Project BETA Medical Evaluation Workgroup. *West J Emerg Med* 2012;13:3-10.
- 4) Olshaker JS, Browne B, Jerrard DA, Prendergast H, Stair TO. Medical clearance and screening of psychiatric patients in the emergency department. *Acad Emerg Med* 1997;4:124-8.

- 5) Rogers JP, Oldham MA, Fricchione G, et al. Evidence-based consensus guidelines for the management of catatonia: recommendations from the British Association for Psychopharmacology. *J Psychopharmacol* 2023;37:327-69.
- 6) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 7) Wilson MP, Pepper D, Currier GW, et al. The psychopharmacology of agitation: consensus statement of the American Association for Emergency Psychiatry Project BETA Psychopharmacology Workgroup. *West J Emerg Med* 2012;13:26-34.
- 8) Klein LR, Driver BE, Miner JR, et al. Intramuscular midazolam, olanzapine, ziprasidone, or haloperidol for treating acute agitation in the emergency department. *Ann Emerg Med* 2018;72:374-85.
- 9) Huf G, Coutinho ES, Adams CE; TREC Collaborative Group. Rapid tranquillisation in psychiatric emergency settings in Brazil: pragmatic randomised controlled trial of intramuscular haloperidol versus intramuscular haloperidol plus promethazine. *BMJ* 2007;335:869.
- 10) Ishida T, Katagiri T, Uchida H, et al. Incidence of deep vein thrombosis in restrained psychiatric patients. *Psychosomatics* 2014;55:69-75.
- 11) 日本臨床救急医学会. 自殺未遂患者への対応：救急外来（ER）・救急科・救命救急センターのスタッフのための手引き. 2009.
- 12) Runeson B, Tidemalm D, Dahlin M, Lichtenstein P, Långström N. Method of attempted suicide as predictor of subsequent successful suicide: national long term cohort study. *BMJ* 2010;341:c3222.
- 13) Quinlivan L, Cooper J, Meehan D, et al. Predictive accuracy of risk scales following self-harm: multicentre, prospective cohort study. *Br J Psychiatry* 2017;210:429-36.
- 14) Large M, Kaneson M, Myles N, Myles H, Gunaratne P, Ryan C. Meta-analysis of longitudinal cohort studies of suicide risk assessment among psychiatric patients: heterogeneity in results and lack of improvement over time. *PLoS One* 2016;11:e0156322.
- 15) National Institute for Health and Care Excellence. Self-harm: assessment, management and preventing recurrence (NICE guideline NG225). 2022.
- 16) Stanley B, Brown GK, Brenner LA, et al. Comparison of the Safety Planning Intervention with follow-up vs usual care of suicidal patients treated in the emergency department. *JAMA Psychiatry* 2018;75:894-900.
- 17) Inagaki M, Kawashima Y, Kawanishi C, et al. Interventions to prevent repeat suicidal behavior in patients admitted to an emergency department for a suicide attempt: a meta-analysis. *J Affect Disord* 2015;175:66-78.
- 18) 精神保健及び精神障害者福祉に関する法律（昭和25年法律第123号）第20条、第21条、第23条、第29条、第29条の2、第33条、第33条の6.
- 19) 警察官職務執行法（昭和23年法律第136号）第3条.

6-4 熱中症・偶発性低体温症

- 1) 日本救急医学会 熱中症および低体温症に関する委員会. 熱中症診療ガイドライン2024. 2024.
- 2) Lott C, Karageorgos V, Abelairas-Gomez C, et al. European Resuscitation Council Guidelines 2025: special circumstances in resuscitation. *Resuscitation* 2025;215 Suppl 1:110753.
- 3) Takauji S, Hifumi T, Saijo Y, et al. Accidental hypothermia: characteristics, outcomes, and prognostic factors—a nationwide observational study in Japan (Hypothermia study 2018 and 2019). *Acute Med Surg* 2021;8:e694.
- 4) Dow J, Giesbrecht GG, Danzl DF, et al. Wilderness Medical Society clinical practice guidelines for the out-of-hospital evaluation and treatment of accidental hypothermia: 2019 update. *Wilderness Environ Med* 2019;30(4S):S47-69.

- 5) Hayashida K, Kondo Y, Hifumi T, et al. A novel early risk assessment tool for detecting clinical outcomes in patients with heat-related illness (J-ERATO score): development and validation in independent cohorts in Japan. *PLoS One* 2018;13(5):e0197032.
- 6) Shimazaki J, Hifumi T, Shimizu K, et al. Clinical characteristics, prognostic factors, and outcomes of heat-related illness (Heatstroke Study 2017-2018). *Acute Med Surg* 2020;7(1):e516.
- 7) Paal P, Pasquier M, Darocha T, et al. Accidental hypothermia: 2021 update. *Int J Environ Res Public Health* 2022;19(1):501.
- 8) Watanabe M, Matsuyama T, Morita S, et al. Impact of rewarming rate on the mortality of patients with accidental hypothermia: analysis of data from the J-Point registry. *Scand J Trauma Resusc Emerg Med* 2019;27:105.
- 9) Dezfulian C, McCallin TE, Bierens J, et al. 2024 American Heart Association and American Academy of Pediatrics focused update on special circumstances: resuscitation following drowning. *Circulation* 2024;150:e501-16.
- 10) Watson RS, Cummings P, Quan L, Bratton S, Weiss NS. Cervical spine injuries among submersion victims. *J Trauma* 2001;51:658-62.
- 11) Schmidt AC, Sempstrott JR, Hawkins SC, Arastu AS, Cushing TA, Auerbach PS. Wilderness Medical Society clinical practice guidelines for the treatment and prevention of drowning: 2019 update. *Wilderness Environ Med* 2019;30(4S):S70-86.
- 12) Tenda ED, Henrina J, Setiadharm A, et al. Systemic corticosteroid as an adjunct for acute respiratory distress syndrome in non-fatal fresh water drowning: an evidence-based case report. *Acta Med Indones* 2024;56:253-9.
- 13) Causey AL, Tilelli JA, Swanson ME. Predicting discharge in uncomplicated near-drowning. *Am J Emerg Med* 2000;18:9-11.

6-5 軽い外傷と創傷

- 1) National Institute for Health and Care Excellence. Head injury: assessment and early management (NG232). 2023.
- 2) Foks KA, van den Brand CL, Lingsma HF, et al. External validation of computed tomography decision rules for minor head injury: prospective, multicentre cohort study in the Netherlands. *BMJ* 2018;362:k3527.
- 3) Hoffman JR, Mower WR, Wolfson AB, Todd KH, Zucker MI. Validity of a set of clinical criteria to rule out injury to the cervical spine in patients with blunt trauma. *N Engl J Med* 2000;343:94-9.
- 4) Stiell IG, Clement CM, McKnight RD, et al. The Canadian C-spine rule versus the NEXUS low-risk criteria in patients with trauma. *N Engl J Med* 2003;349:2510-8.
- 5) Centers for Disease Control and Prevention. HEADS UP: Returning to sports. (2025年9月更新。2026年10月に確認)
- 6) Fernandez R, Green HL, Griffiths R, Atkinson RA, Ellwood LJ. Water for wound cleansing. *Cochrane Database Syst Rev* 2022;9:CD003861.
- 7) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認。キシロカイン注射液、キシロカイン注射液エピレナミン含有、沈降破傷風トキソイド、テタノブリンIH静注、オーグメンチン配合錠、サワシリンカプセル、ラビピュール筋注用、乾燥まむしウマ抗毒素）。
- 8) Perelman VS, Francis GJ, Rutledge T, Foote J, Martino F, Dranitsaris G. Sterile versus nonsterile gloves for repair of uncomplicated lacerations in the emergency department: a randomized controlled trial. *Ann Emerg Med* 2004;43:362-70.
- 9) Farion K, Osmond MH, Hartling L, et al. Tissue adhesives for traumatic lacerations in children and adults. *Cochrane Database Syst Rev* 2002;(3):CD003326.

- 10) Quinn JV, Polevoi SK, Kohn MA. Traumatic lacerations: what are the risks for infection and has the 'golden period' of laceration care disappeared? *Emerg Med J* 2014;31:96-100.
- 11) Cummings P, Del Beccaro MA. Antibiotics to prevent infection of simple wounds: a meta-analysis of randomized studies. *Am J Emerg Med* 1995;13:396-400.
- 12) Seaberg DC, Angelos WJ, Paris PM. Treatment of subungual hematomas with nail trephination: a prospective study. *Am J Emerg Med* 1991;9:209-10.
- 13) Caputo ND, Raja A, Shields C, Menke N. Re-evaluating the diagnostic accuracy of the tongue blade test: still useful as a screening tool for mandibular fractures? *J Emerg Med* 2013;45:8-12.
- 14) Fouad AF, Abbott PV, Tsilingaridis G, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 2. Avulsion of permanent teeth. *Dent Traumatol* 2020;36:331-42.
- 15) Heal C, Buettner P, Raasch B, et al. Can sutures get wet? Prospective randomised controlled trial of wound management in general practice. *BMJ* 2006;332:1053-6.
- 16) 国立感染症研究所. 破傷風 2008年末現在. 病原微生物検出情報 (IASR) 2009;30:65-6.
- 17) 吉野雄一郎, 天野正宏, 飯野志郎, ほか. 創傷・褥瘡・熱傷ガイドライン (2023) —6 熱傷診療ガイドライン (第3版). 日皮会誌 2024;134:509-57.
- 18) Liang JL, Tiwari T, Moro P, et al. Prevention of pertussis, tetanus, and diphtheria with vaccines in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep* 2018;67(2):1-44.
- 19) Stevens DL, Bisno AL, Chambers HF, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis* 2014;59:e10-52.
- 20) Medeiros I, Saconato H. Antibiotic prophylaxis for mammalian bites. *Cochrane Database Syst Rev* 2001;(2):CD001738.
- 21) Talan DA, Citron DM, Abrahamian FM, Moran GJ, Goldstein EJ. Bacteriologic analysis of infected dog and cat bites. *N Engl J Med* 1999;340:85-92.
- 22) 厚生労働省. 狂犬病に関するQ&Aについて. (2026年10月に確認)
- 23) Hifumi T, Sakai A, Kondo Y, et al. Venomous snake bites: clinical diagnosis and treatment. *J Intensive Care* 2015;3:16.
- 24) 厚生労働省. ダニ媒介感染症. (2026年10月に確認)
- 25) Djärv T, Douma M, Palmieri T, et al. Duration of cooling with water for thermal burns as a first aid intervention: a systematic review. *Burns* 2022;48:251-62.
- 26) 日本熱傷学会. 熱傷診療ガイドライン [改訂第3版]. 熱傷 2021;47(Suppl):S1-108.
- 27) Searle J, Slagman A, Maaß W, Möckel M. Cardiac monitoring in patients with electrical injuries: an analysis of 268 patients at the Charité Hospital. *Dtsch Arztebl Int* 2013;110:847-53.
- 28) Bachmann LM, Kolb E, Koller MT, Steurer J, ter Riet G. Accuracy of Ottawa ankle rules to exclude fractures of the ankle and mid-foot: systematic review. *BMJ* 2003;326:417.
- 29) Bachmann LM, Haberzeth S, Steurer J, ter Riet G. The accuracy of the Ottawa knee rule to rule out knee fractures: a systematic review. *Ann Intern Med* 2004;140:121-4.
- 30) Mallee WH, Henny EP, van Dijk CN, Kamminga SP, van Enst WA, Kloen P. Clinical diagnostic evaluation for scaphoid fractures: a systematic review and meta-analysis. *J Hand Surg Am* 2014;39:1683-91.e2.
- 31) Bulger EM, Arneson MA, Mock CN, Jurkovich GJ. Rib fractures in the elderly. *J Trauma* 2000;48:1040-6.
- 32) Maffulli N. The clinical diagnosis of subcutaneous tear of the Achilles tendon: a prospective study in 174 patients. *Am J Sports Med* 1998;26:266-70.

- 33) British Orthopaedic Association. BOA Standard: Diagnosis and management of compartment syndrome of the extremities. 2014 (2025年7月改訂) .
- 34) Donaldson J, Haddad B, Khan WS. The pathophysiology, diagnosis and current management of acute compartment syndrome. *Open Orthop J* 2014;8:185-93.
- 35) 配偶者からの暴力の防止及び被害者の保護等に関する法律（平成13年法律第31号）．第6条、第28条の2.
- 36) 高齢者虐待の防止、高齢者の養護者に対する支援等に関する法律（平成17年法律第124号）．第7条.
- 37) 障害者虐待の防止、障害者の養護者に対する支援等に関する法律（平成23年法律第79号）．第7条.
- 38) 児童虐待の防止等に関する法律（平成12年法律第82号）．第6条.
- 39) 内閣府男女共同参画局．DV相談について．（2026年10月に確認）
- 40) Lalonde D, Bell M, Benoit P, Sparkes G, Denkler K, Chang P. A multicenter prospective study of 3,110 consecutive cases of elective epinephrine use in the fingers and hand: the Dalhousie Project clinical phase. *J Hand Surg Am* 2005;30:1061-7.
- 41) Ilicki J. Safety of epinephrine in digital nerve blocks: a literature review. *J Emerg Med* 2015;49:799-809.
- 42) Prabhakar H, Rath S, Kalaivani M, Bhandari N. Adrenaline with lidocaine for digital nerve blocks. *Cochrane Database Syst Rev* 2015;(3):CD010645.
- 43) Paschos NK, Makris EA, Gantsos A, Georgoulis AD. Primary closure versus non-closure of dog bite wounds: a randomised controlled trial. *Injury* 2014;45:237-40.

6-6 咽頭・鼻・眼・耳の救急

- 1) Sideris A, Holmes TR, Cumming B, Havas T. A systematic review and meta-analysis of predictors of airway intervention in adult epiglottitis. *Laryngoscope* 2020;130:465-73.
- 2) Valerio L, Zane F, Sacco C, et al. Patients with Lemierre syndrome have a high risk of new thromboembolic complications, clinical sequelae and death: an analysis of 712 cases. *J Intern Med* 2021;289:325-39.
- 3) Loureiro RM, Sumi DV, da Costa VLT, Gomes RLE, Soares CR. Clinical, CT, and MRI features of acute calcific tendinitis of the longus colli: a case series with novel imaging findings. *Dentomaxillofac Radiol* 2025;54:420-6.
- 4) Booth AWG, Pungsornruk K, Llewellyn S, Sturgess D, Vidhani K. Airway management of adult epiglottitis: a systematic review and meta-analysis. *BJA Open* 2024;9:100250.
- 5) Chan TV. The patient with sore throat. *Med Clin North Am* 2010;94:923-43.
- 6) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認） .
- 7) Tunkel DE, Anne S, Payne SC, et al. Clinical practice guideline: nosebleed (epistaxis). *Otolaryngol Head Neck Surg* 2020;162(1_suppl):S1-S38.
- 8) Reuben A, Appelboom A, Stevens KN, et al. The use of tranexamic acid to reduce the need for nasal packing in epistaxis (NoPAC): randomized controlled trial. *Ann Emerg Med* 2021;77:631-40.
- 9) Min HJ, Kang H, Choi GJ, Kim KS. Association between hypertension and epistaxis: systematic review and meta-analysis. *Otolaryngol Head Neck Surg* 2017;157:921-7.
- 10) Koskinas I, Terzis T, Georgalas C, et al. Posterior epistaxis management: review of the literature and proposed guidelines of the Hellenic Rhinological-Facial Plastic Surgery Society. *Eur Arch Otorhinolaryngol* 2024;281:1613-27.
- 11) Tran QK, Rehan MA, Haase DJ, Matta A, Pourmand A. Prophylactic antibiotics for anterior nasal packing in emergency department: a systematic review and meta-analysis of clinically-significant infections. *Am J Emerg Med* 2020;38:983-9.

- 12) Fang C, Leavitt JA, Hodge DO, Holmes JM, Mohny BG, Chen JJ. Incidence and etiologies of acquired third nerve palsy using a population-based method. *JAMA Ophthalmol* 2017;135:23-8.
- 13) Optic Neuritis Study Group. The clinical profile of optic neuritis: experience of the Optic Neuritis Treatment Trial. *Arch Ophthalmol* 1991;109:1673-8.
- 14) Bourquard F, Foy JP, Hennocq Q, et al. Diagnosis and management of orbital compartment syndrome caused by retrobulbar hematoma following orbital fractures. *J Craniomaxillofac Surg* 2026;54:104445.
- 15) Dua HS, Ting DSJ, Al Saadi A, Said DG. Chemical eye injury: pathophysiology, assessment and management. *Eye (Lond)* 2020;34:2001-19.
- 16) Mac Grory B, Schrag M, Biousse V, et al. Management of central retinal artery occlusion: a scientific statement from the American Heart Association. *Stroke* 2021;52:e282-94.
- 17) Ryan SJ, Jørstad ØK, Skjelland M, et al. A randomized trial of tenecteplase in acute central retinal artery occlusion. *N Engl J Med* 2026;394:442-50.
- 18) 日本緑内障学会緑内障診療ガイドライン改訂委員会. 緑内障診療ガイドライン（第5版）. 日眼会誌 2022;126:85-177.
- 19) Omar M, Ghouse N, Azar A, Atta S, Malak W. Diagnostic test accuracy of CT for open globe injury: a systematic review and meta-analysis of overall impression and individual imaging signs. *Injury* 2026;57:113041.
- 20) Lim CH, Turner A, Lim BX. Patching for corneal abrasion. *Cochrane Database Syst Rev* 2016;7:CD004764.
- 21) Chandrasekhar SS, Tsai Do BS, Schwartz SR, et al. Clinical practice guideline: sudden hearing loss (update). *Otolaryngol Head Neck Surg* 2019;161(1_suppl):S1-S45.
- 22) Sullivan FM, Swan IR, Donnan PT, et al. Early treatment with prednisolone or acyclovir in Bell's palsy. *N Engl J Med* 2007;357:1598-607.
- 23) Sweeney CJ, Gilden DH. Ramsay Hunt syndrome. *J Neurol Neurosurg Psychiatry* 2001;71:149-54.
- 24) Kim JP, Kwon OJ, Shim HS, Kim RB, Kim JH, Woo SH. Analysis of clinical feature and management of fish bone ingestion of upper gastrointestinal tract. *Clin Exp Otorhinolaryngol* 2015;8:261-7.
- 25) Birk M, Bauerfeind P, Deprez PH, et al. Removal of foreign bodies in the upper gastrointestinal tract in adults: European Society of Gastrointestinal Endoscopy (ESGE) clinical guideline. *Endoscopy* 2016;48:489-96.
- 26) Litovitz T, Whitaker N, Clark L, White NC, Marsolek M. Emerging battery-ingestion hazard: clinical implications. *Pediatrics* 2010;125:1168-77.
- 27) Hayward GN, Hay AD, Moore MV, et al. Effect of oral dexamethasone without immediate antibiotics vs placebo on acute sore throat in adults: a randomized clinical trial. *JAMA* 2017;317:1535-43.
- 28) Sadeghirad B, Siemieniuk RAC, Brignardello-Petersen R, et al. Corticosteroids for treatment of sore throat: systematic review and meta-analysis of randomised trials. *BMJ* 2017;358:j3887.
- 29) Aertgeerts B, Agoritsas T, Siemieniuk RAC, et al. Corticosteroids for sore throat: a clinical practice guideline. *BMJ* 2017;358:j4090.
- 30) Schrag M, Youn T, Schindler J, Kirshner H, Greer D. Intravenous fibrinolytic therapy in central retinal artery occlusion: a patient-level meta-analysis. *JAMA Neurol* 2015;72:1148-54.

6-7 妊娠の可能性のある女性

- 1) 日本産科婦人科学会, 日本産婦人科医会編. 産婦人科診療ガイドライン—産科編2023. 日本産科婦人科学会; 2023.

- 2) Crochet JR, Bastian LA, Chireau MV. Does this woman have an ectopic pregnancy?: the rational clinical examination systematic review. JAMA 2013;309:1722-9.
- 3) Moore C, Todd WM, O'Brien E, et al. Free fluid in Morison's pouch on bedside ultrasound predicts need for operative intervention in suspected ectopic pregnancy. Acad Emerg Med 2007;14:755-8.
- 4) 各薬剤の電子添文（KEGG MEDICUSで2026年10月に確認）。
- 5) 日本医学放射線学会. 画像診断ガイドライン（ウェブ版）産婦人科「妊娠中のCT・MRIは胎児に影響を及ぼすか？ 造影剤投与は胎児に影響を及ぼすか？ 造影剤投与後の授乳は可能か？」. 2026年8月公開.
- 6) 日本医学放射線学会 造影剤安全性委員会. 授乳中の女性に対する造影剤投与後の授乳の可否に関する提言. 2019年6月.
- 7) Kave M, Parooie F, Salarzaei M. Pregnancy and appendicitis: a systematic review and meta-analysis on the clinical use of MRI in diagnosis of appendicitis in pregnant women. World J Emerg Surg 2019;14:37.
- 8) Pasternak B, Svanström H, Mølgaard-Nielsen D, et al. Metoclopramide in pregnancy and risk of major congenital malformations and fetal death. JAMA 2013;310:1601-11.
- 9) Soma-Pillay P, Nelson-Piercy C, Tolppanen H, et al. Physiological changes in pregnancy. Cardiovasc J Afr 2016;27:89-94.
- 10) van der Pol LM, Tromeur C, Bistervels IM, et al. Pregnancy-adapted YEARS algorithm for diagnosis of suspected pulmonary embolism. N Engl J Med 2019;380:1139-49.
- 11) Ramoska EA, Sacchetti AD, Nepp M. Reliability of patient history in determining the possibility of pregnancy. Ann Emerg Med 1989;18:48-50.
- 12) Strote J, Chen G. Patient self assessment of pregnancy status in the emergency department. Emerg Med J 2006;23:554-7.
- 13) Minnerop MH, Garra G, Chohan JK, Troxell RM, Singer AJ. Patient history and physician suspicion accurately exclude pregnancy. Am J Emerg Med 2011;29:212-5.
- 14) Stengel CL, Seaberg DC, MacLeod BA. Pregnancy in the emergency department: risk factors and prevalence among all women. Ann Emerg Med 1994;24:697-700.

6-8 帰宅・診療拒否・警察・書類

- 1) Sessums LL, Zembruska H, Jackson JL. Does this patient have medical decision-making capacity? JAMA 2011;306:420-7.
- 2) 個人情報の保護に関する法律（平成15年法律第57号）. 第27条第1項.
- 3) 個人情報保護委員会, 厚生労働省. 医療・介護関係事業者における個人情報の適切な取扱いのためのガイダンス. 2017（2026年4月一部改正）. 同 Q&A（事例集）Q4-3、Q4-24、Q4-25.
- 4) 山縣然太郎（研究代表者）. 身寄りがいない人の入院及び医療に係る意思決定が困難な人への支援に関するガイドライン. 2019年5月（厚生労働省医政局総務課長通知, 医政総発0603第1号, 令和元年6月3日）.
- 5) 民法（明治29年法律第89号）. 第4条、第698条.
- 6) 警察官職務執行法（昭和23年法律第136号）. 第3条.
- 7) 医療法（昭和23年法律第205号）. 第1条の4第2項.
- 8) 医師法（昭和23年法律第201号）. 第19条、第20条.
- 9) 厚生労働省医政局長通知. 応招義務をはじめとした診察治療の求めに対する適切な対応の在り方等について（医政発1225第4号）. 令和元年12月25日.
- 10) 厚生労働省医政局医事課長通知. 身元保証人等がないことのみを理由に医療機関において入院を拒否することについて（医政医発0427第2号）. 平成30年4月27日.
- 11) 刑法（明治40年法律第45号）. 第130条、第222条、第223条、第234条、第249条.

- 12) 刑事訴訟法（昭和23年法律第131号）．第197条第2項、第218条、第221条、第225条、第239条第2項．
- 13) 覚醒剤取締法（昭和26年法律第252号）．
- 14) 最高裁判所第一小法廷決定．平成17年7月19日．平成17年（あ）第202号．刑集59巻6号600頁．
- 15) 麻薬及び向精神薬取締法（昭和28年法律第14号）．第2条、第58条の2．
- 16) 厚生労働省医薬局長通知．大麻取締法及び麻薬及び向精神薬取締法の一部を改正する法律の施行等について（医薬発1212第1号）．令和6年12月12日．
- 17) 道路交通法（昭和35年法律第105号）．第72条．
- 18) 消防庁．救急業務実施基準（昭和39年自消甲教発第6号．令和7年6月27日 消防救第236号で一部改正）．第24条．
- 19) 消防庁次長，厚生労働省医政局長通知．転院搬送における救急車の適正利用の推進について（消防救第34号，医政発0331第48号）．平成28年3月31日．
- 20) 生活保護法（昭和25年法律第144号）．第25条．
- 21) 厚生労働省保険局保険課長ほか通知．自殺未遂による傷病に係る保険給付等について（保保発0521第1号，保国発0521第2号，保高発0521第1号）．平成22年5月21日．
- 22) Ding R, Jung JJ, Kirsch TD, Levy F, McCarthy ML. Uncompleted emergency department care: patients who leave against medical advice. Acad Emerg Med 2007;14:870-6.
- 23) Hoek AE, Anker SCP, van Beeck EF, et al. Patient discharge instructions in the emergency department and their effects on comprehension and recall of discharge instructions: a systematic review and meta-analysis. Ann Emerg Med 2020;75:435-44.

内科大全 救急編

救急外来と急変

2026年10月07日23:17 下書き（入稿用ではない）

本書に記載した薬剤の用量・用法、診断・治療の基準は、発行時点の電子添文とガイドラインに基づく。実際の使用にあたっては、最新の電子添文とガイドラインを確認すること。

本書の内容の無断での複製・転載を禁じる。