

# Galois Cohomology and Kummer Theory

## 1 A Question about Cyclic Field Extensions

Here's a natural question.

### Question 1.1

*Let  $L/K$  be a Galois extension, and suppose that the Galois group  $G = \text{Gal}(L/K) \cong \mathbb{Z}/n\mathbb{Z}$ . Then is  $L = K(\sqrt[n]{a})$  for some  $a \in K$ ?*

The answer, it turns out, is no, and one doesn't have to look far to find a counterexample, just take any Galois extension  $K/\mathbb{Q}$  with Galois group  $\mathbb{Z}/3\mathbb{Z}$  (for example, the splitting field of  $x^3 + x^2 - 2x - 1$ ). The proof isn't too tricky, if  $K = \mathbb{Q}(\sqrt[3]{a})$ , then as a  $\mathbb{Q}$ -vector space  $K = \mathbb{Q} \oplus \mathbb{Q}\sqrt[3]{a} \oplus \mathbb{Q}\sqrt[3]{a}^2 \subseteq \mathbb{R}$ . But the Galois conjugates of  $\sqrt[3]{a}$  are:

$$\sqrt[3]{a}, \zeta \sqrt[3]{a}, \zeta^2 \sqrt[3]{a},$$

where  $\zeta$  is a primitive cube root of 1. In particular, the last two of these are not contained in  $K$ , so that  $K$  is not Galois. (You can also factor  $x^3 - a = (x - \sqrt[3]{a})(x^2 + x\sqrt[3]{a} + \sqrt[3]{a}^2)$  and see that the discriminant of the latter is negative, so that it can't factor any further and thus the minimal polynomial of  $\sqrt[3]{a}$  doesn't split).

Notice that if  $\mathbb{Q}$  had contained  $\zeta$ , then we would have had no trouble observing that the extension  $K$  was Galois. Indeed, an extension  $K$  is Galois precisely when all the conjugates of a primitive element are in  $K$ . Alternatively, one could factor the minimal polynomial  $x^3 - a = (x - \sqrt[3]{a})(x - \zeta \sqrt[3]{a})(x - \zeta^2 \sqrt[3]{a})$  and see it splits.

Let's summarize our observations so far, but in a slightly more general context. Suppose  $L/K$  is a Galois extension with  $\text{Gal}(L/K) \cong \mathbb{Z}/n\mathbb{Z}$ . If we want  $L \cong K(\sqrt[n]{a})$  for some  $a \in K$ , (notice this implicitly assumes  $x^n - a$  is irreducible in  $K[x]$ ), then we need the  $n$ -th roots of unity to be in  $L$ . Indeed, as  $L$  is Galois, the minimal polynomial of  $a$  must split in  $L$ :

$$x^n - a = (x - \sqrt[n]{a})(x - \zeta \sqrt[n]{a}) \cdots (x - \zeta^{n-1} \sqrt[n]{a}).$$

Here  $\zeta$  is now a primitive  $n$ 'th root of 1. Since Galois acts transitively on the roots of this polynomial, this says that  $\zeta \sqrt[n]{a} \in L$  so that  $\zeta = \zeta \sqrt[n]{a} / \sqrt[n]{a} \in L$ .

### Question 1.2

*Can one use that the Galois group is  $\mathbb{Z}/n\mathbb{Z}$  to show that in fact  $\zeta \in K$ ?*

In particular, we've seen that that a positive answer to Question 1.1 has an explicit obstruction: if  $K$  does not contain a primitive  $n$ 'th root of unity, then the answer is no. On its surface, this obstruction seems much stronger than what is necessary. It tells us that if  $K$  does not contain a primitive  $n$ 'th root of unity, then  $K(\sqrt[n]{a})$  isn't even Galois of degree  $n$ . That is, it tells us that the answer to Question 1.1 is **always no**. That is, at a glance, giving  $K$  a primitive root of 1 puts us in the situation where maybe the answer to Question 1.1 is **sometimes**. The remarkable fact is that this is the *only* obstruction. That is, if  $K$  has a primitive root of unity, the answer to Question 1.1 is **always yes!** (With some restrictions on the characteristic of  $K$ ).

**Theorem 1.3**

Let  $L/K$  be a Galois extension and suppose that the Galois group  $G = \text{Gal}(L/K) \cong \mathbb{Z}/n\mathbb{Z}$ , and suppose that the characteristic of  $K$  does not divide  $n$ . If  $K$  contains a primitive  $n$ 'th root of 1, then  $L \cong K(\sqrt[n]{a})$  for some  $a \in K$ .

The goal of this project is to prove this fact. Notice that there is nothing cohomological in nature about this statement. But a remarkably clever proof of this fact can be extracted from the long exact sequence on cohomology for a particular (right) derived functor.

## 2 Group Cohomology

To turn this problem into a cohomological one we use the following definition.

**Definition 2.1.** Let  $G$  be a group. A  $G$  module is an abelian group  $A$  equipped with an action by  $G$  by automorphisms.

**Exercise 2.2**

Show that the following characterizations of the notion of a  $G$ -module are equivalent.

1. A  $G$ -module  $A$  (as in Definition 2.1).
2. An abelian group  $A$  together with a group homomorphism  $G \rightarrow \text{Aut } A$ .
3. A (ring theoretic) module  $A$  over the group algebra  $\mathbb{Z}[G]$ .

The crucial example for this project is the following.

**Example 2.3**

Let  $K$  be a field and  $L/K$  a Galois field extension with Galois group  $G$ . Then the Galois action naturally makes  $K$  and  $L$  into  $G$ -modules with their underlying additive abelian group structure. (What is the action on  $K$ ?). Similarly, the multiplicative groups  $K^\times$  and  $L^\times$  have natural  $G$ -module structures (with their multiplicative abelian group structures).

We can make  $G$ -modules into a category as follows.

**Definition 2.4.** Let  $A$  and  $B$  be  $G$ -modules. A homomorphism  $\varphi : A \rightarrow B$  is called  $G$ -equivariant if for any  $g \in G$  and  $a \in A$  one has:

$$\varphi(g \cdot a) = g \cdot \varphi(a).$$

**Exercise 2.5**

1. Show that the category of  $G$ -modules with  $G$ -equivariant homomorphisms is an abelian category.
2. Show that the category of  $G$ -modules has enough injectives.

We now can define the following functor from  $G$ -modules to abelian groups:

**Definition 2.6.** Let  $A$  be a  $G$ -module. The  $G$ -invariance of  $A$  is  $A^G = \{a \in A : g \cdot a = a \text{ for all } g \in G\}$ .

**Exercise 2.7**

Let  $K$  and  $L$  be as in example . Compute  $L^G, (L^\times)^G, K^G, (K^\times)^G$ .

**Exercise 2.8**

Show that  $A \mapsto A^G$  is a left exact functor from the category of  $G$ -modules to the category of abelian groups.

Due to Exercises 2.5 and 2.8, we may define the right derived functors of invariance, which is the *group cohomology*. The  $i$ 'th right derived functor will be denoted:

$$H^i(G, \bullet).$$

**Exercise 2.9**

Let  $\mathbb{Z}$  be a  $G$ -module equipped with a trivial action, and  $A$  any  $G$ -module. View both as (ring theoretic)  $\mathbb{Z}[G]$  modules.

1. Show  $A^G \cong \text{Hom}_{\mathbb{Z}[G]}(\mathbb{Z}, A)$ .
2. Show  $H^i(G, A) \cong \text{Ext}_{\mathbb{Z}[G]}^i(\mathbb{Z}, A)$