

# Searching for Rigidity in Algebraic Starscapes

Gabriel Dorfsman-Hopkins

ILAS 2023, Madrid: June 16, 2023

Joint work with Shuchang Xu



ST. LAWRENCE UNIVERSITY

# Land Acknowledgement

# Land Acknowledgement

The University of California Berkeley sits on the territory of xučyun, the ancestral and unceded land of the Chochenyo speaking Ohlone people. This land was and continues to be of great importance to the Muwekma Ohlone Tribe.

It started with a picture...

# It started with a picture...



Image Credit: Stephen Brooks and David Moore



Call this an Algebraic Starscape

# Call this an Algebraic Starscape

## Definition

An algebraic starscape is a plot of algebraic integers,

# Call this an Algebraic Starscape

## Definition

An algebraic starscape is a plot of algebraic integers, complex numbers which are roots of a monic polynomial with integer coefficients.

# Call this an Algebraic Starscape

## Definition

An algebraic starscape is a plot of algebraic integers, complex numbers which are roots of a monic polynomial with integer coefficients.

## Example's of Algebraic Integers:

- $i$  is a root of  $x^2 + 1$ .

# Call this an Algebraic Starscape

## Definition

An algebraic starscape is a plot of algebraic integers, complex numbers which are roots of a monic polynomial with integer coefficients.

## Example's of Algebraic Integers:

- $i$  is a root of  $x^2 + 1$ .
- $e^{2\pi i/n}$  is a root of  $x^n - 1$ .


# Call this an Algebraic Starscape

## Definition

An algebraic starscape is a plot of algebraic integers, complex numbers which are roots of a monic polynomial with integer coefficients.

## Example's of Algebraic Integers:


- $i$  is a root of  $x^2 + 1$ .
- $e^{2\pi i/n}$  is a root of  $x^n - 1$ .
- The golden ratio is a root of  $x^2 - x - 1$ .




Harris, Stange, Trettel



Brooks, Moore



DH, Xu



Derbyshire

# Bohemian Eigenvalue Plots are Starscapes!

# Bohemian Eigenvalue Plots are Starscapes!









The eigenvalues of a square integer matrix algebraic numbers

# Bohemian Eigenvalue Plots are Starscapes!

The eigenvalues of a square integer matrix algebraic numbers, because they are roots of the characteristic polynomial, which is a monic integer polynomial.

# Bohemian Eigenvalue Plots are Starscapes!

The eigenvalues of a square integer matrix algebraic numbers, because they are roots of the characteristic polynomial, which is a monic integer polynomial.



Many different looking starscapes arise. What they look like depends on some choices:

Many different looking starscapes arise. What they look like depends on some choices:

- Which polynomials do we sample from?

Many different looking starscapes arise. What they look like depends on some choices:

- Which polynomials do we sample from?
- Which Bohemian Matrices do we sample from?

Many different looking starscapes arise. What they look like depends on some choices:

- Which polynomials do we sample from?
- Which Bohemian Matrices do we sample from?
- What aesthetic choices are we making in how we plot the points?

Many different looking starscapes arise. What they look like depends on some choices:

- Which polynomials do we sample from?
- Which Bohemian Matrices do we sample from?
- What aesthetic choices are we making in how we plot the points?

## Slogan

Aesthetic choices we make when we draw a collection of algebraic numbers can uncover mathematical structures that could otherwise be hidden.

Many different looking starscapes arise. What they look like depends on some choices:


- Which polynomials do we sample from?
- Which Bohemian Matrices do we sample from?
- What aesthetic choices are we making in how we plot the points?


## Slogan


Aesthetic choices we make when we draw a collection of algebraic numbers can uncover mathematical structures that could otherwise be hidden.


To start let's naively plot quadratics in Sagemath.





















































# Observation

# Observation

As you add more roots the space seemed to fill and lose some sharpness and clarity. Can we deal with this in a meaningful way?



# Observation

- Not all the points are the same size.

# Observation

- Not all the points are the same size. This turns out to be key.

# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial.

# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial. (More precisely, inversely to the magnitude of the vector they form.)


# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial. (More precisely, inversely to the magnitude of the vector they form.)




# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial. (More precisely, inversely to the magnitude of the vector they form.)




# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial. (More precisely, inversely to the magnitude of the vector they form.)




# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial. (More precisely, inversely to the magnitude of the vector they form.)



# Observation

- Not all the points are the same size. This turns out to be key.
- Brooks and Moore size points inversely to the size of the coefficients of the generating polynomial. (More precisely, inversely to the magnitude of the vector they form.)



One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$  we can use the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2}.$$

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$  we can use the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2}.$$

Write  $\sqrt{b^2 - 4c} = n\sqrt{\Delta}$  where  $n$  is the largest integer you can pull out of the square root.

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$  we can use the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2}.$$

Write  $\sqrt{b^2 - 4c} = n\sqrt{\Delta}$  where  $n$  is the largest integer you can pull out of the square root. I'm going to call  $\Delta$  the **discriminant**.

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$  we can use the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2}.$$

Write  $\sqrt{b^2 - 4c} = n\sqrt{\Delta}$  where  $n$  is the largest integer you can pull out of the square root. I'm going to call  $\Delta$  the **discriminant**. Then

$$x = \frac{-b \pm n\sqrt{\Delta}}{2}$$

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$  we can use the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2}.$$

Write  $\sqrt{b^2 - 4c} = n\sqrt{\Delta}$  where  $n$  is the largest integer you can pull out of the square root. I'm going to call  $\Delta$  the **discriminant**. Then

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2} = \frac{-b \pm n\sqrt{\Delta}}{2}.$$

One can also size points by a slightly more refined measure of algebraic complexity: the **discriminant**.

### Example (Discriminants of Quadratics)

If we trying to solve for roots of  $x^2 + bx + c$  we can use the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2}.$$

Write  $\sqrt{b^2 - 4c} = n\sqrt{\Delta}$  where  $n$  is the largest integer you can pull out of the square root. I'm going to call  $\Delta$  the **discriminant**. Then


$$x = \frac{-b \pm \sqrt{b^2 - 4c}}{2} = \frac{-b \pm n\sqrt{\Delta}}{2}.$$

$\Delta$  is a measure of the **irrationality** of  $x$ .

Small discriminant

=


Larger point



Less irrational  $\approx$  Small discriminant

=

Larger point



# Other Approaches

# Other Approaches

- One should ask what happens if you don't vary the size of the points.

# Other Approaches

- One should ask what happens if you don't vary the size of the points.
- It turns out there are lots and lots of algebraic integers.

# Other Approaches

- One should ask what happens if you don't vary the size of the points.
- It turns out there are lots and lots of algebraic integers.
- So many, that you would eventually fill the space.

# Other Approaches

- One should ask what happens if you don't vary the size of the points.
- It turns out there are lots and lots of algebraic integers.
- So many, that you would eventually fill the space. That is, algebraic integers are **dense** in the complex numbers, so every pixel would be colored.

# Other Approaches

- One should ask what happens if you don't vary the size of the points.
- It turns out there are lots and lots of algebraic integers.
- So many, that you would eventually fill the space. That is, algebraic integers are **dense** in the complex numbers, so every pixel would be colored.
- Instead, what if you color a pixel by the density of algebraic integers it contains?

# Other Approaches

- One should ask what happens if you don't vary the size of the points.
- It turns out there are lots and lots of algebraic integers.
- So many, that you would eventually fill the space. That is, algebraic integers are **dense** in the complex numbers, so every pixel would be colored.
- Instead, what if you color a pixel by the density of algebraic integers it contains?

More Integers → Brighter

Fewer Integers → Darker




Image by Sam Derbyshire






Image by Sam Derbyshire

Looks a lot like a dragon fractal!



# Density Plots Reveal Analytic Properties

# Density Plots Reveal Analytic Properties

- Fractal-like behavior.

# Density Plots Reveal Analytic Properties

- Fractal-like behavior.
- (Greg Egan) Connections to the Julia-Mandelbrot correspondence.

# Density Plots Reveal Analytic Properties

- Fractal-like behavior.
- (Greg Egan) Connections to the Julia-Mandelbrot correspondence.

On the other hand...

When you plot individual points sized by algebraic invariants (like the discriminant), you seem to get algebraic curves, like circles, parabolas and lines.




Image by Ed Harris, Katherine Stange, and Steve Treitel

# Paradigm Shifts


# Paradigm Shifts

Compare these two perspectives on drawing a collection of algebraic numbers.

# Paradigm Shifts

Compare these two perspectives on drawing a collection of algebraic numbers.


Density coloring revealed analytic properties.




# Paradigm Shifts

Compare these two perspectives on drawing a collection of algebraic numbers.

Density coloring revealed analytic properties.




Plots sizing according to discriminant revealed algebraic properties.




# Paradigm Shifts

Compare these two perspectives on drawing a collection of algebraic numbers.

Density coloring revealed analytic properties.



Plots sizing according to discriminant revealed algebraic properties.



Let's explain the image on the right.

# Rational Geodesics

What exactly are these curves and why do they appear?

# Rational Geodesics

What exactly are these curves and why do they appear?

**Proposition [HST21]**

The set of complex numbers  $z \in \mathbb{C}$  such that  $\{1, z + \bar{z}, z\bar{z}\}$  are linearly dependent over  $\mathbb{Q}$

# Rational Geodesics

What exactly are these curves and why do they appear?

## Proposition [HST21]

The set of complex numbers  $z \in \mathbb{C}$  such that  $\{1, z + \bar{z}, z\bar{z}\}$  are linearly dependent over  $\mathbb{Q}$  form a collection of vertical lines with rational x-coordinates, and circles centered at rational points with rational radii.

# Rational Geodesics

What exactly are these curves and why do they appear?

## Proposition [HST21]

The set of complex numbers  $z \in \mathbb{C}$  such that  $\{1, z + \bar{z}, z\bar{z}\}$  are linearly dependent over  $\mathbb{Q}$  form a collection of vertical lines with rational x-coordinates, and circles centered at rational points with rational radii. Call these curves **rational geodesics**.

# Rational Geodesics

What exactly are these curves and why do they appear?

## Proposition [HST21]

The set of complex numbers  $z \in \mathbb{C}$  such that  $\{1, z + \bar{z}, z\bar{z}\}$  are linearly dependent over  $\mathbb{Q}$  form a collection of vertical lines with rational x-coordinates, and circles centered at rational points with rational radii. Call these curves **rational geodesics**.

## Corollary [HST21]

If  $z$  is a quadratic algebraic integer, then  $z$  lies on a rational geodesic.

# Rational Geodesics

What exactly are these curves and why do they appear?

## Proposition [HST21]

The set of complex numbers  $z \in \mathbb{C}$  such that  $\{1, z + \bar{z}, z\bar{z}\}$  are linearly dependent over  $\mathbb{Q}$  form a collection of vertical lines with rational x-coordinates, and circles centered at rational points with rational radii. Call these curves **rational geodesics**.

## Corollary [HST21]

If  $z$  is a quadratic algebraic integer, then  $z$  lies on a rational geodesic.


## Proof.

$\{1, z + \bar{z}, z\bar{z}\}$  are 3 elements of a 2 dimensional  $\mathbb{Q}$ -vector space  $\mathbb{Q}(z)$ . □


# Paradigm Shifts

We saw two perspectives on drawing a collection of algebraic numbers.

Density coloring revealed analytic properties.




Plots sizing according to discriminant revealed algebraic properties.




# Paradigm Shifts

We saw two perspectives on drawing a collection of algebraic numbers.

Density coloring revealed analytic properties.



Plots sizing according to discriminant revealed algebraic properties.



In each case, the images inspired new research and new theorems!

An idea:

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special?

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns?

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

- Galois groups

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

- Galois groups
- Ramification

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

- Galois groups
- Ramification
- Inertia

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

- Galois groups
- Ramification
- Inertia
- ...

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

- Galois groups
- Ramification
- Inertia
- ...

The field of algebraic number theory essentially studies how to understand and classify these invariants.

# An idea:

**Slogan:** See interesting new patterns by increasing the size of certain special points.

What if, instead points with small discriminants, we used a different invariant to define special? This would emphasize a different set of points. Would there be new patterns? Algebraic integers have all sorts of fascinating invariants.

- Galois groups
- Ramification
- Inertia
- ...

The field of algebraic number theory essentially studies how to understand and classify these invariants. Let's focus on Galois theory.

# Algebraic Symmetries and Galois Theory

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

This is a symmetry of the complex plane, reflecting it vertically over the real axis.

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

This is a symmetry of the complex plane, reflecting it vertically over the real axis.  
But actually it's even better. If  $z, w \in \mathbb{C}$  :

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

This is a symmetry of the complex plane, reflecting it vertically over the real axis.  
But actually it's even better. If  $z, w \in \mathbb{C}$  :

$$\overline{z + w} = \bar{z} + \bar{w}$$

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

This is a symmetry of the complex plane, reflecting it vertically over the real axis.  
But actually it's even better. If  $z, w \in \mathbb{C}$  :

$$\overline{z + w} = \bar{z} + \bar{w} \quad \text{and} \quad \overline{zw} = \bar{z} \cdot \bar{w}.$$

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

This is a symmetry of the complex plane, reflecting it vertically over the real axis.  
But actually it's even better. If  $z, w \in \mathbb{C}$  :

$$\overline{z + w} = \bar{z} + \bar{w} \quad \text{and} \quad \overline{zw} = \bar{z} \cdot \bar{w}.$$

Complex conjugation is an algebraic symmetry, because it respects addition and multiplication.

# Algebraic Symmetries and Galois Theory

Consider complex conjugation:

$$(z = a + bi) \mapsto (\bar{z} = a - bi).$$

This is a symmetry of the complex plane, reflecting it vertically over the real axis.  
But actually it's even better. If  $z, w \in \mathbb{C}$  :

$$\overline{z + w} = \bar{z} + \bar{w} \quad \text{and} \quad \overline{zw} = \bar{z} \cdot \bar{w}.$$

Complex conjugation is an algebraic symmetry, because it respects addition and multiplication. (You might also see this called an automorphism.)


# Algebraic Symmetries and Permutations

# Algebraic Symmetries and Permutations

Let's apply complex conjugation to the roots of  $x^4 + 2x^2 + 2$ .


# Algebraic Symmetries and Permutations

Let's apply complex conjugation to the roots of  $x^4 + 2x^2 + 2$ .




# Algebraic Symmetries and Permutations

Let's apply complex conjugation to the roots of  $x^4 + 2x^2 + 2$ .




# Algebraic Symmetries and Permutations

Let's apply complex conjugation to the roots of  $x^4 + 2x^2 + 2$ .




# Algebraic Symmetries and Permutations

Let's apply complex conjugation to the roots of  $x^4 + 2x^2 + 2$ .



# Algebraic Symmetries and Permutations

Let's apply complex conjugation to the roots of  $x^4 + 2x^2 + 2$ .



**Slogan:** An algebraic symmetry of the roots gives rise to a permutation of the roots.

Does every permutation give rise to an algebraic symmetry?

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

Example

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula).

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\sigma(\zeta^2)$$

$$\sigma(\zeta)^2$$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\sigma(\zeta^2) = \zeta$$

$$\sigma(\zeta)^2$$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\sigma(\zeta^2) = \zeta$$

$$\sigma(\zeta)^2 = (\zeta^2)^2$$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\sigma(\zeta^2) = \zeta$$

$$\sigma(\zeta)^2 = (\zeta^2)^2 = \zeta^4$$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\begin{aligned}\sigma(\zeta^2) &= \zeta \\ &\qquad\qquad\qquad \times \\ \sigma(\zeta)^2 &= (\zeta^2)^2 = \zeta^4\end{aligned}$$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\begin{array}{rcl} \sigma(\zeta^2) & = & \zeta \\ \nparallel & & \bowtie \\ \sigma(\zeta)^2 & = & (\zeta^2)^2 = \zeta^4 \end{array}$$

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\begin{array}{rcl} \sigma(\zeta^2) & = & \zeta \\ \nparallel & & \bowtie \\ \sigma(\zeta)^2 & = & (\zeta^2)^2 = \zeta^4 \end{array}$$

$\sigma$  does not respect multiplication, and is therefore not algebraic.

Does every permutation give rise to an algebraic symmetry?

- (Gallagher 1973) Almost always: yes!
- In general: no!

### Example

The roots of  $x^5 - 1$  are  $\{1, \zeta, \zeta^2, \zeta^3, \zeta^4\}$ , where  $\zeta = e^{\frac{2\pi i}{5}}$ . (By Euler's formula). Consider a permutation  $\sigma$  swapping  $\zeta$  and  $\zeta^2$ .

$$\begin{array}{rcl} \sigma(\zeta^2) & = & \zeta \\ \swarrow & & \searrow \\ \sigma(\zeta)^2 & = & (\zeta^2)^2 = \zeta^4 \end{array}$$

$\sigma$  does not respect multiplication, and is therefore not algebraic. Since not every permutation of the roots of  $x^5 - 1$  is algebraic, we will call the roots **rigid**.

# Galois Theory

# Galois Theory

Translating what we've said to the language of Galois theory:

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .
- Group of all permutations of roots  $\Leftrightarrow$  Symmetric Group  $S_n$ .

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .
- Group of all permutations of roots  $\Leftrightarrow$  Symmetric Group  $S_n$ .
- Our **slogan** says  $G \leq S_n$ .

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .
- Group of all permutations of roots  $\Leftrightarrow$  Symmetric Group  $S_n$ .
- Our **slogan** says  $G \leq S_n$ .
- Gallagher's result says usually  $G = S_n$ .

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .
- Group of all permutations of roots  $\Leftrightarrow$  Symmetric Group  $S_n$ .
- Our **slogan** says  $G \leq S_n$ .
- Gallagher's result says usually  $G = S_n$ .
- Example shows sometimes  $G \subsetneq S_n$ . (This is the **rigid** case).

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .
- Group of all permutations of roots  $\Leftrightarrow$  Symmetric Group  $S_n$ .
- Our **slogan** says  $G \leq S_n$ .
- Gallagher's result says usually  $G = S_n$ .
- Example shows sometimes  $G \subsetneq S_n$ . (This is the **rigid** case).

## Question

Do rigid algebraic integers satisfy interesting geometric relationships?

# Galois Theory

Translating what we've said to the language of Galois theory:

- Group of algebraic symmetries of roots  $\Leftrightarrow$  Galois Group  $G$ .
- Group of all permutations of roots  $\Leftrightarrow$  Symmetric Group  $S_n$ .
- Our **slogan** says  $G \leq S_n$ .
- Gallagher's result says usually  $G = S_n$ .
- Example shows sometimes  $G \subsetneq S_n$ . (This is the **rigid** case).

## Question

Do rigid algebraic integers satisfy interesting geometric relationships? To explore this we introduce an invariant which measures how far  $G$  is from  $S_n$ , called **rigidity**.

# Rigidity

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ .

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ . Recall that  $\#S_n = n!$ .

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ . Recall that  $\#S_n = n!$ .

## Definition

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ . Recall that  $\#S_n = n!$ .

## Definition

Let  $z$  be an algebraic integer of degree  $n$ , and let  $G$  be its Galois group.

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ . Recall that  $\#S_n = n!$ .

## Definition

Let  $z$  be an algebraic integer of degree  $n$ , and let  $G$  be its Galois group. The rigidity of  $z$  is:

$$\text{rig}(z) := \frac{n! - \#G}{n!}.$$

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ . Recall that  $\#S_n = n!$ .

## Definition

Let  $z$  be an algebraic integer of degree  $n$ , and let  $G$  be its Galois group. The **rigidity** of  $z$  is:

$$\text{rig}(z) := \frac{n! - \#G}{n!}.$$

If  $\text{rig}(z) > 0$  then  $z$  is **rigid**.

# Rigidity

Rigidity measures how far  $G$  is from being all of  $S_n$ . Recall that  $\#S_n = n!$ .

## Definition

Let  $z$  be an algebraic integer of degree  $n$ , and let  $G$  be its Galois group. The **rigidity** of  $z$  is:

$$\text{rig}(z) := \frac{n! - \#G}{n!}.$$

If  $\text{rig}(z) > 0$  then  $z$  is **rigid**.


Notice that if  $G = S_n$  then  $\#G = n!$  so that  $\text{rig}(z) = 0$ .

Let's plot algebraic integers again,

Let's plot algebraic integers again, but now increasing the size of rigid points.


Let's plot algebraic integers again, but now increasing the size of rigid points.

Quartics sized by discriminant.




Let's plot algebraic integers again, but now increasing the size of rigid points.

Quartics sized by discriminant.



Quartics sized by rigidity.




Clearly hidden geometry appears...

Clearly hidden geometry appears... but wait a minute, this looks a bit familiar.


Clearly hidden geometry appears... but wait a minute, this looks a bit familiar.

Are these circles and lines...






Clearly hidden geometry appears... but wait a minute, this looks a bit familiar.



Are these circles and lines...





...the same as these ones?











- Quartics (degree 4 polynomials).



- Quartics (degree 4 polynomials).
- Quadratics (degree 2 polynomials).



- Quartics (degree 4 polynomials).
- Quadratics (degree 2 polynomials).
- Uses Galois theory.



- Quartics (degree 4 polynomials).
- Uses Galois theory.
- Quadratics (degree 2 polynomials).
- Ignores Galois theory.

And yet...

And yet...

Theorem (DH, Xu)

# And yet...

Theorem (DH, Xu)

- *If a quartic lies on a rational geodesic, it is rigid*

# And yet...

Theorem (DH, Xu)

- *If a quartic lies on a rational geodesic, it is rigid of rigidity  $\frac{2}{3}$  or  $\frac{5}{6}$ .*

# And yet...

Theorem (DH, Xu)

- If a quartic lies on a rational geodesic, it is rigid of rigidity  $\frac{2}{3}$  or  $\frac{5}{6}$ .
- If a complex quartic  $z \in \mathbb{C} \setminus \mathbb{R}$

# And yet...

## Theorem (DH, Xu)

- *If a quartic lies on a rational geodesic, it is rigid of rigidity  $\frac{2}{3}$  or  $\frac{5}{6}$ .*
- *If a complex quartic  $z \in \mathbb{C} \setminus \mathbb{R}$  has rigidity  $\frac{5}{6}$ ,*

# And yet...

## Theorem (DH, Xu)

- *If a quartic lies on a rational geodesic, it is rigid of rigidity  $\frac{2}{3}$  or  $\frac{5}{6}$ .*
- *If a complex quartic  $z \in \mathbb{C} \setminus \mathbb{R}$  has rigidity  $\frac{5}{6}$ , it lies on a rational geodesic.*

# And yet...

## Theorem (DH, Xu)

- *If a quartic lies on a rational geodesic, it is rigid of rigidity  $\frac{2}{3}$  or  $\frac{5}{6}$ .*
- *If a complex quartic  $z \in \mathbb{C} \setminus \mathbb{R}$  has rigidity  $\frac{5}{6}$ , it lies on a rational geodesic.*

This result explains the hidden patterns in the image of the rigid quartics, and was completely inspired by that image.

# And yet...

## Theorem (DH, Xu)

- *If a quartic lies on a rational geodesic, it is rigid of rigidity  $\frac{2}{3}$  or  $\frac{5}{6}$ .*
- *If a complex quartic  $z \in \mathbb{C} \setminus \mathbb{R}$  has rigidity  $\frac{5}{6}$ , it lies on a rational geodesic.*

This result explains the hidden patterns in the image of the rigid quartics, and was completely inspired by that image. It also exhibits a previously hidden geometric connection between quadratics and rigid quartics.

# Conclusion

## Slogan

Aesthetic choices we make when we draw a collection of algebraic numbers can uncover mathematical structures that could otherwise be hidden.

# Conclusion

## Slogan

Aesthetic choices we make when we draw a collection of algebraic numbers can uncover mathematical structures that could otherwise be hidden.

- Draw with analytic measurements (e.g. density): get analytic results.

# Conclusion

## Slogan

Aesthetic choices we make when we draw a collection of algebraic numbers can uncover mathematical structures that could otherwise be hidden.

- Draw with analytic measurements (e.g. density): get analytic results.
- Draw arithmetic properties: get arithmetic results.

# Conclusion

## Slogan

Aesthetic choices we make when we draw a collection of algebraic numbers can uncover mathematical structures that could otherwise be hidden.

- Draw with analytic measurements (e.g. density): get analytic results.
- Draw arithmetic properties: get arithmetic results.

## Question

What choices can we make when drawing Bohemian starscapes that could reveal interesting patterns in matrix theory?



Thank You!