

1 Étude d'une fonction

Soit les fonctions $f(x) = \exp(-x^2)$ et $g(x) = -x^4 + 2x^2 + 1$ et $h(x) = \frac{x + 1}{x - 2}$

x	$-\infty$	0	$+\infty$
$f'(x)$	$+$	0	$-$
$f(x)$	0	$\nearrow 1 \searrow$	0

x	$-\infty$	-1	0	1	$+\infty$				
$g'(x)$		$+$	0	$-$	0	$+$	0	$-$	
$g(x)$	$-\infty$	$\nearrow$	2	$\searrow$	1	$\nearrow$	2	$\searrow$	$-\infty$

x	$-\infty$	2	$+\infty$
$h'(x)$	$-$	$\parallel$	$-$
$h(x)$	$1 \searrow$	$-\infty \parallel +\infty \searrow$	1

x	$-\infty$	$+\infty$
e^{x^2}	$+$	
$f(x)$	$+$	

x	$-\infty$	$-\sqrt{1+\sqrt{2}}$	$\sqrt{1+\sqrt{2}}$	$+\infty$	
x^4-2x^2-1	$+$		$-$	$+$	
$g(x)$	$-$	0	$+$	0	$-$

x	$-\infty$	-1	2	$+\infty$
$x + 1$	$-$	0	$+$	$+$
$x - 2$	$-$	$-$	0	$+$
$h(x)$	$+$	0	$- \parallel +$	

$f'(x) = -2xe^{-x^2}$

$g'(x) = 4x(1 - x^2)$

$h'(x) = -\frac{3}{(x - 2)^2}$

$f''(x) = 2(2x^2 - 1)e^{-x^2}$

$g''(x) = 4 - 12x^2$

$$h''(x) = \frac{6}{(x-2)^3}$$

$$f(x) = 0 \iff x \in \{\emptyset\}$$

$$g(x) = 0 \iff x \in \left\{-\sqrt{1+\sqrt{2}}, \sqrt{1+\sqrt{2}}\right\}$$

$$h(x) = 0 \iff x \in \{-1\}$$



