

# Expectation Argument

## Lemma

Let  $X$  be a random variable on a discrete space  $S$  such that  $E(X) = \mu$ . Then  $\Pr[X \geq \mu] > 0$  and  $\Pr[X \leq \mu] > 0$ .

# Max cut

## Instance

Undirected graph  $G = \langle V, E \rangle$

## Feasible solutions

Bipartition  $(A, B)$  of  $V$

## Objective function

$|E \cap (A \times B)|$

# Max cut

```
Data: graph  $G = \langle V, E \rangle$   
foreach vertex  $v \in V$  do  
    | Assign  $v$  to A or to B with probability  $1/2$   
return (A, B)
```

# Max cut

```
Data: graph  $G = \langle V, E \rangle$   
foreach vertex  $v \in V$  do  
    | Assign  $v$  to A or to B with probability  $1/2$   
return (A, B)
```

## Lemma 1 (to prove)

The expected value of the number of edges in the cut (A, B) is at least  $m/2$

## Lemma 2 (to prove)

The probability that the cut (A, B) has at least  $m/2$  edges is  $\geq \frac{1}{m/2+1}$

# Proof of Lemma 2

## Lemma 2

$$p = \Pr \left( C(A, B) \geq \frac{m}{2} \right) \geq \frac{1}{m/2 + 1}$$

# Proof of Lemma 2

## Lemma 2

$$p = \Pr \left( C(A, B) \geq \frac{m}{2} \right) \geq \frac{1}{m/2 + 1}$$

$$\frac{m}{2} = \mathbb{E}[C(A, B)] = \sum_{i \leq \frac{m}{2} - 1} i \Pr(C(A, B) = i) + \sum_{i \geq \frac{m}{2}} i \Pr(C(A, B) = i)$$

$$\Rightarrow p \geq \frac{1}{m/2 + 1}$$

# Proof of Lemma 2

## Lemma 2

$$p = \Pr\left(C(A, B) \geq \frac{m}{2}\right) \geq \frac{1}{m/2 + 1}$$

$$\begin{aligned}\frac{m}{2} = \mathbb{E}[C(A, B)] &= \sum_{i \leq \frac{m}{2} - 1} i \Pr(C(A, B) = i) + \sum_{i \geq \frac{m}{2}} i \Pr(C(A, B) = i) \\ &\leq (1 - p) \left(\frac{m}{2} - 1\right) + pm \\ \Rightarrow p &\geq \frac{1}{m/2 + 1}\end{aligned}$$

# Sample and modifying

## Lemma (to prove)

Let  $G = \langle V, E \rangle$  be a undirected graph. Then  $G = \langle V, E \rangle$  has an independent set with at least  $\frac{n^2}{4m}$  vertices.



# Sample and modifying

## Lemma (to prove)

Let  $G = \langle V, E \rangle$  be a undirected graph. Then  $G = \langle V, E \rangle$  has an independent set with at least  $\frac{n^2}{4m}$  vertices.

Hint 1:  $d = \frac{2m}{n}$  is the average degree.

Hint 2: the proof is a probabilistic algorithm

# Sample and modifying

- 1 Sample  $S \leftarrow$  keep each vertex with probability  $\frac{1}{d}$
- 2 For each edge in the sample, remove one of its endpoints.

# Sample and modifying

- 1 Sample  $S \leftarrow$  keep each vertex with probability  $\frac{1}{d}$
- 2 For each edge in the sample, remove one of its endpoints.

Proof: after step 1

$$\mathbb{E}[|S|] = \frac{n}{d}; \quad \mathbb{E}[\text{number of edges in } S] \geq \frac{nd}{2} \cdot \frac{1}{d^2} = \frac{n}{2d}$$

# Sample and modifying

- 1 Sample  $S \leftarrow$  keep each vertex with probability  $\frac{1}{d}$
- 2 For each edge in the sample, remove one of its endpoints.

Proof: after step 1

$$\mathbb{E}[|S|] = \frac{n}{d}; \quad \mathbb{E}[\text{number of edges in } S] \geq \frac{nd}{2} \cdot \frac{1}{d^2} = \frac{n}{2d}$$

Proof: after step 2

For each edge, we remove at most a vertex. Hence

$$\mathbb{E}[\text{number of surviving vertices}] = \frac{n}{d} - \frac{n}{2d} = \frac{n}{2d} = \frac{n^2}{4m}$$

# Min cut

## Instance

Undirected graph  $G = \langle V, E \rangle$

## Feasible solutions

Bipartition  $(A, B)$  of  $V$

## Objective function

$\max |E \cap (A \times B)|$

# Karger's Algorithm

**Data:** Undirected graph  $G = \langle V, E \rangle$

**while**  $|V| > 2$  **do**

    | Pick a random edge  $(v, w)$ ;

    | Merge  $v$  and  $w$ ;

$\alpha, b \leftarrow$  the two vertices of  $G$ ;

**return** (*vertices merged into  $\alpha$ , vertices merged into  $b$* )

# Karger, Stein Algorithm

**Data:** Undirected graph  $G = \langle V, E \rangle$ , parameter  $t \geq 2$

**while**  $|V| > t$  **do**

- Pick a random edge  $(v, w)$ ;
- Merge  $v$  and  $w$ ;

**return** *optimal solution on G*

# License

This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-sa/4.0/>).

