# ST117 Individual DRAFT Written Report - Task C

My WARWICK ID 5645242 (Report Pod 041)

Today's date in the format 2025-04-10

#### Question 1

Setting up data frames before for phase 2 part C

```
#Import the dataframes from phase 1, precipitation for our pod.
df_precipitation_wide <- readRDS("/Users/danielguo/Desktop/University/Year 1/ST117/5645242_041_WR/df_precipitation
#There is no LCODE or RID for precipitation data set, no averaging is needed

#STAGE, VACUUM, and VOLUME are only contained in some of the datasets and may be dropped unless needed.
df_precipitation_wide <- df_precipitation_wide %>% select(-VOLUME)

#filter out the assigned data for our pod: TO2, TO7
df_precipitation_Afiltered <- df_precipitation_wide %>%
filter(SITECODE %in% c("TO2", "TO7")) %>%
mutate(
    SDATE = as.Date(SDATE),
    YEAR = as.numeric(format(SDATE, "%Y"))
)
```

### Question 2

```
# load and reshape data
df_precip_tall <- df_precipitation_Afiltered %>%
 pivot_longer(-c(SDATE, SITECODE, YEAR), names_to = "Variables", values_to = "VALUE") %>%
 drop_na() %>%
 mutate(DAYS = as.numeric(SDATE - min(SDATE)))
#plot year boxplots
ggplot(df_precip_tall, aes(x = factor(YEAR), y = VALUE)) +
 geom_boxplot(outlier.size = 0.3) +
  facet_grid(Variables ~ SITECODE, scales = "free_y", switch = "y", labeller = label_wrap_gen()) +
 labs(title = "Yearly Boxplots of Chemical Concentrations by Site",
       x = "Year", y = "Concentration") +
  theme_minimal(base_size = 9) +
   axis.text.x = element_text(angle = 90, vjust = 0.5, size = 6),
   axis.text.y = element_text(size = 6),
   strip.text = element_text(size = 7),
   plot.title = element_text(hjust = 0.5, size = 14),
   panel.spacing = unit(0.4, "lines")
```

1

## Yearly Boxplots of Chemical Concentrations by Site

```
Concentration

TAI DII GAL 455 GAL PH OSI HAI NEI ROI DOC DNI. ON LCI MIN LK

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995

1995
```

```
# Compute R2 for each sites and chemicals
mdl_summary <- df_precip_tall %>%
  group_by(SITECODE, Variables) %>%
  group_map(~{
    mod \leftarrow lm(VALUE \sim DAYS, data = .x)
    s <- summary(mod)
    data.frame(
      SITECODE = .y$SITECODE,
      Variables = .y$Variables,
      r.squared = s$r.squared,
      p.value = pf(s$fstatistic[1], s$fstatistic[2], s$fstatistic[3], lower.tail = FALSE)
  }) %>%
  bind_rows()
#pick examples for good poor fit
good_fit <- mdl_summary %>% filter(r.squared > 0.7) %>% slice(1)
poor_fit <- mdl_summary %>% filter(r.squared < 0.1) %>% slice(1)
#function to plot fit
plot_fit <- function(data, fit_info, color, title_prefix) {</pre>
  ggplot(data, aes(x = DAYS, y = VALUE)) +
    geom_point(alpha = 0.6) +
    geom_smooth(method = "lm", color = color) +
      title = paste(title_prefix, "Linear Fit:", fit_info$Variables, "at", fit_info$SITECODE),
      x = "Days",
      y = "concentration"
    )
}
#plot good fit if available, if not say no good fit
if (nrow(good_fit) > 0) {
  good_data <- df_precip_tall %>%
```

```
filter(SITECODE == good_fit$SITECODE[1], Variables == good_fit$Variables[1])
print(plot_fit(good_data, good_fit, "blue", "Good"))
} else {
    message("No good fit found R2 > 0.7")
}

## No good fit found R2 > 0.7

# Plot poor fit, if not say no poor fit
if (nrow(poor_fit) > 0) {
    poor_data <- df_precip_tall %>%
        filter(SITECODE == poor_fit$SITECODE[1], Variables == poor_fit$Variables[1])
    print(plot_fit(poor_data, poor_fit, "red", "Poor"))
} else {
    message("No poor fit found R2 < 0.1.")
}</pre>
```

##  $geom_smooth()$  using formula = 'y ~ x'

#### Poor Linear Fit: ALKY at T02

