

# Math 4610 Task Sheet 2

Manuel Santana

September 21, 2020

## 1 First Order Centered Difference

Here we analyse the error of the approximation

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h}.$$

Consider the following consequences of the Taylor Remainder Theorem

$$f(x+h) = f(x) + f'(x)h + \frac{1}{2}f''(x)h^2 + \frac{1}{6}f'''(\zeta_1)h^3 \quad (1)$$

$$f(x-h) = f(x) + f'(x)(-h) + \frac{1}{2}f''(x)h^2 - \frac{1}{6}f'''(\zeta_2)h^3 \quad (2)$$

with  $x \leq \xi_1 \leq x+h$  and  $x-h \leq \xi_2 \leq x$ . Now let our error term  $\psi$  be

$$\psi = f'(x) - \frac{f(x+h) - f(x-h)}{2h}.$$

Substituting in equations 1, 2 we get

$$\psi = f'(x) - \frac{1}{2h} \left( 2f'(x)h + \frac{1}{6}f'''(\zeta_1)h^3 + \frac{1}{6}f'''(\zeta_2)h^3 \right).$$

Which upon further simplification we see that our error term is second order for some constant  $c$

$$\psi = h^2 \left( \frac{1}{6}f'''(\zeta_1) + \frac{1}{6}f'''(\zeta_2) \right) < ch^2$$

## 2 Second Order Centered Difference

We want do analyze the error approximation

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}.$$

$$f(x+h) = f(x) + f'(x)(h) + \frac{1}{2}f''(x)h^2 + \frac{1}{6}f'''(x)h^3 + \frac{1}{24}f''''(\xi_1)h^4 \quad (3)$$

$$f(x-h) = f(x) - f'(x)(h) + \frac{1}{2}f''(x)h^2 - \frac{1}{6}f'''(x)h^3 + \frac{1}{24}f''''(\xi_2)h^4 \quad (4)$$

with  $x \leq \xi_1 \leq x+h$  and  $x-h \leq \xi_2 \leq x$ . Now let our error term  $\varepsilon$  be

$$\varepsilon = f''(x) - \frac{f(x+h) - 2f(x) + f(x-h)}{h}.$$

Substituting in equations 3, 4 we get

$$\varepsilon = f''(x) - \frac{1}{h^2} \left( \frac{1}{2}f''(x)h^2 + \frac{1}{24}f''''(\xi_1)h^4 + \frac{1}{24}f''''(\xi_2)h^4 \right)$$

This shows our error term is

$$\varepsilon = h^2 \left( \frac{1}{24}f''''(\xi_1) + \frac{1}{24}f''''(\xi_2) \right) < ch^2.$$

Which we see is second order for some constant  $c$ .