

CENG 384 - Signals and Systems for Computer Engineers
Spring 2023
Homework 3

Geçit, Emre
e2521581@ceng.metu.edu.tr

Yancı, Baran
xxxxxxx@ceng.metu.edu.tr

May 13, 2023

1.

$$\begin{aligned}\int_{-\infty}^t x(s)ds &= \int_{-\infty}^t \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 s} ds \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} \Big|_{-\infty}^t \right) \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} - a_k \cdot \frac{e^{jk\omega_0(-\infty)}}{jk\omega_0} \right) \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} - a_k \cdot \frac{0}{jk\omega_0} \right) \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} \right)\end{aligned}$$

This equation is in the synthesis equation form where $a_k \frac{1}{jk\omega_0}$ is the Fourier series coefficients of the integrated signal.

Since ω_0 is the frequency of the signal, $\omega_0 = \frac{2\pi}{T}$ where T is the period of the signal.

Substituting ω_0 in the equation above, we prove the integration property of the Fourier series.

2. (a) $x(t)x(t) \leftrightarrow a_k * a_k$ (Multiplication Property)

(b) $\mathcal{E}\{x(t)\} \leftrightarrow b_k$ (Even Property)

$$b_k = \begin{cases} a_k & k \geq 0 \\ a_{-k} & k < 0 \end{cases}$$

(c) $x(t+t_0) + x(t-t_0) \leftrightarrow a_k e^{jk\omega_0 t_0} + a_{-k} e^{-jk\omega_0 t_0}$ (Shifting and Linearity Properties)

3.

$$x(t) = \begin{cases} 2 & x \in (0, 1) \\ 0 & x \in (1, 2) \\ -2 & x \in (2, 3) \\ 0 & x \in (3, 4) \\ \text{Periodic} & x \notin (0, 4) \end{cases}$$

$$\begin{aligned}
a_k &= \frac{1}{T} \int_0^T x(t) e^{-jk w_0 t} dt \\
&= \frac{1}{4} \left(\int_0^1 2e^{-jk w_0 t} dt + \int_1^2 0 dt + \int_2^3 -2e^{-jk w_0 t} dt + \int_3^4 0 dt \right) \\
&= \frac{1}{4} \left(2 \frac{e^{-jk w_0 t}}{-jk w_0} \Big|_0^1 + 2 \frac{e^{-jk w_0 t}}{-jk w_0} \Big|_2^3 \right) \\
&= \frac{1}{4} \left(2 \frac{e^{-jk w_0}}{-jk w_0} - \frac{2}{-jk w_0} + 2 \frac{e^{-3jk w_0}}{-jk w_0} - 2 \frac{e^{-2jk w_0}}{-jk w_0} \right) \\
&= \frac{1}{-2jk w_0} (e^{-jk w_0} - 1 + e^{-3jk w_0} - e^{-2jk w_0})
\end{aligned}$$

Substitute $w_0 = \frac{2\pi}{T} = \frac{2\pi}{4} = \frac{\pi}{2}$

$$\begin{aligned}
a_k &= \frac{1}{-2jk \frac{\pi}{2}} (e^{-jk \frac{\pi}{2}} - 1 + e^{-3jk \frac{\pi}{2}} - e^{-2jk \frac{\pi}{2}}) \\
&= \frac{1}{-jk \pi} (e^{-jk \frac{\pi}{2}} - 1 + e^{-3jk \frac{\pi}{2}} - e^{-jk \pi}) \\
&= \frac{1}{-jk \pi} (\cos(-k \frac{\pi}{2}) + j \sin(-k \frac{\pi}{2}) - 1 + \cos(-3k \frac{\pi}{2}) + j \sin(-3k \frac{\pi}{2}) - \cos(-k \pi) - j \sin(-k \pi)) \\
&= \frac{1}{-jk \pi} (2 \cos(k \frac{\pi}{2}) - 1 - \cos(k \pi)) \\
&= \frac{1}{jk \pi} (1 - 2 \cos(k \frac{\pi}{2}) + \cos(k \pi))
\end{aligned}$$

4. (a)
- (b)
- (c)
- (d)
5. (a)
- (b)
- (c)
- (d)
6. (a)
- (b)
7. (a)
- (b)
- 8.