

CENG 384 - Signals and Systems for Computer Engineers
Spring 2023
Homework 3

Geçit, Emre
e2521581@ceng.metu.edu.tr

Yancı, Baran
e2449015@ceng.metu.edu.tr

May 17, 2023

1.

$$\begin{aligned}\int_{-\infty}^t x(s)ds &= \int_{-\infty}^t \sum_{k=-\infty}^{\infty} a_k e^{jk\omega_0 s} ds \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} \Big|_{-\infty}^t \right) \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} - a_k \cdot \frac{e^{jk\omega_0(-\infty)}}{jk\omega_0} \right) \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} - a_k \cdot \frac{0}{jk\omega_0} \right) \\ &= \sum_{k=-\infty}^{\infty} \left(a_k \cdot \frac{e^{jk\omega_0 t}}{jk\omega_0} \right)\end{aligned}$$

This equation is in the synthesis equation form where $a_k \frac{1}{jk\omega_0}$ is the Fourier series coefficients of the integrated signal.

Since ω_0 is the frequency of the signal, $\omega_0 = \frac{2\pi}{T}$ where T is the period of the signal.

Substituting ω_0 in the equation above, we prove the integration property of the Fourier series.

2. (a) $x(t)x(t) \leftrightarrow a_k * a_k$ (Multiplication Property)

(b) $\mathcal{E}v\{x(t)\} \leftrightarrow b_k$ (Even Property)

$$b_k = \begin{cases} a_k & k \geq 0 \\ a_{-k} & k < 0 \end{cases}$$

(c) $x(t+t_0) + x(t-t_0) \leftrightarrow a_k e^{jk\omega_0 t_0} + a_{-k} e^{-jk\omega_0 t_0}$ (Shifting and Linearity Properties)

3.

$$x(t) = \begin{cases} 2 & x \in (0, 1) \\ 0 & x \in (1, 2) \\ -2 & x \in (2, 3) \\ 0 & x \in (3, 4) \\ \text{Periodic} & x \notin (0, 4) \end{cases}$$

$$\begin{aligned}
a_k &= \frac{1}{T} \int_0^T x(t) e^{-jk\omega_0 t} dt \\
&= \frac{1}{4} \left(\int_0^1 2e^{-jk\omega_0 t} dt + \int_1^2 0 dt + \int_2^3 -2e^{-jk\omega_0 t} dt + \int_3^4 0 dt \right) \\
&= \frac{1}{4} \left(2 \frac{e^{-jk\omega_0 t}}{-jk\omega_0} \Big|_0^1 - 2 \frac{e^{-jk\omega_0 t}}{-jk\omega_0} \Big|_2^3 \right) \\
&= \frac{1}{4} \left(2 \frac{e^{-jk\omega_0}}{-jk\omega_0} - \frac{2}{-jk\omega_0} - 2 \frac{e^{-3jk\omega_0}}{-jk\omega_0} + 2 \frac{e^{-2jk\omega_0}}{-jk\omega_0} \right) \\
&= \frac{1}{-2jk\omega_0} (e^{-jk\omega_0} - 1 - e^{-3jk\omega_0} + e^{-2jk\omega_0})
\end{aligned}$$

Substitute $\omega_0 = \frac{2\pi}{T} = \frac{2\pi}{4} = \frac{\pi}{2}$

$$\begin{aligned}
a_k &= \frac{1}{-2jk\frac{\pi}{2}} (e^{-jk\frac{\pi}{2}} - 1 - e^{-3jk\frac{\pi}{2}} + e^{-2jk\frac{\pi}{2}}) \\
&= \frac{1}{-jk\pi} (e^{-jk\frac{\pi}{2}} - 1 - e^{-3jk\frac{\pi}{2}} + e^{-jk\pi}) \\
&= \frac{1}{-jk\pi} (\cos(-k\frac{\pi}{2}) + j\sin(-k\frac{\pi}{2}) - 1 - \cos(-3k\frac{\pi}{2}) - j\sin(-3k\frac{\pi}{2}) + \cos(-k\pi) + j\sin(-k\pi)) \\
&= \frac{1}{-jk\pi} (-2j\sin(k\frac{\pi}{2}) - 1 + \cos(-k\pi))
\end{aligned}$$

4. (a)

$$\begin{aligned}
x(t) &= 1 + \sin(\omega_0 t) + 2\cos(\omega_0 t) + \cos(2\omega_0 t + \frac{\pi}{4}) \\
&= 1 + \sin(\omega_0 t) + 2\cos(\omega_0 t) + \cos(2\omega_0 t) \cos(\frac{\pi}{4}) - \sin(2\omega_0 t) \sin(\frac{\pi}{4}) \\
&= 1 + \sin(\omega_0 t) + 2\cos(\omega_0 t) + \frac{\sqrt{2}}{2} \cos(2\omega_0 t) - \frac{\sqrt{2}}{2} \sin(2\omega_0 t) \\
&= 1 + \frac{e^{j\omega_0 t} - e^{-j\omega_0 t}}{2j} + 2 \frac{e^{j\omega_0 t} + e^{-j\omega_0 t}}{2} + \frac{\sqrt{2}}{2} \frac{e^{j2\omega_0 t} + e^{-j2\omega_0 t}}{2} - \frac{\sqrt{2}}{2} \frac{e^{j2\omega_0 t} - e^{-j2\omega_0 t}}{2j} \\
&= 1 + \frac{1}{2j} e^{j\omega_0 t} - \frac{1}{2j} e^{-j\omega_0 t} + e^{j\omega_0 t} + e^{-j\omega_0 t} + \frac{\sqrt{2}}{4} e^{j2\omega_0 t} + \frac{\sqrt{2}}{4} e^{-j2\omega_0 t} - \frac{\sqrt{2}}{4j} e^{j2\omega_0 t} + \frac{\sqrt{2}}{4j} e^{-j2\omega_0 t}
\end{aligned}$$


$$\alpha_0 = 1$$

$$\alpha_1 = 1 + \frac{1}{2j}$$

$$\alpha_{-1} = 1 - \frac{1}{2j}$$

$$\alpha_2 = \frac{\sqrt{2}}{4} - \frac{\sqrt{2}}{4j}$$

$$\alpha_{-2} = \frac{\sqrt{2}}{4} + \frac{\sqrt{2}}{4j}$$



(b)

$$\begin{aligned}
 x(t) &= e^{j\omega_0 t} \\
 y(t) &= H(jk\omega_0)e^{jk\omega_0 t} \\
 y'(t) + y(t) &= x(t) \\
 jk\omega_0 H(jk\omega_0)e^{jk\omega_0 t} + H(jk\omega_0)e^{jk\omega_0 t} &= e^{jk\omega_0 t} \\
 H(jk\omega_0) &= \frac{1}{1 + jk\omega_0}
 \end{aligned}$$

(c)

$$\begin{aligned}
 y(t) &= H(jk\omega_0)e^{jk\omega_0 t} = \frac{1}{1 + jk\omega_0}e^{jk\omega_0 t} \\
 b_k &= a_k \frac{1}{1 + jk\omega_0} \\
 b_1 &= \frac{1 + \frac{1}{2j}}{1 + j\omega_0} \\
 b_{-1} &= \frac{1 - \frac{1}{2j}}{1 - j\omega_0} \\
 b_2 &= \frac{\frac{\sqrt{2}}{4} - \frac{\sqrt{2}}{4j}}{1 + 2j\omega_0} \\
 b_{-2} &= \frac{\frac{\sqrt{2}}{4} + \frac{\sqrt{2}}{4j}}{1 - 2j\omega_0}
 \end{aligned}$$

(d)

$$\begin{aligned}
y(t) &= \sum_{k=-\infty}^{\infty} b_k e^{jk\omega_0 t} \\
&= b_1 e^{j\omega_0 t} + b_{-1} e^{-j\omega_0 t} + b_2 e^{j2\omega_0 t} + b_{-2} e^{-j2\omega_0 t} \\
&= \frac{1 + \frac{1}{2j}}{1 + j\omega_0} e^{j\omega_0 t} + \frac{1 - \frac{1}{2j}}{1 - j\omega_0} e^{-j\omega_0 t} + \frac{\frac{\sqrt{2}}{4} - \frac{\sqrt{2}}{4j}}{1 + 2j\omega_0} e^{j2\omega_0 t} + \frac{\frac{\sqrt{2}}{4} + \frac{\sqrt{2}}{4j}}{1 - 2j\omega_0} e^{-j2\omega_0 t}
\end{aligned}$$

5. (a)

$$\begin{aligned}
x[n] &= \sin\left(\frac{\pi}{2}n\right) \\
&= \frac{e^{j\frac{\pi}{2}n} - e^{-j\frac{\pi}{2}n}}{2j} \\
&= \frac{1}{2j} e^{j\frac{\pi}{2}n} - \frac{1}{2j} e^{-j\frac{\pi}{2}n} \\
\alpha_1 &= \frac{1}{2j} \\
\alpha_{-1} &= -\frac{1}{2j}
\end{aligned}$$

(b)

$$\begin{aligned}
y[n] &= 1 + \cos\left(\frac{\pi}{2}n\right) \\
&= 1 + \frac{e^{j\frac{\pi}{2}n} + e^{-j\frac{\pi}{2}n}}{2} \\
&= 1 + \frac{1}{2} e^{j\frac{\pi}{2}n} + \frac{1}{2} e^{-j\frac{\pi}{2}n} \\
\alpha_0 &= 1 \\
\alpha_1 &= \frac{1}{2} \\
\alpha_{-1} &= \frac{1}{2}
\end{aligned}$$

(c)

$$\begin{aligned}
x[n]y[n] &\leftrightarrow \alpha_k * \beta_k \\
&= \sum_{k=0}^{N-1} \alpha_l \beta_{k-l} \\
&= \sum_{k=0}^3 \alpha_l \beta_{k-l} \\
&= \alpha_0 \beta_{k-0} + \alpha_1 \beta_{k-1} + \alpha_2 \beta_{k-2} + \alpha_3 \beta_{k-3} \\
c_k &= \frac{1}{2} \beta_{k-1} + \frac{1}{2} \beta_{k-3} \\
c_1 &= 0 \\
c_2 &= \frac{-1}{2j} \\
c_3 &= 0 \\
c_4 &= \frac{1}{2j}
\end{aligned}$$


(d)


$$\begin{aligned}
c_k &= \frac{1}{N} \sum_{n=0}^{N-1} x[n]y[n]e^{-j\frac{2\pi}{N}kn} \\
&= \frac{1}{4} \sum_{n=0}^3 x[n]y[n]e^{-j\frac{2\pi}{4}kn} \\
&= \frac{1}{4} (x[0]y[0]e^{-j\frac{2\pi}{4}k0} + x[1]y[1]e^{-j\frac{2\pi}{4}k1} + x[2]y[2]e^{-j\frac{2\pi}{4}k2} + x[3]y[3]e^{-j\frac{2\pi}{4}k3}) \\
&= \frac{1}{4} (0 \cdot 2 \cdot e^{-j\frac{2\pi}{4}k0} + 1 \cdot 1 \cdot e^{-j\frac{2\pi}{4}k1} + 0 \cdot 0 \cdot e^{-j\frac{2\pi}{4}k2} + (-1) \cdot 1 \cdot e^{-j\frac{2\pi}{4}k3}) \\
&= \frac{1}{4} (e^{-j\frac{2\pi}{4}k} - e^{-j\frac{2\pi}{4}k3}) \\
c_1 &= 0 \\
c_2 &= \frac{-1}{2j} \\
c_3 &= 0 \\
c_4 &= \frac{1}{2j}
\end{aligned}$$

The results are the same.

6. (a)

$$\begin{aligned}
x[n] &= 1 - \cos\left(\frac{n\pi}{2}\right) \\
&= 1 - \frac{e^{j\frac{n\pi}{2}} + e^{-j\frac{n\pi}{2}}}{2} \\
&= 1 - \frac{1}{2}e^{j\frac{n\pi}{2}} - \frac{1}{2}e^{-j\frac{n\pi}{2}} \\
\alpha_0 &= 1 \\
\alpha_1 &= -\frac{1}{2} \\
\alpha_{-1} &= -\frac{1}{2}
\end{aligned}$$







(b) i.

ii.

$$\begin{aligned}
 y[n] &= 1 + \sin\left(\frac{n\pi}{2}\right) - \cos\left(\frac{n\pi}{2}\right) \\
 &= 1 + \frac{e^{j\frac{n\pi}{2}} - e^{-j\frac{n\pi}{2}}}{2j} - \frac{e^{j\frac{n\pi}{2}} + e^{-j\frac{n\pi}{2}}}{2} \\
 &= 1 + \frac{1}{2j}e^{j\frac{n\pi}{2}} - \frac{1}{2j}e^{-j\frac{n\pi}{2}} - \frac{1}{2}e^{j\frac{n\pi}{2}} - \frac{1}{2}e^{-j\frac{n\pi}{2}} \\
 &= 1 - \frac{1}{2}e^{j\frac{n\pi}{2}} + \frac{1}{2j}e^{j\frac{n\pi}{2}} - \frac{1}{2j}e^{-j\frac{n\pi}{2}} - \frac{1}{2}e^{-j\frac{n\pi}{2}} \\
 \alpha_0 &= 1 \\
 \alpha_1 &= -\frac{1}{2} + \frac{1}{2j} \\
 \alpha_{-1} &= -\frac{1}{2} - \frac{1}{2j}
 \end{aligned}$$



7. (a)

(b)

8. (a) `from numpy import exp, pi`

```
def spectral_coefficients(signal, period, num_coefficients):  
    coefficients = []  
    for k in range(num_coefficients + 1):  
        S = 0  
        for n in range(period):  
            S += signal[n] * exp(-1j * 2 * pi * n * k / period)  
        coefficients.append(S / period)  
    return coefficients
```

(b) `from matplotlib import pyplot`

`from numpy import exp, pi, linspace`

`SAVE_FOLDER = "figures"`

`t = linspace(-0.5, 0.5, 1000)`

```
class SignalFromSpectralCoefficients:  
    def __init__(self, coefficients, period):  
        self.coefficients = coefficients  
        self.period = period  
  
    def __getitem__(self, n):  
        S = 0  
        for k, coefficient in enumerate(self.coefficients):  
            S += coefficient * exp(1j * 2 * pi * n * k / self.period)  
        return S  
  
    def __iter__(self):  
        for n in range(self.period):  
            yield self[n]  
  
    def __len__(self):  
        return self.period  
  
    def plot(self, name):  
        pyplot.plot(t, self, label="Reconstructed Signal")  
        pyplot.legend()  
        pyplot.savefig(SAVE_FOLDER + "/" + name + ".svg", format = "svg")  
        pyplot.clf()
```

(c) `from matplotlib import pyplot`

`from scipy.signal import sawtooth`

`from q8a import spectral_coefficients`

`from q8b import SignalFromSpectralCoefficients, t`

`square_wave = [-10] * 500 + [10] * 500`

`for n in (1, 5, 10, 50, 100):`

```
    pyplot.plot(t, square_wave, label="Square Wave")  
    coefficients = spectral_coefficients(square_wave, len(square_wave), n)  
    reconstructed = SignalFromSpectralCoefficients(coefficients, 1000)  
    reconstructed.plot(f"square_wave_{n}")
```

Figure 1: Approximated Square Wave with 1 Spectral Coefficient

Figure 2: Approximated Square Wave with 5 Spectral Coefficients

Figure 3: Approximated Square Wave with 10 Spectral Coefficients

Figure 4: Approximated Square Wave with 50 Spectral Coefficients

Figure 5: Approximated Square Wave with 100 Spectral Coefficients

```
(d) import numpy
from matplotlib import pyplot
from scipy.signal import sawtooth
from q8a import spectral_coefficients
from q8b import SignalFromSpectralCoefficients, t

sawtooth_wave = sawtooth(2 * numpy.pi * t)
for n in (1, 5, 10, 50, 100):
    pyplot.plot(t, sawtooth_wave, label="Sawtooth Wave")
    coefficients = spectral_coefficients(sawtooth_wave, len(sawtooth_wave), n)
    reconstructed = SignalFromSpectralCoefficients(coefficients, 1000)
    reconstructed.plot(f"sawtooth_wave_{n}")
```

Figure 6: Approximated Sawtooth Wave with 1 Spectral Coefficient

Figure 7: Approximated Sawtooth Wave with 5 Spectral Coefficients

Figure 8: Approximated Sawtooth Wave with 10 Spectral Coefficients

Figure 9: Approximated Sawtooth Wave with 50 Spectral Coefficients

Figure 10: Approximated Sawtooth Wave with 100 Spectral Coefficients

Although, increasing the number of spectral coefficients increases the accuracy of the approximation, there is a scaling difference between the original and the approximated wave. This is because the number of coefficients used is far less than the number of points in the original wave. For an accurate approximation, the number of coefficients should be equal to the number of points in a period of the original wave.