

# ANOVA

```
graph TD; ANOVA --> OneWay[One-Way ANOVA]; ANOVA --> TwoWay[Two way ANOVA];
```

## One-Way ANOVA

- One independent variable
- Only one 'p' value is obtained.

## Two way ANOVA

- Two independent Variables.
- Three different 'p' values are obtained.
- Outcome of factorial Design.

# Why do an ANOVA?

- when there are 3 or more means being compared, statistical significance can be ascertained by conducting one statistical test, ANOVA, or by repeated t-tests.
- Why not conduct repeated t-tests?
- Each statistical test is conducted with a specified chance of making a type I-error—the alpha level.

# Why do an ANOVA?

- **Why not several t-tests?**
- Imagine we have a design with three groups that have to be compared:  
G1, G2, G3
- We will have to run several separate t-tests  
(one to compare G1 with G2, one to compare G1 with G3, and one to compare G2 with G3)
- For every test we use a general  $\alpha$ -level of 0.05

- **$\alpha$  -level=0.05**
- 5% possibility to make Type I error, i.e. rejecting  $H_0$ , when  $H_0$  is actually true.
- Our scope is too reduce the possibilities to have Type I error
- If we were to run 3 separate t-tests to compare G1, G2 and G3, each with a  $\alpha$ -level of 0.05, the overall possibility not to make Type I error would be 0.857 [i.e.  $(0.95)^3$ ]
- Therefore subtracting that from the overall possibility not to make Type 1 error (1=100%)
 
$$1 - 0.857 = 0.14$$
- We have 14% of possibilities to make Type 1 error.
- 14% >> than the usual 5%
- We can't be happy with that!

# Assumptions of one way ANOVA

- The data are randomly sampled and independently chosen from the populations
- The variances of each sample are assumed equal.
- The residuals are normally distributed.

## State the null hypothesis and alpha level

The null hypothesis is that all the groups have equal means.

$$H_0: \mu_1 = \mu_2 = \mu_3$$

The alternative hypothesis is that there is at least one significant difference between the means

Level of significance  $\alpha$  is selected as 0.05

# Computing a one-way ANOVA

Here is the basic one-way ANOVA table

Source	SS	df	MS	F	p	F <sub>crit</sub>
Between						
Within						
Total						

## Sum of squares (SS)

- Sum of squares are nothing but sum of squared deviations around the mean i.e.  $(X - \bar{X})^2$
- To calculate the various sums of squares, we need:
  - i) the  $\Sigma X^2$  (sum of each score squared) in the whole sample
  - ii) the  $\Sigma X$  (sum of all scores) for the whole sample
  - iii) and the Group  $\Sigma X$  (sum of scores separately for each group): represented as  $\Sigma X_g$  in as follows



FRONT ROW	MIDDLE ROW	BACK ROW
8	13	9
8	11	11
6	15	14
9	14	12
10	14	9
13	12	11
9	11	12
11	11	11
8	11	9
11	11	9
10	17	11
9	13	11
112	153	129
1082	1993	1413
9.33	12.75	10.75

$\Sigma X_g =$

$\Sigma X^2 =$

Mean =

= 394

= 4488

	FRONT ROW	MIDDLE ROW	BACK ROW	TOTAL
$\Sigma X_g$	112	153	129	394
$\Sigma X^2$	1082	1993	1413	4488

- $$SS_{\text{total}} = \Sigma X^2 - \frac{(\Sigma X)^2}{N}$$

$$= 4488 - 394^2 / 36$$

$$= 4488 - 4312.11$$

$$= 175.88$$

	FRONT ROW	MIDDLE ROW	BACK ROW	TOTAL
$\Sigma X_g$	112	153	129	394
$\Sigma X^2$	1082	1993	1413	4488

- $$SS_{\text{between}} = \frac{(\Sigma X_g)^2}{n} - \frac{(\Sigma X)^2}{N}$$

$$= \frac{112^2 + 153^2 + 129^2}{12} - (394^2 / 36)$$

$$= 4382.83 - 4312.11$$

$$= \mathbf{70.72}$$

$$SS_{\text{total}} = 175.88$$

$$SS_{\text{between}} = 70.72$$

$$SS_{\text{within}} = SS_{\text{total}} - SS_{\text{between}}$$

$$= 175.88 - 70.72$$

$$= 105.16$$

After filling in the sum of squares, we have ...

Source	SS	df	MS	F	p	F <sub>crit</sub>
Between	70.72					
Within	105.16					
Total	175.88					

# Degrees of Freedom

- The degrees of freedom, noted in are calculated as  $N_i - 1$  for the total( $N_i$  is the total number of observations);
- number of groups minus one for the between groups;
- And for the within error, subtract d.f. for groups from the total degrees of freedom.

So ,

$$\begin{aligned}\text{Total degrees of freedom} &= \text{total number of observations} - 1 \\ &= 36-1 \\ &= \mathbf{35}\end{aligned}$$

$$\begin{aligned}\text{Degrees of freedom between groups} &= \text{Total number of groups} - 1 \\ &= 3-1 \\ &= \mathbf{2}\end{aligned}$$

$$\begin{aligned}\text{Degrees freedom within groups} &= \text{Total d.f} - \text{Between group d.f} \\ &= 35-2 \\ &= \mathbf{33}\end{aligned}$$

- Filling in the degrees of freedom gives this ...

<b>Source</b>	<b>SS</b>	<b>df</b>	<b>MS</b>	<b>F</b>	<b>p</b>	<b>F<sub>crit</sub></b>
Between	70.72	2				
Within	105.16	33				
Total	175.88	35				



# Mean Squares-MS (Variance)

- **Why variance estimates are called Mean Squares?**

A population variance is equal to the sum of the squared deviations about the mean divided by N.

$$\frac{\Sigma (X - \bar{X})^2}{N}$$

So the population variance is really the mean of the squared deviations about the mean, or the mean square(d) deviation about the mean. This is why the term mean square is used in place of variance.

It is obtained by dividing each sum of squares with corresponding degrees of freedom.

So,

<b>MS<sub>between</sub></b>	<b>MS<sub>within</sub></b>
$= 70.7222 / 2$	$= 105.1667 / 33$
$= 35.36$	$= 3.18$

<b>Source</b>	<b>SS</b>	<b>df</b>	<b>MS</b>	<b>F</b>	<b>p</b>	<b>F<sub>crit</sub></b>
Between	70.72	2	35.36			
Within	105.16	33	3.18			
Total	175.88	35				

## $F_{\text{calculated}}$ value


- It is the ratio of  $MS_{\text{between}}$  and  $MS_{\text{within}}$

$$F = MS(B) / MS(W)$$

- So,  $F = \frac{35.3611}{3.1869} = 11.10$

Source	SS	df	MS	F	p	F <sub>crit</sub>
Between	70.72	2	35.36	11.10		
Within	105.16	33	3.18			
Total	175.88	35				

Table F.1  
Critical Values of the F Distribution for  $\alpha = .05$




0		F																		df NUMERATOR																	
df	Denominator	1	2	3	4	5	6	7	8	9	10	12	15	20	25	30	40	60	120	∞																	
3		10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.74	8.70	8.66	8.63	8.62	8.59	8.57	8.55	8.53																	
4		7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.91	5.86	5.80	5.77	5.75	5.72	5.69	5.66	5.63																	
5		6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.68	4.62	4.56	4.52	4.50	4.46	4.43	4.40	4.36																	
6		5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.00	3.94	3.87	3.83	3.81	3.77	3.74	3.70	3.67																	
7		5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.57	3.51	3.44	3.40	3.38	3.34	3.30	3.27	3.23																	
8		5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.28	3.22	3.15	3.11	3.08	3.04	3.01	2.97	2.93																	
9		5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.07	3.01	2.94	2.89	2.86	2.83	2.79	2.75	2.71																	
10		4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.91	2.85	2.77	2.73	2.70	2.66	2.62	2.58	2.54																	
11		4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.79	2.72	2.65	2.60	2.57	2.53	2.49	2.45	2.40																	
12		4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.69	2.62	2.54	2.50	2.47	2.43	2.38	2.34	2.30																	
13		4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.60	2.53	2.46	2.41	2.38	2.34	2.30	2.25	2.21																	
14		4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.53	2.46	2.39	2.34	2.31	2.27	2.22	2.18	2.13																	
15		4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.48	2.40	2.33	2.28	2.25	2.20	2.16	2.11	2.07																	
16		4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	2.49	2.42	2.35	2.28	2.23	2.19	2.15	2.11	2.06	2.01																	
17		4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	2.45	2.38	2.31	2.23	2.18	2.15	2.10	2.06	2.01	1.96																	
18		4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	2.41	2.34	2.27	2.19	2.14	2.11	2.06	2.02	1.97	1.92																	
19		4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	2.38	2.31	2.23	2.16	2.11	2.07	2.03	1.98	1.93	1.88																	
20		4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	2.35	2.28	2.20	2.12	2.07	2.04	1.99	1.95	1.90	1.84																	
21		4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37	2.32	2.25	2.18	2.10	2.04	2.01	1.96	1.92	1.87	1.81																	
22		4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	2.30	2.23	2.15	2.07	2.02	1.98	1.94	1.89	1.84	1.78																	
23		4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	2.27	2.20	2.13	2.05	2.00	1.96	1.91	1.86	1.81	1.76																	
24		4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.18	2.11	2.03	1.97	1.94	1.89	1.84	1.79	1.73																	
25		4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	2.24	2.16	2.09	2.01	1.95	1.92	1.87	1.82	1.77	1.71																	
26		4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	2.22	2.15	2.07	1.99	1.94	1.90	1.85	1.80	1.75	1.69																	
27		4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25	2.20	2.13	2.06	1.97	1.92	1.88	1.84	1.79	1.73	1.67																	
28		4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	2.19	2.12	2.04	1.96	1.91	1.87	1.82	1.77	1.71	1.65																	
29		4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22	2.18	2.10	2.03	1.94	1.90	1.85	1.81	1.75	1.70	1.64																	
30		4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.09	2.01	1.93	1.88	1.84	1.79	1.74	1.68	1.62																	
40		4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.00	1.92	1.84	1.78	1.74	1.69	1.64	1.58	1.51																	
60		4.00	3.15	2.76	2.53	2.37	2.25	2.17	2.10	2.04	1.99	1.92	1.84	1.75	1.69	1.65	1.59	1.53	1.47	1.39																	
120		3.92	3.07	2.68	2.45	2.29	2.17	2.09	2.02	1.96	1.91	1.83	1.75	1.66	1.60	1.55	1.50	1.43	1.35	1.25																	
∞		3.84	3.00	2.60	2.37	2.21	2.10	2.01	1.94	1.88	1.83	1.75	1.67	1.57	1.51	1.46	1.39	1.32	1.22	1.00																	

Putting value of F crit to ANOVA table

<b>Source</b>	<b>SS</b>	<b>df</b>	<b>MS</b>	<b>F</b>	<b>p</b>	<b>F<sub>crit</sub></b>
Between	70.72	2	35.36	11.10		3.32
Within	105.16	33	3.18			
Total	175.88	35				

# *P*-value

- It is the area under the F distribution curve that is to the right of your observed F-statistic.
- This area may be obtained by integral calculus



- If the *P*-value is less than or equal to  $\alpha$ , reject  $H_0$ .
- If the *P*-value is greater than  $\alpha$ , fail to reject  $H_0$ .



# One-Way ANOVA: Interpretation

- Our obtained F of 11.10 is greater than the critical F of 3.32 i.e.  $F_{\text{calculated}} > F_{\text{critical}}$  and p- value is less than 0.05
- We reject the null hypothesis and accept the alternative hypothesis that there is at least a difference between two of the group means.
- If  $F_{\text{cal}} < F_{\text{crit}}$ , We fail to reject the null hypothesis and conclude that there are no significant differences between the group means.

New Microsoft Office Excel Worksheet.xlsx - Microsoft Excel

Home Insert Page Layout Formulas Data Review View

Get External Data Refresh All Edit Links Connections Sort & Filter Filter Sort Clear Reapply Advanced Data Tools Text to Columns Remove Duplicates Data Validation Consolidate What-If Analysis Group Ungroup Subtotal Outline Data Analysis Analysis

E8

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
1	FRONT ROW	MIDDLE ROW	BACK ROW												
2	8	13	9												
3	8	11	11												
4	6	15	14												
5	9	14	12												
6	10	14	9												
7	13	12	11												
8	9	11	12												
9	11	11	11												
10	8	11	9												
11	11	11	9												
12	10	17	11												
13	9	13	11												
14															
15															
16															
17															
18															
19															
20															
21															
22															
23															
24															
25															

Sheet1 Sheet2 Sheet3



New Microsoft Office Excel Worksheet.xlsx - Microsoft Excel

Home Insert Page Layout Formulas Data Review View

Get External Data Refresh All Edit Links Connections Sort & Filter Filter Clear Reapply Advanced Text to Columns Remove Duplicates Data Validation Consolidate What-If Analysis Group Ungroup Subtotal Outline Data Analysis Analysis

E8

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
1	FRONT ROW	MIDDLE ROW	BACK ROW												
2	8	13	9												
3	8	11	11												
4	6	15	14												
5	9	14	12												
6	10	14	9												
7	13	12	11												
8	9	11	12												
9	11	11	11												
10	8	11	9												
11	11	11	9												
12	10	17	11												
13	9	13	11												
14															
15															
16															
17															
18															
19															
20															
21															
22															
23															
24															

Data Analysis

Analysis Tools

- Anova: Single Factor
- Anova: Two-Factor With Replication
- Anova: Two-Factor Without Replication
- Correlation
- Covariance
- Descriptive Statistics
- Exponential Smoothing
- F-Test Two-Sample for Variances
- Fourier Analysis
- Histogram

OK Cancel Help

New Microsoft Office Excel Worksheet.xlsx - Microsoft Excel

Home Insert Page Layout Formulas Data Review View

Get External Data Refresh All Edit Links Connections Sort & Filter Filter Clear Reapply Advanced Text to Columns Remove Duplicates Data Validation Consolidate What-If Analysis Group Ungroup Subtotal Data Analysis Analysis

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O
1	FRONT ROW	MIDDLE ROW	BACK ROW												
2	8	13	9												
3	8	11	11												
4	6	15	14												
5	9	14	12												
6	10	14	9												
7	13	12	11												
8	9	11	12												
9	11	11	11												
10	8	11	9												
11	11	11	9												
12	10	17	11												
13	9	13	11												
14															
15															
16															
17															
18															
19															
20															
21															
22															
23															
24															

Anova: Single Factor

Input

Input Range: \$A\$1:\$C\$13

Grouped By: ☒ Columns ☐ Rows

☒ Labels in first row

Alpha: 0.05


Output options

☒ Output Range: \$I\$11

☐ New Worksheet Ply:


☐ New Workbook

OK Cancel Help



# HOW TO CALCULATE P- VALUE IN EXCEL?





New Microsoft Office Excel Worksheet.xlsx - Microsoft Excel

Home Insert Page Layout Formulas Data Review View

Paste Font Alignment Number Styles Cells Editing

FDIST X 11.09 Deg\_freedom1 3 Deg\_freedom2 36 = 2.66227E-05

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26

Sheet1 Sheet2 Sheet3

### Function Arguments

FDIST

X	11.09	= 11.09
Deg_freedom1	3	= 3
Deg_freedom2	36	= 36

= 2.66227E-05

Returns the F probability distribution (degree of diversity) for two data sets.

**Deg\_freedom2** is the denominator degrees of freedom, a number between 1 and 10^10, excluding 10^10.

Formula result = 2.66227E-05

[Help on this function](#)

OK Cancel

New Microsoft Office Excel Worksheet.xlsx - Microsoft Excel

Home Insert Page Layout Formulas Data Review View

Clipboard Font Alignment Number Styles Cells Editing

Calibri 11 A A

B I U

General

\$ %

Conditional Formatting Format as Table Cell Styles

Insert Delete Format

Σ Sort & Filter Find & Select

A1 fx =FDIST(11.09,3,36)

	A	B	C	D	E	F	G	H	I	J	K	L	M	N
1	2.66227E-05													
2														
3														
4														
5														
6														
7														
8														
9														
10														
11														
12														
13														
14														
15														
16														
17														
18														
19														
20														
21														
22														
23														
24														
25														
26														

Sheet1 Sheet2 Sheet3