

EE510 Linear Algebra for Engineering



Week 1 Session 1

Review:

Logical Inference

Logical Statement P and Q

| P | Q | $\neg P$ | $\neg Q$ | $P \wedge Q$ | $P \vee Q$ | $P \implies Q$ | $P \iff Q$ |
|-----|-----|----------|----------|--------------|------------|----------------|------------|
| 1 | 1 | 0 | 0 | 1 | 1 | 1 | 1 |
| 0 | 1 | 1 | 0 | 0 | 1 | 1 | 0 |
| 1 | 0 | 0 | 1 | 0 | 1 | 0 | 0 |
| 0 | 0 | 1 | 1 | 0 | 0 | 1 | 1 |

\wedge is AND

\vee is OR

\implies If then

\iff If and only if

Conditional: $P \implies Q$

Contrastive: $\neg P \implies \neg Q$

Converse: $Q \implies P$

Predicate: Px means x is P

Quantifier: $\forall x$ (universal) means "for all x "

$\exists x$ (existential) means "for some x "

$\forall x: Px$ means "Everything is P "

$Px_1 \text{ AND } Px_2 \text{ AND } Px_3 \text{ AND}$

$\exists x: Px$ means "Something is P "

$Px_1 \text{ OR } Px_2 \text{ OR } Px_3 \text{ OR}$

Rules of Inference:

- Modus Ponens: Affirming the antecedent

Premise 1: $P \implies Q$

Premise 2: P

Conclusion: Q

- Modus Tollens: Denying the consequent

Premise 1: $P \implies Q$

Premise 2: $\neg Q$

Conclusion: $\neg P$

- Mathematical Induction

Goal: Proof that $P_n \forall n \geq n_0$ where n_0 is usually 0 or a positive number

1. Basis step: P_{n_0}

2. Induction step:

$$P_{n_0} \& P_{n-1} \implies P_n$$

Assume P_{n_0} and P_{n-1} then show P_n

Set Theory

set: a collection of elements

$x \in A$, where x is element, A is set, $\in \equiv$ Element hood

$$A = \{a_1, a_2, \dots, a_n\}$$

Subset: $A \subset X, B \subset X$

$A \subset X$ if and only if $\forall x \in A, x \in X$

$$A^c = \{x \in X : x \notin A\}$$

$$A \cup B = \{x \in X : x \in A \text{ OR } x \in B\}$$

$$A \cap B = \{x \in X : x \in A \text{ AND } x \in B\}$$


De Morgan's Law:

$$A \cup B = (A^c \cap B^c)^c$$


$$A \cap B = (A^c \cup B^c)^c$$

Function

$$f : X \implies Y$$




Injective function:




f is injective if and only if $\forall x, z \in X, f(x) = f(z) \implies x = z$

Surjective function:



$\forall y \in Y, \exists x \in X : f(x) = y$

Bijjective Function (1-1 correspondence)



f is bijective if and only if f is injective and surjective.

Cardinality of a set

Finite set:

$$A = \{a_1, \dots, a_n\}, \text{ where } n \in \mathbb{Z}^+$$

Infinite set:

1. Uncountably infinite

$$\mathbb{R}$$

2. Countably infinite

$$\mathbb{Z}^+$$

Example:

$$f: \mathbb{Z}^+ (1-1 \text{ correspondence}) \implies \mathbb{Z}^-$$

Vectors

A vector is a 1-dimensional array of scalars over a field.

$$\text{Let } V \in \mathbb{R}^{(n)} : v_1, \dots, v_n \in \mathbb{R}$$

$$\text{For } u, v \in \mathbb{R}^n$$

- Vector Addition:

$$u + v = \begin{bmatrix} u_1 + v_1 \\ \dots \\ u_n + v_n \end{bmatrix} \in \mathbb{R}^n$$

- Scalar Multiplication:


$$\text{For } a \in \mathbb{R}, v \in \mathbb{R}^n$$

$$\text{Then } av = \begin{bmatrix} av_1 \\ \dots \\ av_n \end{bmatrix}$$

- Linear Combination:

$$\text{For } a, b \in \mathbb{R} \text{ and } u, v \in \mathbb{R}^n$$

$$au + bv = \begin{bmatrix} au_1 \\ \dots \\ au_n \end{bmatrix} + \begin{bmatrix} bv_1 \\ \dots \\ bv_n \end{bmatrix} = \begin{bmatrix} au_1 + bv_1 \\ \dots \\ au_n + bv_n \end{bmatrix}$$



- Inner Product


$$u, v \in \mathbb{R}^n$$

$$u \cdot v = \sum_{k=1}^n u_k v_k$$

Length:

$$\|u\|^2 = u \cdot u = \sum_{k=1}^n (u_k)^2$$

$$\|u\| = \sqrt{\sum_{k=1}^n (u_k)^2}$$



$$u = v$$

$$u \cdot v = \|u\| \|v\| \cos \theta$$

$$\|u\|^2 = u \cdot u$$

$$\|v\|^2 = v \cdot v$$

$$u \cdot v = \|u\| \|v\| \cos(\theta)$$

$$\|v\|^2 = v \cdot v$$

$$\cos(\theta) = \frac{u \cdot v}{\|u\| \|v\|}$$

Thm: Cauchy Schwartz Inequality

$$\text{Let } u, v \in \mathbb{R}^n, |u \cdot v| \leq \|u\| \|v\|$$

Proof:

Case 1: $\|u\| = 0$ or $\|v\| = 0$

$$\text{If } \|u\| = 0 : |0 \cdot v| = 0 \leq \|u\| \|v\| = 0 \|v\| = 0$$

$$\text{If } \|v\| = 0 : |u \cdot 0| = 0 \leq \|u\| \|v\| = \|u\| 0 = 0$$

Case 2: $\|u\| \neq 0$ and $\|v\| \neq 0$

Lemma 1: If $a, b \in \mathbb{R}$, then $a^2 + b^2 \geq 2ab$

Proof: $(a - b)^2 \geq 0$ for $a, b \in \mathbb{R}$

$$a^2 + b^2 - 2ab \geq 0$$

$$a^2 + b^2 \geq 2ab$$

Lemma 2: If $a, b \in \mathbb{R}$, then $a^2 + b^2 \geq -2ab$

Proof: $(a + b)^2 \geq 0$ for $a, b \in \mathbb{R}$

$$a^2 + b^2 + 2ab \geq 0$$

$$a^2 + b^2 \geq -2ab$$

$$\text{Let } a_k \equiv \frac{u_k}{\|u\|}, b_k \equiv \frac{v_k}{\|v\|}$$

$$(a_k)^2 + (b_k)^2 \geq 2a_k b_k \quad \text{using Lemma 1}$$

$$\sum_{k=1}^n \left(\frac{(u_k)^2}{(\|u\|)^2} + \frac{(v_k)^2}{(\|v\|)^2} \right) \geq \sum_{k=1}^n \left(2 \frac{u_k}{\|u\|} \frac{v_k}{\|v\|} \right)$$

$$\frac{1}{(\|u\|)^2} \sum_{k=1}^n (u_k)^2 + \frac{1}{(\|v\|)^2} \sum_{k=1}^n (v_k)^2 \geq \frac{2}{\|u\|\|v\|} \sum_{k=1}^n u_k v_k$$

$$\frac{(\|u\|)^2}{(\|u\|)^2} + \frac{(\|v\|)^2}{(\|v\|)^2} \geq \frac{2}{\|u\|\|v\|} (u \cdot v)$$

$$2 \geq \frac{2}{\|u\|\|v\|} (u \cdot v)$$

$$\|u\| \|v\| \geq (u \cdot v)$$

Similarly,

$$\|u\| \|v\| \geq -(u \cdot v) \quad \text{using Lemma 2}$$

Therefore $\|u\| \|v\| = 0$

Week 1 Session 2

Outline

Vectors: Dot Products, Norm, Minkowski Inequality

Matrices: Matrix multiplication, Transpose, Trace, Block matrices

$$u, v \in \mathbb{R}^n$$

$$\text{Inner Product: } u \cdot v = \sum_{k=1}^n u_k v_k$$

$$\text{Length (Norm): } \|u\|^2 = u \cdot u = \sum_{k=1}^n (u_k)^2$$

Properties: For $k \in \mathbb{R}$, $u, v, w \in \mathbb{R}^n$

$$1. u \cdot v = v \cdot u$$

$$2. u \cdot (v + w) = (u \cdot v) + (u \cdot w)$$

$$3. ku \cdot v = k(u \cdot v)$$

$$4. u \cdot u \geq 0 \text{ and } u \cdot u = 0 \text{ if and only if } u = \mathbf{0}$$

$$|u \cdot v| \leq \|u\| \|v\| : \text{Cauchy Schwartz Inequality}$$

Minkowski Inequality

$$||u + v|| \leq ||u|| + ||v||$$

Proof:


$$\begin{aligned} ||u + v||^2 &= (u + v) \cdot (u + v) \\ &= (u \cdot u) + (u \cdot v) + (v \cdot u) + (v \cdot v) \\ &= ||u||^2 + 2(u \cdot v) + ||v||^2 \\ &\leq ||u||^2 + 2|u \cdot v| + ||v||^2 \quad (u \cdot v) \in \mathbb{R} \\ &\leq ||u||^2 + 2||u|| ||v|| + ||v||^2 \quad \text{Cauchy Schwartz Inequality} \\ &= (||u|| + ||v||)^2 \end{aligned}$$

Therefore:

$$\begin{aligned} ||u + v||^2 &\leq (||u|| + ||v||)^2 \\ ||u + v|| &\leq ||u|| + ||v|| \end{aligned}$$

u and v are orthogonal (perpendicular) $\implies u \cdot v = 0$

Normalizing a vector: $\frac{v}{||v||}$



$u^* \equiv$ Projection of u onto v

$$u^* \equiv Proj(u, v) = \frac{u \cdot v}{||v||^2} v$$

$u^* \equiv Proj(u, v) = ||u|| \frac{v}{||v||}$, where $||u||$ is the magnitude, $\frac{v}{||v||}$ is the direction

$$\begin{aligned} &= ||u|| \cos(\theta) \frac{v}{||v||} \\ &= ||u|| ||v|| \cos(\theta) \frac{v}{||v||^2} \\ &= \frac{u \cdot v}{||v||^2} v \end{aligned}$$

Complex Vectors

$$u, v \in \mathbb{C}^n$$

$$u \cdot v = \sum_{k=1}^n u_k v_k^*$$

where $v_k \in \mathbb{C}$, $v_k = a_k + j b_k$, where a_k is the real part, and b_k is the imaginary part

Matrices

$$A \equiv [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

A is $m \times n$ with m rows and n columns

$$A \in \mathbb{R}^{m \times n}$$

$$A = \begin{bmatrix} 2 & 1 & 0 \\ 4 & 2 & -1 \\ 3 & 3 & 0 \\ 2 & 4 & 2 \end{bmatrix} \in \mathbb{R}^{4 \times 3}$$

A row vector: $v = [v_1, v_2, \dots, v_n] \in K^{1 \times n}$

$$\text{A column vector: } v = \begin{bmatrix} v_1 \\ v_2 \\ \dots \\ v_m \end{bmatrix} \in K^{m \times 1}$$

Matrix Addition

$$\begin{aligned} A, B \in K^{m \times n} \quad A + B &= \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} + \begin{bmatrix} b_{11} & b_{12} & \dots & b_{1n} \\ \dots & \dots & \dots & \dots \\ b_{m1} & b_{m2} & \dots & b_{mn} \end{bmatrix} \\ &= \begin{bmatrix} a_{11} + b_{11} & a_{12} + b_{12} & \dots & a_{1n} + b_{1n} \\ \dots & \dots & \dots & \dots \\ a_{m1} + b_{m1} & a_{m2} + b_{m2} & \dots & a_{mn} + b_{mn} \end{bmatrix} \end{aligned}$$

Scalar Multiplication

If $k \in K, A \in K^{m \times n}$

$$kA = \begin{bmatrix} ka_{11} & ka_{12} & \dots & ka_{1n} \\ \dots & \dots & \dots & \dots \\ ka_{m1} & ka_{m2} & \dots & ka_{mn} \end{bmatrix}$$

Null Matrix

$$A \equiv [a_{ij}] = \mathbf{0}$$

$$\forall i, j, a_{ij} = 0$$

$$A = \begin{bmatrix} 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 \end{bmatrix}$$

Linear Combination:

$$a, b \in K, A, B \in K^{M \times N}$$

$$aA + bB = \begin{bmatrix} aa_{11} + bb_{11} & aa_{12} + bb_{12} & \dots & aa_{1n} + bb_{1n} \\ \dots & \dots & \dots & \dots \\ aa_{m1} + bb_{m1} & aa_{m2} + bb_{m2} & \dots & aa_{mn} + bb_{mn} \end{bmatrix}$$

Properties:

If $k, k' \in K$ and $A, B, C \in K^{m \times n}$

1. $A + B = B + A$ Commutativity
2. $A + (B + C) = (A + B) + C$ Associativity
3. $k(A + B) = kA + kB$
4. $kk'A = k(k'A)$

$$5. A + -A = 0$$

$$6. A + 0 = A$$

Transpose:

If $A \in K^{m \times n}$ and $A = [a_{ij}]$, then

$A^T \in K^{n \times m}$ and $A^T = B = [b_{ij}]$ where $b_{ij} = a_{ji}$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix} \text{ where dimension is } m \times n$$

$$A^T = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{m1} \\ \dots & \dots & \dots & \dots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{bmatrix} \text{ where dimension is } n \times m$$

Example.

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$$

then

$$A^T = \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}$$

Properties:

If $A, B \in K^{m \times n}$

$$1. (A + B)^T = A^T + B^T$$

$$2. (A^T)^T = A$$

Let $u, v \in K^{m \times 1}$

then $u \cdot v = u^T v$

Square Matrix

$A = [a_{ij}]$ is a square matrix if and only if the number of rows equal the number of columns.

$$m = n$$

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

Diagonal Matrix:

$$A \equiv [a_{ij}]$$

A square matrix such that $\forall i \neq j, a_{ij} = 0$

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

Triangular Matrices:

Upper triangular

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

$$\forall i > j, a_{ij} = 0$$

Lower triangular

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\forall i < j, a_{ij} = 0$$

Matrix Multiplication

| A | B | C |
|--------------|--------------|--------------|
| $m \times n$ | $n \times p$ | $m \times p$ |
| $[a_{ij}]$ | $[b_{ij}]$ | $[c_{ij}]$ |

$$c_{ij} = \sum_{k=1}^n a_{ik} b_{kj}$$

| B | A | D |
|--------------|--------------|--------------|
| $n \times m$ | $m \times p$ | $n \times p$ |
| $[a_{ij}]$ | $[b_{ij}]$ | $[c_{ij}]$ |

$$BA = D$$

$$d_{ij} = \sum_{k=1}^m b_{ik} a_{kj}$$

$$c_{ij} = i^{\text{th}} \text{ row of } A \cdot j^{\text{th}} \text{ column of } B$$

$$i^{\text{th}} \text{ row of } A: A_{i:}$$

$$j^{\text{th}} \text{ column of } B: B_{:j}$$

$$\begin{bmatrix} \vdots & \text{A}_{i:} & \vdots \end{bmatrix} \begin{bmatrix} \vdots & \text{B}_{:j} & \vdots \end{bmatrix} = \begin{bmatrix} c_{ij} \end{bmatrix}$$

Properties: If A, B, C are conformable for multiplication

1. $(AB)C = A(BC)$ Associativity
2. $A(B + C) = AB + AC$ Left distribution
3. $(A + B)C = AC + BC$ Right distribution
4. $(AB)^T = B^T A^T$
5. $c(AB) = (cA)B = A(cB)$ if c is a scalar
6. $AB \neq BA$

Trace

$$A \in K^{n \times n}$$

$$\text{Tr}(A) = \sum_{k=1}^n a_{kk}$$

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 0 & 2 & 5 \end{bmatrix}$$

$$\text{Tr}(A) = a_{11} + a_{22} + a_{33} = 1 + 1 + 5 = 7$$

Properties:

If A, B, C are conformable for multiplication

1. $Tr(A) = Tr(A^T)$
2. $Tr(BA) = Tr(AB)$
3. $Tr(ABC) = Tr(BCA) = Tr(CAB)$

Cyclic Property of Trace

Thm:

$$Tr(A_1 A_2 \dots A_{n-1} A_n) = Tr(A_n A_1 \dots A_{n-1})$$

If the matrices A_k are conformable for matrix multiplication where Tr is the trace operator:

$$Tr(A) = \sum_{k=1}^p a_{kk} \text{ if } A \text{ is a square matrix}$$

$$A_k \in \mathbb{C}^{m_k \times n_k}$$

Proof:

Lemma 1:

$$Tr(AB) = Tr(BA)$$

Lemma 2:

$$A \times (B \times C) = (A \times B) \times C$$

Lemma 1:

A dimension is $m \times n$

B dimension is $n \times m$

Then $A \times B$ is $m \times m$, $B \times A$ is $n \times n$

$$\begin{aligned} Tr(AB) &= \sum_{k=1}^m (AB)_{kk} && \text{def of } Tr \\ &= \sum_{k=1}^m (\sum_{l=1}^n a_{kl} b_{lk}) && \text{def of matrices, multiplication} \\ &= \sum_{k=1}^m \sum_{l=1}^n a_{kl} b_{lk} && \text{distribution} \\ &= \sum_{l=1}^n \sum_{k=1}^m a_{kl} b_{lk} && \text{finite sum} \\ &= \sum_{l=1}^n \sum_{k=1}^m b_{lk} a_{kl} && \text{complex number} \\ &= \sum_{l=1}^n (\sum_{k=1}^m b_{lk} a_{kl}) && \text{distribution} \\ &= \sum_{l=1}^n (BA)_{ll} && \text{def of matrix multiplication} \\ &= Tr(BA) && \text{def of } Tr \end{aligned}$$

Lemma 2:

A dimension is $u \times v$

B dimension is $v \times w$

C dimension is $w \times r$

Then $A \times (B \times C)$ is $u \times r$, $(A \times B) \times C$ is $u \times r$

say $M \equiv [m_{ij}]$, $N \equiv [n_{ij}]$

$$m_{ij} = n_{ij}$$

$$m_{ij} = (A(BC))_{ij}$$

$$\begin{aligned}
&= \sum_{k=1}^v a_{ik} (BC)_{kj} && \text{def of matrix multiplication} \\
&= \sum_{k=1}^v a_{ik} \left(\sum_{l=1}^w b_{kl} c_{lj} \right) && \text{def of matrix multiplication} \\
&\text{where } \left(\sum_{l=1}^w b_{kl} c_{lj} \right) = (BC)_{kj} \\
&= \sum_{k=1}^v \sum_{l=1}^w a_{ik} b_{kl} c_{lj} && \text{distribution} \\
&= \sum_{l=1}^w \left(\sum_{k=1}^v a_{ik} b_{kl} \right) c_{lj} && \text{finite sum} \\
&\text{where } \left(\sum_{k=1}^v a_{ik} b_{kl} \right) = (AB)_{il} \\
&= \sum_{l=1}^w (AB)_{il} c_{lj} && \text{def of matrix multiplication} \\
&= ((AB)C)_{ij} && \text{def of matrix multiplication} \\
&= n_{ij}
\end{aligned}$$

$$\begin{aligned}
&Tr(A_1 A_2 \dots A_{n-1} A_n) = Tr((A_1 A_2 \dots A_{n-1}) A_n) \\
&= Tr(A_n (A_1 A_2 \dots A_{n-1})) \\
&= Tr(A_n A_1 \dots A_{n-1})
\end{aligned}$$

| A | B | $A + B$ |
|--------------|--------------|--------------|
| $n \times n$ | $n \times n$ | $n \times n$ |
| diagonal | diagonal | diagonal |
| triangular | triangular | triangular |
| upper | upper | upper |
| lower | lower | lower |

Invertible Matrices

A is invertible if and only if $\exists B : AB = BA = I_n$

$$I = \begin{bmatrix} 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

Properties:

1. $A^{-1}A = I_n$
 2. $(AB)^{-1} = B^{-1}A^{-1}$
 3. $(A^T)^{-1} = (A^{-1})^T$
-

$$A \in \mathbb{C}^{m \times n}$$

Hermitian

$$A^H = (A^*)^T = (A^T)^*$$

$$\text{If } A \in \mathbb{R}^{m \times n}, A^H = (A^*)^T = (A^T)^*$$

Normal Matrices

$$A^T A = A A^H$$

Complex:

- Hermitian matrices: $A = A^H$
- Skew Hermitian: $A = -A^H$

- Unitary: $A^{-1} = A^H$

Real:

- Symmetric: $A = A^T$
- Skew symmetric: $A = -A^T$
- Orthogonal: $A^{-1} = A^T$

Block Matrices

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \\ a_{41} & a_{42} & a_{43} & a_{44} \end{bmatrix}$$

Week 2 Session 1

Outlines

Linear System: Lines, Hyperplane, Normal

Equivalent Systems: Elementary row operations


Echelon Form: Gaussian Elimination

Row Canonical Form: Gauss-Jordan

Located Vectors


$$P(u_1, \dots, u_n)$$

$$Q(v_1, \dots, v_n)$$



$$\vec{PQ} = \vec{Q} - \vec{P} = \begin{bmatrix} v_1 \\ \dots \\ v_n \end{bmatrix} - \begin{bmatrix} u_1 \\ \dots \\ u_n \end{bmatrix} = \begin{bmatrix} v_1 - u_1 \\ \dots \\ v_n - u_n \end{bmatrix}$$

Lines



$$L = \{x \in \mathbb{R}^n : x = p + ut, t \in \mathbb{R}^n\}$$

L is a line that passes through point P with direction $u \in \mathbb{R}^n$

Linear Systems

Linear Equation

$$a_1x_1 + \dots + a_nx_n = b$$

$$\sum_{j=1}^n a_jx_j = b$$

where a_j are the coefficients, and x_j are the unknowns


Hyperplane H :

$$H = \{x \in \mathbb{R}^n : \sum_{j=1}^n a_jx_j = b\}$$

Example:

$$6x = 6, H = \{1\}$$

$$x + y = 2$$



$$x + y + z = 1$$

Normal to H : $\sum_{j=1}^n a_jx_j = b$

$w \in \mathbb{R}^n$ such that for all any located vector \overrightarrow{PQ} in H , w is orthogonal to \overrightarrow{PQ}

$$w = \begin{bmatrix} a_1 \\ \dots \\ a_n \end{bmatrix}$$

Proof:

$$\sum_{j=1}^n a_jx_j = b$$

$$P(u_1, \dots, u_n) \in H \implies \sum_{j=1}^n a_ju_j = b$$

$$Q(v_1, \dots, v_n) \in H \implies \sum_{j=1}^n a_jv_j = b$$

$$w \perp \overrightarrow{PQ}$$

$$w = \begin{bmatrix} a_1 \\ \dots \\ a_n \end{bmatrix}$$

$$w \cdot \overrightarrow{PQ} = \begin{bmatrix} a_1 \\ \dots \\ a_n \end{bmatrix} \cdot \begin{bmatrix} v_1 - u_1 \\ \dots \\ v_n - u_n \end{bmatrix}$$

$$= \sum_{j=1}^n a_j(v_j - u_j)$$

$$= \sum_{j=1}^n a_j v_j - \sum_{j=1}^n a_j u_j$$

$$= b - b$$

$$= 0$$

Linear Systems

A list of linear equations with the same unknowns

m equations and n unknowns

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

...

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

- Unique solution
- Infinite solution
- No solution

| A | x | b |
|--------------|--------------|--------------|
| $m \times n$ | $n \times 1$ | $m \times 1$ |

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

$$x = \begin{bmatrix} x_1 \\ \dots \\ x_n \end{bmatrix}$$

$$b = \begin{bmatrix} b_1 \\ \dots \\ b_m \end{bmatrix}$$

Degenerate linear equation:

$$0x_1 + \dots + 0x_n = b$$

1. $b = 0$, every $x \in \mathbb{R}^n$ is a solution
2. $b \neq 0$, no solution

Homogenous system: $Ax = b = \mathbf{0}$

Equivalent Systems

$Ax = b$, $A'x = b'$ where x is in dimension $n \times 1$

Theorem:

Let L be a linear combination of the equations m $Ax = b$, then x is a solution to L

Proof:

$$Ax = b$$

$$\sum_{j=1}^n a_{ij}x_j = b_i \text{ where } 1 \leq i \leq m$$

$$\text{Let } s = \begin{bmatrix} s_1 \\ \dots \\ s_n \end{bmatrix} \text{ is a solution to } Ax = b$$

$$\text{Then: } \sum_j \sum_{j=1}^n a_{ij}x_j = \sum_j b_i \quad \text{Integration}$$

$$\begin{aligned}
 \sum_{i=1}^m c_i (\sum_{j=1}^n a_{ij} s_j) &= \sum_{i=1}^m \sum_{j=1}^n c_i a_{ij} s_j \\
 &= \sum_{j=1}^n (\sum_{i=1}^m c_i a_{ij}) s_j \\
 &= \sum_{j=1}^m c_j b_j
 \end{aligned}$$

x is also a solution to L

$$Ax = b \text{ Linear combination} \rightarrow A'x = b'$$

Elementary Row Operations

1. Row swap: $R_i \leftrightarrow R_j$
2. Scalar multiplication: $R_i \rightarrow kR_i$
3. Sum of a row with a scalar multiple of another row: $R_i \rightarrow R_i + kR_j$

Thm:

$Ax = b$ and $A'x = b'$ where A' (b') is obtained from the elementary row operations on $Ax = b$ then they have same solutions.

Geometry: Linear System Solutions

$$Ax = b$$

Row:

$$\sum_{j=1}^n a_{ij} x_j = b_i$$

$$\text{Row 1: } a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1$$

$$\text{Row 2: } a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2$$

...


$$\text{Row m: } a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m$$

Example 1:

$$x + y = 2$$

$$x - y = 0$$

$x = 1, y = 1$ is the unique solution




Example 2:

$$x + y = 2$$

$$2x + 2y = 4$$

Infinite solution




Example 3:

$$x + y = 2$$

$$x + y = 0$$

No solution



Column

$$Ax = b$$

$$A = \begin{bmatrix} \dots & \dots & \dots & \dots \\ a_{11} & a_{12} & \dots & a_{1n} \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

$$x = \begin{bmatrix} x_1 \\ \dots \\ x_n \end{bmatrix}$$

$$b = \begin{bmatrix} b_1 \\ \dots \\ b_m \end{bmatrix}$$


$$\sum_{j=1}^n A_{ij}x_j = b$$

Example 1:

$$x + y = 2$$

$$x - y = 0$$

$$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$$




Example 2:

$$x + y = 2$$

$$2x + 2y = 4$$

$$\begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$



Echelon Form

$$2x_1 + 3x_2 + x_3 + x_4 - x_5 = 2$$

$$x_2 + x_3 + 0x_4 + x_5 = 2$$

$$x_4 + x_5 = 1$$

$$m = 3, n = 5$$

Pivot variables: x_1, x_2, x_4 (leading variables)

Free variables: x_3, x_5 (non-leading variables)

Special case (Triangular Form)

$$2x_1 + 3x_2 + 4x_3 = 5$$

$$2x_2 + x_3 = 6$$

$$3x_3 = 1$$

$$m = 3, n = 3$$

Gaussian Elimination

Two step process for solving linear systems of form $Ax = b$

1. Forward elimination: Reduce to Echelon Form
2. Backward substitution

Example 1:

$$R1 : 2x + y + z = 5$$

$$R2 : 4x - 6y = -2$$

$$R3 : -2x + 7y + 2z = 9$$

Forward Elimination:

$$R1 : R1$$

$$R2 : R2 - 2R1$$

$$R3 : R3 + R1$$

$$2x + y + z = 5$$

$$0x - 8y - 2z = -12$$

$$0x + 8y + 3z = 14$$

$$R1 : R1$$

$$R2 : R2$$

$$R3 : R3 + R2$$

$$2x + y + z = 5$$

$$0x - 8y - 2z = -12$$

$$0x + 0y + z = 2$$

Backward Substitution:

$$z = 2$$

$$y = 1$$

$$x = 1$$

Augmented Matrix (M)

| A | x | b | M |
|--------------|--------------|--------------|--------------------|
| $m \times n$ | $n \times 1$ | $m \times 1$ | $m \times (n + 1)$ |

$$M \equiv [A \mid b]$$

$$M = \left[\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 4 & -6 & 0 & -2 \\ -2 & 7 & 2 & 9 \end{array} \right]$$

$$\text{Where } A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}, b = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

Echelon Matrix:

$$M = \left[\begin{array}{cccc} 2 & 1 & 2 & 1 \\ 0 & 4 & 3 & 2 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 5 \end{array} \right]$$

Week 2 Session 2

Outline

Row Canonical Form: Gauss Jordan Elimination

Elementary Matrix Operations

LU Decomposition: LDU

Vector Spaces

Echelon Matrix

$$\left[\begin{array}{ccccc} 1 & 1 & 2 & 3 & 5 \\ 0 & 2 & 1 & 4 & -1 \\ 0 & 0 & 0 & 2 & 1 \end{array} \right]$$

Augmented Matrix

$$Ax = b, M = [A|b]$$

Row Canonical Form (Row-reduced Echelon Form)

1. Echelon Form
2. All non zero leading elements must be equal to 1
3. All the other values above and below a leading element must be 0

$$\left[\begin{array}{ccccc} 1 & 0 & 3 & 0 & 1 \\ 0 & 1 & 2 & 0 & 2 \\ 0 & 0 & 0 & 1 & 2 \end{array} \right]$$

$$M = [A|b]$$

Gauss-Jordan Elimination

$$Ax = b$$

$$M = [A|b] \text{ - Augmented matrix}$$

Reduce M to its row canonical form

$$M' = [A'|b'] \text{ (i.e., } A'x = b')$$

Example:

$$2x_1 + x_2 + x_3 = 5$$

$$4x_1 - 6x_2 = -2$$

$$-2x_1 + 7x_2 + 2x_3 = 9$$

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}$$

$$b = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

$$M \equiv [A|b] = \left[\begin{array}{ccc|c} 2 & 1 & 1 & 5 \\ 4 & -6 & 0 & -2 \\ -2 & 7 & 2 & 9 \end{array} \right]$$

$$R1 : R1$$

$$R2 : R2 - 2R1$$

$$R3 : R3 + R1$$

$$\begin{bmatrix} 2 & 1 & 1 & 5 \\ 0 & 8 & -2 & -12 \\ 0 & 8 & 3 & 14 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2$$

$$R3 : R3 + R2$$

$$\begin{bmatrix} 2 & 1 & 1 & 5 \\ 0 & 8 & -2 & -12 \\ 0 & 0 & 1 & 2 \end{bmatrix} \text{ which is the Echelon Form}$$

$$R1 : R1 - R3$$

$$R2 : R2 + 2R3$$

$$R3 : R3$$

$$\begin{bmatrix} 2 & 1 & 0 & 3 \\ 0 & -8 & 0 & -8 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : -1/8R2$$

$$R3 : R3$$

$$\begin{bmatrix} 2 & 1 & 0 & 3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

$$R1 : R1 - R2$$

$$R2 : R2$$

$$R3 : R3$$

$$\begin{bmatrix} 2 & 0 & 0 & 2 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

$$R1 : 1/2R1$$

$$R2 : R2$$

$$R3 : R3$$

$$\begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}, \text{ which is in row canonical form}$$

$$x_1 = 1, x_2 = 2, x_3 = 2$$

Linear combination of orthogonal vectors

Let $u_1, u_2, \dots, u_n \in \mathbb{R}^n$ are mutually orthogonal

For any vector $v \in \mathbb{R}^n$

$$v = u_1x_1 + \dots + u_nx_n$$

where $x_i = \frac{v \cdot u_i}{||u_i||^2}$ and $u_i \neq \mathbf{0}$ for $1 \leq i \leq n$

$$A = \begin{bmatrix} \dots & \dots & \dots & \dots \\ u_1 & u_2 & \dots & u_n \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

$Ax = v$ what is x ?

Proof:

$$u_i \cdot u_j = \begin{cases} 0, & \text{if } i \neq j \\ ||u_i||^2, & \text{if } i = j \end{cases} \quad \text{Equation 1}$$

$$Ax = v$$

$$\sum_{j=1}^n x_j u_j = v \quad \text{Equation 2}$$

$$v \cdot u_i = \sum_{j=1}^n x_j u_j \cdot u_i$$

$$= \sum_{j=1}^n x_j (u_j \cdot u_i)$$

$$= (u_i \cdot u_i)x_i + \sum_{j=1, j \neq i}^n x_j (u_i \cdot u_j)$$

$$= ||u_i||^2 x_i$$

Therefore, $v \cdot u_i = ||u_i||^2 x_i$ means that $x_i = \frac{v \cdot u_i}{||u_i||^2}$

$$v = \sum_{j=1}^n x_j u_j = \sum_{j=1}^n \frac{v \cdot u_j}{\|u_j\|^2} u_j$$

Inverse Matrix

Using Gauss Jordan Elimination for A^{-1}

If A ($n \times n$) is invertible, $\exists A^{-1}$ such that $AA^{-1} = I$

$$AA^{-1} = I$$

$$\text{say } B = A^{-1}$$

$$A = \begin{bmatrix} \dots & \dots & \dots & \dots \\ a_1 & a_2 & \dots & a_n \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

$$A = \begin{bmatrix} \dots & \dots & \dots & \dots \\ b_1 & b_2 & \dots & b_n \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

$$\begin{bmatrix} \dots & \dots & \dots & \dots \\ a_1 & a_2 & \dots & a_n \\ \dots & \dots & \dots & \dots \end{bmatrix} \begin{bmatrix} \dots & \dots & \dots & \dots \\ b_1 & b_2 & \dots & b_n \\ \dots & \dots & \dots & \dots \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 \end{bmatrix}$$

$$Ab_1 = \begin{bmatrix} 1 \\ 0 \\ \dots \\ 0 \end{bmatrix}$$

$$Ab_2 = \begin{bmatrix} 0 \\ 1 \\ \dots \\ 0 \end{bmatrix}$$

$$M = [A|I] \text{ Row canonical} \rightarrow [I|A^{-1}]$$

Example 1:

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \text{ Find } A^{-1}$$

$$M \equiv \begin{bmatrix} 2 & 1 & 1 & 1 & 0 & 0 \\ 4 & -6 & 0 & 0 & 1 & 0 \\ -2 & 7 & 2 & 0 & 0 & 1 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2$$

$$R3 = R3 + R2$$

$$\begin{bmatrix} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 8 & 3 & 1 & 0 & 1 \end{bmatrix}$$

$$R1 : R1 - R3$$

$$R2 : R2$$

$$R3 : R3 + R2$$

$$\begin{bmatrix} 2 & 1 & 1 & 1 & 0 & 0 \\ 0 & -8 & -2 & -2 & 1 & 0 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{bmatrix}$$

$$R1 : R1 - R3$$

$$R2 : R2 + 2R3$$

$$R3 : R3$$

$$\begin{bmatrix} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & -8 & 0 & -4 & 3 & 2 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{bmatrix}$$

$$R1 : R1$$

$$R2 = -1/8R2$$

$$R3 = R3$$

$$\begin{bmatrix} 2 & 1 & 0 & 2 & -1 & -1 \\ 0 & 1 & 0 & 1/2 & -3/8 & -1/4 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{bmatrix}$$

$$R1 : R1 - R2$$

$$R2 : R2$$

$$R3 = R3$$

$$\begin{bmatrix} 2 & 0 & 0 & 3/2 & -5/8 & -3/4 \\ 0 & 1 & 0 & 1/2 & -3/8 & -1/4 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{bmatrix}$$

$$R1 : 1/2R1$$

$$R2 : R2$$

$$R3 : R3$$

$$\begin{bmatrix} 1 & 0 & 0 & 3/4 & -5/16 & -3/8 \\ 0 & 1 & 0 & 1/2 & -3/8 & -1/4 \\ 0 & 0 & 1 & -1 & 1 & 1 \end{bmatrix}$$

$$\text{where } A^{-1} = \begin{bmatrix} 3/4 & -5/16 & -3/8 \\ 1/2 & -3/8 & -1/4 \\ -1 & 1 & 1 \end{bmatrix}$$

Check:

$$AA^{-1} = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix} \begin{bmatrix} 3/4 & -5/16 & -3/8 \\ 1/2 & -3/8 & -1/4 \\ -1 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$


Elementary Matrix Operations

$$eA \equiv EA$$

where e is the elementary row operation, E is the elementary matrix operation

$$e_n \dots e_1 A = E_n \dots E_1 A$$

1. Row Swap $R_i \leftrightarrow R_j$



$$EA = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} A = \begin{bmatrix} \dots b_1 : \\ \dots b_2 : \\ \dots b_3 : \end{bmatrix}$$


Let $E = I$

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$EA \equiv B$ where $B = [b_{ij}]$

$$\sum_{k=1}^n e_{ik} a_{kj} = b_{ij}$$

where $e_{ik} = [e_{i1}, e_{i2}, \dots, e_{in}]$



$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

2. Scalar Multiplication of a row

$R_i : kR_i$

$EA = B$

$R1 : R1$

$R2 : kR2$

$R3 : R3$

$$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & k & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } E^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/k & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

3. Row addition with a scalar multiple of another row

| Operation | E | E^{-1} |
|-----------|------------|------------------|
| $R1$ | $R1$ | $R1$ |
| $R2$ | $R2 + kR3$ | $R2 + kR3 - kR3$ |
| $R3$ | $R3$ | $R3$ |

This is an operation of E and E^{-1}

$$E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & k \\ 0 & 0 & 1 \end{bmatrix} \text{ and } E^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -k \\ 0 & 0 & 1 \end{bmatrix}$$

LU decomposition

$$A = LU \equiv LDU$$

where A is in dimension $n \times n$, L is the lower triangular, U is the upper triangular, D is the diagonal matrix

A is a nonsingular matrix that can be reduced into triangular from U only row-addition operations

Example:

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}$$

$$e_n \dots e_1 A = U = E_n \dots E_1 A$$

$$E_n \dots E_1 A = U$$

$$(E_n \dots E_1)^{-1} = E_1^{-1} E_2^{-1} \dots E_n^{-1}$$

$$(E_n \dots E_1)^{-1} (E_n \dots E_1) A = E_1^{-1} E_2^{-1} \dots E_n^{-1} U$$

$$LHS : A = LU$$

$$RHS = LU$$

$$R1 : R1$$

$$R2 : R2 - 2R1$$

$$R3 : R3 + R1$$

$$\begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 8 & 3 \end{bmatrix}$$

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

| Operations | E_1 | E_1^{-1} |
|------------|-------------------|------------|
| $R1$ | $R1$ | $R1$ |
| $R2$ | $R2 - 2R1 (+2R1)$ | $R2$ |
| $R3$ | $R3 + R1 (-R1)$ | $R3$ |

$$E_1^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2$$

$$R3 : R3 + R2$$

$$\begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

$$E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

$$(E_2 E_1)A = U$$

$$A = (E_1^{-1} E_2^{-1})U \text{ and } E_1^{-1} E_2^{-1} = L$$

| Operations | E_1 | E_1^{-1} |
|------------|----------------|------------|
| $R1$ | $R1$ | $R1$ |
| $R2$ | $R2$ | $R2$ |
| $R3$ | $R3 + R2(-R2)$ | $R3$ |

$$E_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

$$L = E_1^{-1} E_2^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix}$$

Check:

$$LU = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}$$

Week 3 Session 1

Outlines

LU Decomposition: LDU

Vector Spaces: Fields, Span, Subspaces

Linear Independence: Invertibility

Uniqueness Theorem

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ 2 & 7 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 1 \\ 0 & -8 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

$$A = LU$$

$$A = LDU$$

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 \\ 0 & -8 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1/2 & 1/2 \\ 0 & 1 & 1/4 \\ 0 & 0 & 1 \end{bmatrix}$$

Vector Spaces

Field:

A field F is a collection of elements such that for binary operations: $+$, \times

We have the following: $\forall a, b, c \in F$

- $a + b = b + a$; $a \cdot b = b \cdot a$
- $a + (b + c) = (a + b) + c$; $a \cdot (b \cdot c) = (a \cdot b) \cdot c$
- $\exists 0 \in F: a + 0 = a$
 $\exists 1 \in F: a \cdot 1 = a$
- $\exists a' \in F: a + a' = 0$

$$5. a \times \frac{1}{a} = 1 \text{ if } a \neq 0$$

$$6. a \cdot (b + c) = (a \cdot b) + (a \cdot c)$$

Example:

$\mathbb{R}, \mathbb{Q}, \mathbb{C}$ - field

\mathbb{Z} not a field ($5^{th} \frac{1}{a} \notin \mathbb{Z}$)

A vector V over field F is a collection of elements $\{\alpha, \beta, \gamma, \dots\}$ (typically called vectors) and collection of elements $\{a, b, c, \dots\} \in F$ called scalars such that:

- Commutative group for $(V, +)$

$$1. \alpha + \beta \in V$$

$$2. \alpha + \beta = \beta + \alpha$$

$$3. \alpha + (\beta + \gamma) = (\alpha + \beta) + \gamma$$

$$4. \forall \alpha, \exists \alpha' \in V : \alpha + \alpha' = \mathbf{0}$$

$$5. \exists \mathbf{0} \in V : \forall \alpha \in V, \mathbf{0} + \alpha = \alpha$$

- Properties for combination of $+$ and \times

$$1. a\alpha \in V$$

$$2. a(b\alpha) = (ab)\alpha$$

$$3. a(\alpha + \beta) = a\alpha + a\beta$$

$$4. (a + b)\alpha = a\alpha + b\alpha$$

$$5. \exists 1 \in F : 1\alpha = \alpha$$

1

K is field, K^n

$$\alpha, \beta \in K^n$$

$$\alpha = \begin{bmatrix} a_1 \\ \dots \\ a_n \end{bmatrix}, a_i \in K$$

2

Polynomial Space: $P(t)$

$$p(t) \in P(t)$$

$$p(t) = a_0 + a_1 t^1 + a_2 t^2 + \dots + a_s t^s$$

where $s \in \{1, 2, 3, \dots\}$

3

Matrix over a field: $K_{m \times n}$

$$A \in K_{m \times n}$$

$$A \equiv [a_{ij}] \text{ where } a_{ij} \in K$$

Linear Combination:

Let $\alpha_1, \alpha_2, \dots, \alpha_n \in V$ where V is a vector space over field F

w is a linear combination of the α_i 's if:

$$w = a_1\alpha_1 + \dots + a_n\alpha_n$$

where $a_1, a_2, \dots, a_n \in F$

Alternatively:

$$Ax = b$$

$$\begin{bmatrix} \dots \\ \alpha_1 \\ \dots \end{bmatrix} x_1 + \begin{bmatrix} \dots \\ \alpha_2 \\ \dots \end{bmatrix} x_2 + \dots + \begin{bmatrix} \dots \\ \alpha_n \\ \dots \end{bmatrix} x_n = w$$

Linear Span

Let $S = \{\alpha_1, \dots, \alpha_n\} \subset V$ for a vector space V over field F

S spans V means that $\forall w \in V, \exists a_1, \dots, a_n \in F$ such that:

$$w = a_1\alpha_1 + \dots + a_n\alpha_n$$

Subspace

u is a subspace of vector space V over field F , if

1. $u \subset V$ (u is a subset of V)
2. u is a vector space over F

Thm:

Let V be a vector space over field F and u is a subset of V ($u \subset V$), If:

1. $0 \in u$
2. $\forall \alpha, \beta \in u, \forall a, b \in F, a\alpha + b\beta \in u$

Then u is a subspace of V

Thm:

Let V be a vector space over field F . If u is a subspace of V , and w is a subspace of u , then w is a subspace of V

Thm:

Intersection of any number of subspaces of a vector V over field F is a subspace of V

Proof:

u_1, u_2, \dots are subspaces of V

u_1 is a subspace of V

u_2 is a subspace of V

...

If $\bigcap_{i=1}^n u_i$ a subspace of V ?

Yes.

Example:

$$w = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\text{where } \alpha_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \alpha_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \alpha_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \quad \checkmark$$

$$\text{where } \alpha_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \alpha_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \quad \times$$

$$\mathbb{R}^2 \equiv \{(x, y) : x \in \mathbb{R}, y \in \mathbb{R}\}$$

$$\{\mathbf{0}\} \text{ subspace of } \mathbb{R}^2 \quad \checkmark$$

$$ax + by = 1 \quad \times$$

$$ax + by = 0 \text{ subspace of } \mathbb{R}^2$$

Thm:

Let $S = \{\alpha_1, \dots, \alpha_n\} \subset V$ where V is a vector space over F and $L(s)$ be the set of all linear combinations of S with respect to F . Then $L(s)$ is a subspace of V .

1. Vector space V over field F
2. $S = \{\alpha_1, \dots, \alpha_n\} \subset V$
3. $L(s) = \{w : w = \sum_{i=1}^n a_i \alpha_i, a_i \in F, \alpha_i \in S\}$

$\implies L(s)$ (span of S) is a subspace of V

Proof:

1. Show that $L(s) \subset V$

$$v \in L(s) \implies v \in V$$

Assume that $v \in L(s)$

$$v = \sum_{i=1}^n a_i \alpha_i \quad \text{- Def of } L(s)$$

$$\alpha_1 \in S \implies \alpha_i \in V \quad \text{- because } S \subset V$$

$$v = \sum_{i=1}^n a_i \alpha_i \in V \quad \text{- } V \text{ is a vector space}$$

$$L(s) \subset V$$

2. Show that $\mathbf{0} \in L(s)$

$$\mathbf{0} = 0\alpha_1 + 0\alpha_2 + \dots + 0\alpha_n = \sum_{i=1}^n 0\alpha_i \in L(s) \quad \text{- Def of } S$$

3. Show that for $v, w \in L(s)$ and $c, d \in F$, $cv + dw \in L(s)$

$$cv + dw = c \sum_{i=1}^n a_i \alpha_i + d \sum_{i=1}^n b_i \alpha_i \text{ where } v = \sum_{i=1}^n a_i \alpha_i \text{ and } w = \sum_{i=1}^n b_i \alpha_i$$

$$= \sum_{i=1}^n ca_i \alpha_i + \sum_{i=1}^n db_i \alpha_i$$

$$= \sum_{i=1}^n (ca_i + db_i) \alpha_i \text{ where } ca_i + db_i \in F$$

Therefore, $cv + dw \in L(s)$

$L(s)$ is a subspace of V

Linear Independence

Let v be a vector space over field F

$$S = \{\alpha_1, \dots, \alpha_n\} \subset v$$

s is a linearly dependent set if there exist a_i 's in F such that:

$$a_1\alpha_1 + a_2\alpha_2, \dots, a_n\alpha_n = \mathbf{0}$$

and at least one of the a_i 's is non-zero

Linearly Independent:

s is linearly independent means that:

$$a_1\alpha_1 + a_2\alpha_2, \dots, a_n\alpha_n = \mathbf{0} \text{ only holds when:}$$

$$a_1 = a_2 = \dots = a_n = 0$$

$Ax = \mathbf{0}$ - Homogenous System

$$A = \begin{bmatrix} \dots & \dots & \dots & \dots \\ \alpha_1 & \alpha_2 & \dots & \alpha_n \\ \dots & \dots & \dots & \dots \end{bmatrix}, x = \begin{bmatrix} x_1 \\ \dots \\ x_n \end{bmatrix}, b = \mathbf{0}$$

Note:

Let $S = \{\alpha_1, \dots, \alpha_n\} \subset v$, then:

1. If $\mathbf{0} \in s$, then s is a linearly dependent set
2. If $s = \{\alpha_1\}$, then s is linearly dependent if and only if $\alpha_1 = \mathbf{0}$

Row Equivalence

A, B are in dimension of $m \times n$

A is row equivalent to B if fB can be obtained from a sequence of elementary row operations of A

Example

A row operations $\implies A'$ (Echelon Form) row operations $\implies A''$ (Row Canonical Form)

Say A in dimension of $n \times n$

Echelon Form

$$L = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 4 & 5 \\ 0 & 0 & 3 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \text{ number of pivots } (1, 2, 3, 1) = n$$

$$R = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 4 & 5 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \text{ number of pivots } (1, 2, 2) < n \text{ Linearly dependent, 0 row (R4)}$$

Row Canonical Form

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = I$$

$$R = \begin{bmatrix} 1 & 0 & x & 0 \\ 0 & 1 & y & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \neq I$$

$$I^{-1} = I$$

$$A \in \mathbb{R}^{n \times n} = \begin{cases} A \sim (\text{Row Equivalent}) I \\ A \approx I \end{cases}$$

$$B = \begin{bmatrix} \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & 0 \end{bmatrix} \text{ (zero row)}, B = \begin{bmatrix} \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ \beta_1 & \beta_2 & \beta_3 & \beta_4 \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

$$BB^{-1} \neq \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = I$$

There is no β_4 such that $i_{44} = 1$

So B^{-1} does not exist

Week 3 Session 2

Outlines

Uniqueness Theorem

Basis and Dimension: Dimension Theorem

Subspaces of a matrix

$$A \in \mathbb{R}^{n \times n}$$

1. Linearly independent rows $\leftrightarrow A \sim I$
2. Linearly dependent rows $\leftrightarrow A \sim B$ such that B^{-1} does not exist

Thm:

Let A be a square matrix, the following statement are equivalent:

1. A is invertible
 2. A is row equivalent to I
 3. A is a product of elementary matrices
-

Let P and Q be logical statements

If P then Q ($P \implies Q$)

1. Assume P is TRUE, and then show it logically implies that Q is TRUE
 2. Proof by contradiction: $\sim Q \implies \sim P$
-

If and only if (Equivalence)

P if and only if ($P \leftrightarrow Q$)

- $P \implies Q$
 - $Q \implies P$
-

Proof: $a \implies b, b \implies c, c \implies a$

$a \implies b, b \implies c, c \implies a$

Then, $a \leftrightarrow b$

- $a \implies b$

A is invertible $\implies A$ is row equivalent to I

$P \implies Q$

$\sim Q$: If A is not row equivalent to I , then $A \sim B$ such that B^{-1} does not exist

so, $B = E_n \dots E_1 A$

$$(E_n \dots E_1)^{-1} B = (E_n \dots E_1)^{-1} E_n \dots E_1 A = A$$

Due to $(A_1 A_2)^{-1} = A_2^{-1} A_1^{-1}$

$$A^{-1} = B^{-1} (E_n \dots E_1)$$

So, A is not invertible because B^{-1} does not exist

- $b \implies c$

If A is row equivalent to I then A is a product of elementary matrices

$P \implies Q$

$$E_n \dots E_1 A = I$$

$$(E_n \dots E_1)^{-1} (E_n \dots E_1) A = (E_n \dots E_1)^{-1} I$$

$$\text{so, } A = (E_n \dots E_1)^{-1} = E_1^{-1} \dots E_n^{-1}$$

For an elementary matrix E_i , E_i^{-1} is also an elementary matrix

- $c \implies a$

If A is a product of elementary matrices then A is invertible

$$A = (E_1 \dots E_n)$$

$$A^{-1} = (E_1 \dots E_n)^{-1} = E_n^{-1} \dots E_1^{-1} \text{ because } E_i^{-1} \text{ exists}$$

Therefore, $a \implies b, b \implies c, c \implies a$

Thm:

Let V be a vector space over F and $S = \{\alpha_1, \dots, \alpha_n\} \subset V$. Suppose S is a linearly independent set, then for every $w \in V$ there exist at most one representation as a linear combination of vectors in S .

Sketch:

If $S = \{\alpha_1, \dots, \alpha_n\} \subset V$ (linearly independent set), then $\forall w \in V$, there exist at least one representation:

$$w = \sum_{i=1}^n a_i \alpha_i$$

$P \implies Q$

Proof:

$\sim Q$: Assume that $\exists w \in V$, we have two possible representations

$$w = \sum_{i=1}^n a_i \alpha_i \text{ and } w = \sum_{i=1}^n b_i \alpha_i, \exists k : a_k \neq b_k, 1 \leq k \leq n$$

$$\text{So, } \mathbf{0} = w - w = \sum_{i=1}^n a_i \alpha_i - \sum_{i=1}^n b_i \alpha_i$$

$$= \sum_{i=1}^n (a_i - b_i) \alpha_i$$

$$(0) = \sum_{i=1, i \neq k}^n (a_i - b_i) \alpha_i + (a_k - b_k) \alpha_k, \text{ where } (a_k - b_k) \neq 0$$

Therefore, S is linearly dependent set

$S = \{\alpha_1, \dots, \alpha_n\} \subset V$ (vector space over field F)

If S spans V then $\forall w \in V, \exists a_i$'s $\in F : w = \sum_{i=1}^n a_i \alpha_i$

$$P \implies Q$$

Thm:

Let $S = \{\alpha_1, \dots, \alpha_n\} \subset V$ where V is a vector space over field F

If:

1. S is linearly independent (number of representations ≤ 1)
2. S spans V (number of representations ≥ 1)

then every vector $w \in V$ has a unique representation as a linear combination of vectors in S

Properties

$S = \{\alpha_1, \dots, \alpha_n\} \subset V$ (vector space over field F)

1. If S is linearly dependent, then any larger set of vectors containing S is linearly dependent
 2. If S is linearly independent, then any subset of S is linearly dependent
-

Basis and Dimension

Vector space V over field F

Basis of V is a set of vectors $S \in V$ such that:

1. S span V
2. S is a linearly independent set

Dimension of V is the number of vectors in the basis of V

Example:

$$V \equiv \mathbb{R}^3$$

$$V = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}, x_1, x_2, x_3 \in \mathbb{R}$$

$$\text{Basis: } \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$$

$$x_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\} \checkmark$$

$$\text{Dim}(\mathbb{R}^3) = 3$$

$$P_n(t) = \text{Polynomial of order } \leq n$$

$$p(t) \in P_n(t)$$

$$p(t) = c_0 + c_1 t^1 + \dots + c_n t^n$$

$$\text{Basis} = \{1, t, t^2, \dots, t^n\}$$

$$p(t) = \sum_{i=0}^n c_i t^i$$

$$\text{Dim}(P_n(t)) = n + 1$$

Thm: Dimension Theorem

All basis of a vector space have the same number of vectors

Proof:

If $T = \{\alpha_1, \dots, \alpha_n\}$ (a basis) and $S = \{\beta_1, \dots, \beta_m\}$ (a basis) then $n = m$

$P \implies Q$

Proof by contradiction:

$\sim Q : n \neq m \rightarrow (n < m) \text{ or } (n > m)$

Let $(n < m)$ - (Without Loss Of Generality)

$T = \{\alpha_1, \dots, \alpha_n\}$

$S = \{\beta_1, \dots, \beta_n, \beta_{n+1}, \dots, \beta_m\}$

$A = \{\alpha_1, \dots, \alpha_n\}, B = \{\beta_1, \dots, \beta_n\}$

$$B = \begin{bmatrix} \dots & \beta_1 & \dots \\ \dots & \beta_2 & \dots \\ \dots & \dots & \dots \\ \dots & \beta_n & \dots \end{bmatrix} \in \mathbb{R}^{n \times p}, C = \begin{bmatrix} c_{11} & \dots & c_{1n} \\ \dots & \dots & \dots \\ \dots & \dots & \dots \\ c_{1n} & \dots & c_{nn} \end{bmatrix} \in \mathbb{R}^{n \times n}, A = \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \alpha_2 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_n & \dots \end{bmatrix} \in \mathbb{R}^{n \times p}$$

$B = CA$

$$\begin{bmatrix} \dots & \beta_1 & \dots \\ \dots & \beta_2 & \dots \\ \dots & \dots & \dots \\ \dots & \beta_n & \dots \end{bmatrix} = \begin{bmatrix} c_{11} & \dots & c_{1n} \\ \dots & \dots & \dots \\ \dots & \dots & \dots \\ c_{1n} & \dots & c_{nn} \end{bmatrix} \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \alpha_2 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_n & \dots \end{bmatrix}$$

$\beta_{lj} = \sum_{k=1}^n c_{lk} \alpha_{kj}$ - Matrix Multiplication

$\beta_l = \sum_{k=1}^n c_{lk} \alpha_k$

Lemma 1:

If A and B have linearly independent rows then C is invertible

$P \implies Q$ or $\sim Q \implies \sim P$

Note: C is invertible $\leftrightarrow C$ has linearly independent rows

$\sim Q : C$ has linearly dependent rows

$$c_l = \sum_{i=1, i \neq l}^n a_i c_i, c_{lk} = \sum_{i=1, i \neq l}^n a_i c_{lk}$$

$$\beta_l = \sum_{k=1}^n c_{lk} \alpha_k$$

$$= \sum_{k=1}^n \sum_{i=1, i \neq l}^n a_i c_{lk} \alpha_k$$

$$= \sum_{i=1, i \neq l}^n a_i \sum_{k=1}^n c_{lk} \alpha_k$$

$$= \sum_{i=1, i \neq l}^n a_i \beta_i$$

So for $B = CA$ with invertible C then,

$$A = C^{-1}B$$

$$C^{-1} = D \equiv [d_{ij}]$$

$$\alpha_{ij} = \sum_{k=1}^n d_{ik} \beta_{kj}$$

$$\alpha_i = \sum_{k=1}^n d_{ik} \beta_k$$

$T = \{\alpha_1, \dots, \alpha_n\}$ is a basis of V and $\beta_{m+1} \in V$

$\beta_{n+1} = \sum_{i=1}^n e_i \alpha_i$ for some e_i 's $\in F$

$$\beta_{n+1} = \sum_{i=1}^n e_i (\sum_{k=1}^n d_{ik} \beta_k)$$

$$= \sum_{i=1}^n \sum_{k=1}^n e_i d_{ik} \beta_k$$

$$= \sum_{i=1}^n (\sum_{k=1}^n e_i d_{ik}) \beta_k$$

S is linearly dependent set

so S is a basis

Fundamental subspace of a matrix

$$A \in \mathbb{R}^{m \times n}$$

$$A = \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \dots & \dots & \dots \\ a_{m1} & \dots & a_{mn} \end{bmatrix}$$

$$Ax = b \text{ where } x \in \mathbb{R}^{n \times 1}, b \in \mathbb{R}^{m \times 1}$$

$$T : \mathbb{R}^{n \times 1} \text{ (Domain)} \rightarrow \mathbb{R}^{m \times 1} \text{ (Co-domain)}$$

$$A^T y = d \text{ where } A^T \in \mathbb{R}^{n \times m}, y \in \mathbb{R}^{m \times 1}, d \in \mathbb{R}^{n \times 1}$$

$$T : \mathbb{R}^{m \times 1} \text{ (Domain)} \rightarrow \mathbb{R}^{n \times 1} \text{ (Co-domain)}$$

1. Column Space: $C(A)$

$$C(A) = \{b \in \mathbb{R}^{m \times 1} : Ax = b, x \in \mathbb{R}^{n \times 1}\}$$

2. Row Space: $C(A^T)$

$$C(A^T) = \{d \in \mathbb{R}^{n \times 1} : A^T y = d, y \in \mathbb{R}^{m \times 1}\}$$

3. Null Space: $n(A)$

$$n(A) = \{x \in \mathbb{R}^{n \times 1} : Ax = \mathbf{0}\}$$

4. Left Null Space: $n(A^T)$

$$n(A) = \{y \in \mathbb{R}^{m \times 1} : A^T y = \mathbf{0}\}$$

| Subspaces | Dimension |
|-----------|-------------------------------|
| Domain | $n \equiv \text{order}$ |
| $C(A)$ | $r \equiv \text{rank}$ |
| $n(A)$ | $\zeta \equiv \text{nullity}$ |

Fact: $n = r + \zeta$

$r \equiv \text{rank}$

$r = \text{number of pivots} = \text{Dim}(C(A)) = \text{Dim}(C(A^T))$

$\zeta = \text{number of free variables} = \text{Dim}(n(A))$

Example:

$$A = \begin{bmatrix} 1 & 0 \\ 5 & 4 \\ 2 & 4 \end{bmatrix} \text{ Find } n(A) \text{ and its dimension.}$$

$$Ax = \mathbf{0}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 5 & 4 & 0 \\ 2 & 4 & 0 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2 - 5R1$$

$$R3 : R3 - 2R1$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 4 & 0 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2$$

$$R3 : R3 - R2$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 \\ 0 & 4 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$r = \text{number of pivots} = 2$$

$$\zeta = \text{number of free variables} = 0$$

$$x = 0, y = 0$$

$$n(A) = \{\mathbf{0}\}$$

$$\text{Dim}(n(A)) = \zeta = 0$$

$$A = \begin{bmatrix} 1 & 0 & 1 \\ 5 & 4 & 9 \\ 2 & 4 & 6 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2 - 5R1$$

$$R3 : R3 - 2R1$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 4 & 4 \\ 0 & 4 & 4 \end{bmatrix}$$

$$R1 : R1$$

$$R2 : R2$$

$$R3 : R3 - R2$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 4 & 4 \\ 0 & 0 & 0 \end{bmatrix} r = 2, \zeta = 1$$

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 4 & 4 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

$$\text{Let } x_3 = z, z \in \mathbb{R}$$

$$4x_3 + 4x_3 = 0 \implies x_2 = -Z$$

$$x_1 + x_3 = 0 \implies x_1 = -z$$

$$z \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix}, z \in \mathbb{R}$$

$$n(A) = \text{span}\left\{ \begin{bmatrix} -1 \\ -1 \\ 1 \end{bmatrix} \right\}$$

$$\text{Dim}(C(A)) = 2, \text{Dim}(n(A)) = 1$$

Thm:

Interchanging the rows of a matrix leaves its rank unchanged.

Thm:

If $Ax = 0$ and $Bx = 0$ have the same solution, then A and B have the same column rank

Week 4 Session 1

Outlines

Dimension Theorem

Existence and Uniqueness

Inner Product Space

$$Ax = b \text{ where } A \in \mathbb{R}^{m \times n}, x \in \mathbb{R}^{n \times 1} = y \in \mathbb{R}^{m \times 1}, m \neq n$$

Principal Component Analysis (Dimension Reduction)

$$X \in \mathbb{R}^{n \times p} \text{ where } n \text{ is the number of sample, } p \text{ is the number of features, } p \gg 1$$

$$A \in \mathbb{R}^{p \times s} \text{ where } s \text{ is a very small dimension}$$

$$X \rightarrow \tilde{X} \text{ mean} = 0 \implies K_{xx} \text{ where it is } p \times p \rightarrow \text{Eigenvalues}$$

$$E \in \mathbb{R}^{p \times p} = \begin{bmatrix} \dots & \dots & \dots & \dots & \dots \\ e_1 & \dots & e_s & \dots & e_p \\ \dots & \dots & \dots & \dots & \dots \\ (\lambda_1) & \dots & (\lambda_s) & \dots & (\lambda_p) \end{bmatrix}$$

$$\lambda_1 \gg \lambda_2 \gg \dots \gg \lambda_p$$

$$XE \text{ where } X \text{ is } n \times p, \text{ and } E \text{ is } p \times p$$

$$X\bar{E} = \hat{X} \text{ where } X \text{ is } n \times p, \bar{E} \text{ is } p \times s, \hat{X} \text{ is } n \times s$$

Thm:

If $Ax = 0$ and $Bx = 0$ have the same solution, then A and B have the same column rank.

Proof:

$$P \implies Q$$

Let s be the column rank of A

Let t be the column rank of B

where $t \neq s$ so, $(t > s)/(s > t)$

Let $t > s$ (WLOG)

$$B \in \mathbb{R}^{m \times n} = \begin{bmatrix} \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \beta_1 & \dots & \beta_s & \dots & \beta_t & \dots & \beta_n \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix} \text{ so that } Bx = 0$$

where column $1 \dots t$ are linearly independent, $t + 1 \dots n$ are linearly dependent

$$A \in \mathbb{R}^{m \times n} = \begin{bmatrix} \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \alpha_1 & \dots & \alpha_s & \dots & \alpha_t & \dots & \alpha_n \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \end{bmatrix} \text{ so that } Ax = 0$$

Therefore $\exists d'_i s \neq 0 : \sum_{i=1}^{\sigma} d_i \alpha_i = \mathbf{0}$ where $t > s$

$$\sum_{j=1}^{\sigma} d_j \alpha_j + \sum_{j=t+1}^n 0 \alpha_j = \mathbf{0}$$

$x_1 = d_1, x_2 = d_2, \dots, x_t = d_t$, this is the solution to $Ax = 0$

$$x_{t+1} = \dots = x_n = 0$$

$$\exists d'_i s \neq 0 : \sum_{j=1}^t d_i \beta_j + \sum_{j=t+1}^n 0 \beta_j = \mathbf{0}$$

$$\exists d'_i s \neq 0 : \sum_{j=1}^t d_i \beta_j = \mathbf{0}$$

$\{\beta_1, \dots, \beta_t\}$ is linearly dependent

Contradiction

Thm:

Elementary row operations preserve column rank

$$Ax = b \text{ elementary operations } \implies A'x = b'$$

$$Ax = 0 \implies A'x = 0$$

Thm:

Rank Theorem

Dimension of column space equals the dimension of row space.

$$Ax = b \text{ where } A \in \mathbb{R}^{m \times n}$$

Proof:

Let c be the column rank of A

Let r be the row rank of A

$$c \leq r \text{ or } r \leq c$$

Case 1: $c \leq r$

$$A = \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_r & \dots \\ \dots & \alpha_{r+1} & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_m & \dots \end{bmatrix}$$

$$\text{where } B \in \mathbb{R}^{r \times n} = \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_r & \dots \end{bmatrix} \text{ is linearly independent rows}$$

where $D \in \mathbb{R}^{(m-r) \times n} = \begin{bmatrix} \dots & \alpha_{r+1} & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_m & \dots \end{bmatrix}$ is linearly dependent rows

$\forall_j : r+1 \leq j \leq m, \exists t'_{ji} s :$

$$\alpha_j = \sum_{i=1}^r t_{ji} \alpha_i, \quad T \equiv [t_{ji}]$$

$$D = TB$$

$$(m-r) \times n = (m-r) \times r(t \times n)$$

$$A = \begin{bmatrix} B \\ D \end{bmatrix} = \begin{bmatrix} B \\ TB \end{bmatrix}$$

$$\text{If } Ax = \mathbf{0} \implies \begin{bmatrix} B \\ TB \end{bmatrix} x = \begin{bmatrix} Bx \\ TBx \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}$$

$$\text{If } Bx = \mathbf{0} \implies Ax = \begin{bmatrix} B \\ TB \end{bmatrix} x = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}$$

Case 2: $r \leq c$

$$A^T y = x$$

$$A^T \in \mathbb{R}^{n \times m}, y \in \mathbb{R}^{m \times 1}, x \in \mathbb{R}^{n \times 1}$$

The column rank of A^T is r

The row rank of A^T is c

$$A^T = \begin{bmatrix} \dots & \beta_1 & \dots \\ \dots & \dots & \dots \\ \dots & \beta_c & \dots \\ \dots & \beta_{c+1} & \dots \\ \dots & \dots & \dots \\ \dots & \beta_n & \dots \end{bmatrix}$$

where $E \in \mathbb{R}^{c \times m} = \begin{bmatrix} \dots & \beta_1 & \dots \\ \dots & \dots & \dots \\ \dots & \beta_c & \dots \end{bmatrix}$ is linearly independent rows

where $F \in \mathbb{R}^{(n-c) \times m} = \begin{bmatrix} \dots & \beta_{c+1} & \dots \\ \dots & \dots & \dots \\ \dots & \beta_n & \dots \end{bmatrix}$ is linearly dependent rows

$\forall_j : c+1 \leq j \leq n, \exists r'_{ji} s :$

$$\beta_j = \sum_{i=1}^c r_{ji} \beta_i, \quad R \equiv [r_{ji}]$$

$$F = RE$$

$$A^T = \begin{bmatrix} E \\ F \end{bmatrix} = \begin{bmatrix} E \\ RE \end{bmatrix}$$

$$A^T y = \mathbf{0} \text{ if and only if } Ey = \mathbf{0}$$

The column rank of $E = r$

$$Ey = x \text{ where } E \in \mathbb{R}^{c \times m}, y \in \mathbb{R}^{m \times 1}, x \in \mathbb{R}^{c \times 1}$$

$$r \leq c$$

Therefore $r = c$

$$Bx = y \text{ where } B \in \mathbb{R}^{m \times n}, x \in \mathbb{R}^{n \times 1}, y \in \mathbb{R}^{m \times 1}$$

x is the domain and y is the co-domain

$Bx \equiv$ column space or range space

column space \subset co-domain

Counting Theorem

$$A \in \mathbb{R}^{m \times n}$$

Dimension of column space + Dimension of null sapce = n = number of columns

Proof:

$$A \in \mathbb{R}^{m \times n}$$

$$A \implies R_r \text{ (row reduced Echelon)}$$

$$Ax = 0 \text{ and } R_r x = 0$$

Number of pivots in R_r = column rank (A)

$$R_r x = 0 \text{ and } Ax = 0$$

Dim of null sapce for $R_r = n - r$ where n is the number of columns and r is the number of pivots

Because A and R_r are row equivalent, then

$$Ax = 0 \text{ if and only if } R_r x = \mathbf{0}$$

Dimension of null space of $A = n - r$

$$n - r + r = n$$

$$\text{Dim}(n(A)) + \text{Dim}(C(A)) = \text{number of columns}$$

Thm:

Fundamental Theorem: $A \in \mathbb{R}^{m \times n}$

1. The row space of A and nullspace of A are orthogonal complements in $\mathbb{R}^{n \times 1}$
2. The column space of A and left null sapce of A are orthoginal complements in $\mathbb{R}^{m \times 1}$

Let v be a vector space

u be a subspace of v

w be a subsapce of v

u and w are orthogonal complements means that $\forall \alpha \in u$ and $\forall \beta \in w, \alpha \perp \beta$

$$\alpha \cdot \beta = 0$$

Proof:

$$Ax = y \text{ where } A \in \mathbb{R}^{m \times n}, x \in \mathbb{R}^{n \times 1}, y \in \mathbb{R}^{m \times 1}$$

1. Row Space: $C(A^T) = \{x \in \mathbb{R}^{n \times 1} : A^T y = x, y \in \mathbb{R}^{m \times 1}\}$
Null Space: $n(A) = \{x \in \mathbb{R}^{n \times 1} : Ax = \mathbf{0}\}$
Assume $\alpha \in C(A^T)$ and $\beta \in n(A)$
 $\alpha \cdot \beta = \alpha^T \beta = (A^T y)^T x$
 $= y^T Ax$ where $Ax = 0$
 $= 0$
2. Column Space: $C(A) = \{y \in \mathbb{R}^{m \times 1} : Ax = y, x \in \mathbb{R}^{n \times 1}\}$
Left null space $n(A^T) = \{y \in \mathbb{R}^{m \times 1} : A^T y = 0\}$
Assume $\alpha \in C(A)$ and $\beta \in n(A^T)$

$$\begin{aligned}\alpha^T \beta &= (Ax)^T y \\ &= x^T A^T y \text{ where } A^T y = 0 \\ &= 0\end{aligned}$$

Summary:

$$A \in \mathbb{R}^{m \times n}$$

column rank = r

dimension of null space = $n - r$

row rank = r

dimension of left null space = $m - r$

$$Ax = b$$

$m \equiv$ number of equations

$n \equiv$ number of unknowns

$$M = [Ab]$$

$$\text{Rank}(M) \quad \text{Rank}(A)$$

Existence and Uniqueness

Thm:

Let $Ax = b$ be a system with n -unknowns m equations and augmented matrix $M = [Ab]$

1. The system has at least one solution if and only if $\text{rank}(M) = \text{rank}(A)$

$$M' = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 3 & 1 & 2 \\ 0 & 0 & 2 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix} \quad A' = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 3 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

$$n = 3$$

2. The system has a unique solution if and only if $\text{rank}(M) = n = \text{rank}(A)$

Inner Product Space

Vector space V over a field F

Real Inner Product space:

Let V be a vector space over field \mathbb{R}

$\langle \alpha, \beta \rangle$ assign a real number for $\alpha, \beta \in V$

Then $\langle \alpha, \beta \rangle$ is an inner product if:

[I₁] Linearity: $\langle \alpha, a\beta + b\gamma \rangle = a \langle \alpha, \beta \rangle + b \langle \alpha, \gamma \rangle, \forall \alpha, \beta, \gamma \in V$ and $a, b \in \mathbb{R}$

[I₂] Symmetry: $\langle \alpha, \beta \rangle = \langle \beta, \alpha \rangle, \forall \alpha, \beta \in V$

[I₃] Positive Definite: $\langle \alpha, \alpha \rangle \geq 0$ and $\langle \alpha, \alpha \rangle = 0$ if and only if $\alpha = \mathbf{0}$

Examples:

1. Euclidean \mathbb{R}^n

$$\langle u, v \rangle = u \cdot v = \sum_{i=1}^n u_i v_i$$

2. Function space $C[a, b]$ and polynomial space $P_n(t)$

$C[a, b]$ - vector space of all continuous functions on the closed interval $[a, b]$

$$\langle f, g \rangle = \int_a^b f(x)g(x)dx$$

3. Matrix space $M = \mathbb{R}^{m \times n}$

M - vector space of all real $m \times n$ matrices

$$\langle A, B \rangle = \text{Tr}(B^T A)$$

Week 4 Session 2

Outlines

Orthogonality and Inner Products

Gram-Schmidt Process

Inner Product

$$\langle \alpha, \beta \rangle$$

Complex Inner Product Space

Vector V over field \mathbb{C}

$$\langle \alpha, \beta \rangle = \sum_{i=1}^n a_i b_i^* \text{ where } \alpha = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}, \beta = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$$

$\langle \alpha, \beta \rangle$ must satisfy the following properties:

$$\forall \alpha, \beta, \gamma \in V; \forall a, b \in \mathbb{C}$$

$[I_1]$: Linearity

$$\langle \alpha, a\beta + b\gamma \rangle = a^* \langle \alpha, \beta \rangle + b^* \langle \alpha, \gamma \rangle$$

$[I_2]$: Conjugate Symmetry

$$\langle \alpha, \beta \rangle = \langle \beta, \alpha \rangle^*$$

$[I_3]$: Positive Definite:

$$\langle \alpha, \alpha \rangle \geq 0 \text{ and } \langle \alpha, \alpha \rangle = 0 \text{ if and only if } \alpha = \mathbf{0}$$

Normed Vector Spaces

Let $V = \{\alpha, \beta, \gamma, \dots\}$ be a vector space over a field F . A norm $\|\cdot\|$ of V is a function from the elements of V (vectors in V) into the non-negative real number such that:

$$[N_1]: \|\alpha\| \geq 0, \forall \alpha \in V \text{ and } \|\alpha\| = 0 \text{ if and only if } \alpha = \mathbf{0}$$

$$[N_2]: \|k\alpha\| = |k| \|\alpha\|, \forall \alpha \in V \text{ and } \forall k \in F$$

$$[N_3]: \|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|, \forall \alpha, \beta \in V \text{ (triangle inequality)}$$

Example:

$$1. v = \mathbb{R}^n, \alpha \in V, \alpha = [a_1, \dots, a_n]$$

$$\|\alpha\| = \sqrt{(a_1)^2 + \dots + (a_n)^2} \text{ - Euclidean Norm}$$

$$2. v = \mathbb{C}^n \text{ Complex field}$$

Metric Space

Vector space V over F

$M(\alpha, \beta)$ - metric

Properties of a metric:

$$[M_1]: M(\alpha, \beta) \geq 0 \text{ and } M(\alpha, \beta) = 0 \text{ if and only if } \alpha = \beta$$

$$[M_2]: M(\alpha, \beta) = M(\beta, \alpha)$$

$$[M_3]: M(\alpha, \gamma) \leq M(\alpha, \beta) + M(\beta, \gamma)$$

Norm

$$l^p \text{ - norm: } \sqrt[p]{\sum_{i=1}^n |x_i|^p} = \|x\|_p$$

$$l^p \text{ - distance: } \|x - y\|_p$$

Volume of an Euclidean ball of radius r


$$l^2 \text{ - norm } r = \sqrt{\sum_{i=1}^n x_i^2} = n = 2$$

$$r = \sqrt{x^2 + y^2}$$

Given Conditions:

$$V_n(r) = c_n r^n; c_n = \frac{2\pi}{n} c_{n-2}$$

| n | V_n |
|--|-----------------------|
| $c_1 = 2$ | $2r$ |
| $c_2 = \pi$ | πr^2 |
| $c_3 = \frac{2\pi}{3} c_1 = \frac{4\pi}{3}$ | $\frac{4\pi}{3} r^3$ |
| $c_4 = \frac{2\pi}{4} c_2 = \frac{\pi^2}{2}$ | $\frac{\pi^2}{2} r^4$ |



$$0 < \epsilon < r$$

Volume shell - Entire Volume

$$\frac{c_n r^n - c_n (r - \epsilon)^n}{c_n r^n}$$

$$= \frac{r^n - (r - \epsilon)^n}{r^n}$$

$$= 1 - \left(1 - \frac{\epsilon}{r}\right)^n$$

$$0 < \epsilon < r \implies 0 < \frac{\epsilon}{r} < 1$$

$$1 > 1 - \frac{\epsilon}{r} > 0$$

$$\lim_{n \rightarrow \infty} 1 - (1 - \frac{\epsilon}{r})^n = 1$$

Orthogonality

Vector space V over field F

$$\alpha, \beta \in V$$

$$\alpha \perp \beta \text{ if and only if } \langle \alpha, \beta \rangle = 0$$

Def: Let $S = \{\alpha_1, \dots, \alpha_n\} \subset V$ is mutually orthogonal if and only if

$$\alpha_i \cdot \alpha_j = 0 \text{ for } i \neq j$$

Mutually Orthonormal

A vector is normal if and only if its norm $\|\cdot\|$ is equal to 1

Def: Let $S = \{\beta_1, \dots, \beta_n\} \subset V$ is mutually orthonormal if and only if

$$\alpha_i \cdot \alpha_j = \begin{cases} 0, & \text{if } i \neq j \\ 1, & \text{if } i = j \end{cases}$$

$$S = \{\alpha_1, \dots, \alpha_n\} \text{ which is mutually orthogonal} \implies T = \left\{ \frac{\alpha_1}{\|\alpha_1\|}, \dots, \frac{\alpha_n}{\|\alpha_n\|} \right\} \text{ which is mutually orthonormal}$$

S is linearly independent $\not\Rightarrow S$ is mutually orthogonal

S is mutually orthogonal $\implies S$ is linearly independent

Example.

$$S = \{\alpha_1, \alpha_2, \alpha_3\}$$

$$\text{where } \alpha_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \alpha_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}, \alpha_3 = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$$

$$v = \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}$$

$$\text{where } \alpha_1 \cdot \alpha_2 = 0, \alpha_1 \cdot \alpha_3 = 0, \alpha_2 \cdot \alpha_3 = 0$$

$$c_1 = \frac{v \cdot \alpha_1}{\alpha_1 \cdot \alpha_1} = \frac{3}{1}$$

$$c_2 = \frac{v \cdot \alpha_2}{\alpha_2 \cdot \alpha_2} = \frac{7}{2}$$

$$c_3 = \frac{v \cdot \alpha_3}{\alpha_3 \cdot \alpha_3} = \frac{3}{2}$$

$$\text{Therefore: } v = 3\alpha_1 + \frac{7}{2}\alpha_2 + \frac{3}{2}\alpha_3$$

Thm:

If $S = \{\alpha_1, \dots, \alpha_n\}$ is in a vector space V and S is mutually orthogonal (with $\alpha_i \neq 0$), then S is linearly independent

Proof:

$$c_1\alpha_1 + \dots + c_n\alpha_n = \mathbf{0}$$

$$(c_1\alpha_1 + \dots + c_n\alpha_n) \cdot \alpha_i = \mathbf{0} \cdot \alpha_i = \mathbf{0}$$

$$\sum_{j=1}^n c_j(\alpha_j \cdot \alpha_i) = 0$$

$$\sum_{j=1, j \neq i}^n c_j(\alpha_j \cdot \alpha_i) + c_i(\alpha_i \cdot \alpha_i) = 0$$

$\sum_{j=1, j \neq i}^n c_j (\alpha_j \cdot \alpha_i) = 0$ because S is mutually orthogonal

$$c_i (\alpha_i \cdot \alpha_i) = 0$$

$$c_i = \frac{0}{\alpha_i \cdot \alpha_i} \text{ where } \alpha_i \neq 0$$

Therefore, $c_1 = c_2 = \dots = c_n$ is the only solution

Therefore, S is linearly independent

$$S = \{\alpha_1, \alpha_2, \alpha_3\}$$

$$\text{where } \alpha_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \alpha_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \alpha_3 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

It is linearly independent but not mutually orthogonal

$$\text{where } \alpha_1 \cdot \alpha_2 = 1, \alpha_1 \cdot \alpha_3 = 2, \alpha_2 \cdot \alpha_3 = 1$$

Thm:

If $S = \{\alpha_1, \dots, \alpha_n\}$ is in a vector space V , S is a basis of V and S is mutually orthogonal, then $\forall \beta \in V, \exists a'_i$ such that

$$a_1 \alpha_1 + \dots + a_n \alpha_n = \beta$$

$$a_i = \frac{\beta \cdot \alpha_i}{\alpha_i \cdot \alpha_i}$$

Proof:

S is a basis for V

$$\forall \beta \in V$$

$$a_1 \alpha_1 + \dots + a_n \alpha_n = \beta$$

$$(a_1 \alpha_1 + \dots + a_n \alpha_n) \alpha_i = \beta \cdot \alpha_i$$

$$\sum_{j=1}^n a_j (\alpha_j \cdot \alpha_i) = \beta \cdot \alpha_i$$

$$\sum_{j=1, j \neq i}^n a_j (\alpha_j \cdot \alpha_i) + a_i (\alpha_i \cdot \alpha_i) = \beta \cdot \alpha_i$$

$$a_i (\alpha_i \cdot \alpha_i) = \beta \cdot \alpha_i$$


$$\text{Therefore, } a_i = \frac{\beta \cdot \alpha_i}{\alpha_i \cdot \alpha_i}$$

$S = \{\alpha_1, \dots, \alpha_n\}$ is an orthogonal basis of V

1. Basis
 2. Mutually orthogonal
-

Projection

Projection of α onto β



$$\gamma = \text{Proj}_{\beta}(\alpha) = c\beta$$

$$\omega = \alpha - \gamma \text{ and } \omega \perp \beta$$

$$(\alpha - \gamma) \perp \beta$$

$$(\alpha - c\beta) \perp \beta$$

$$(\alpha - c\beta) \cdot \beta = 0$$

$$\alpha \cdot \beta - c\beta \cdot \beta = 0$$

$$c = \frac{\alpha \cdot \beta}{\beta \cdot \beta}$$

$$\text{Proj}_{\beta}(\alpha) = c\beta = \frac{\alpha \cdot \beta}{\beta \cdot \beta} \beta$$


$$\text{Orth}_{\beta}(\alpha) \equiv \alpha - \gamma = \alpha - \frac{\alpha \cdot \beta}{\beta \cdot \beta} \beta$$

$$\text{Proj}_{\beta}(\alpha) + \text{Orth}_{\beta}(\alpha) = \alpha$$

w is a subspace of V

Projection of α onto w

$S = \{\beta_1, \dots, \beta_m\}$ is an orthogonal basis for w



$$\gamma = \text{Proj}_w(\alpha) = c_1\beta_1 + \dots + c_m\beta_m$$

$$1. \gamma \cdot \beta_i = (c_1\beta_1 + \dots + c_m\beta_m) \cdot \beta_i \\ = c_i(\beta_i \cdot \beta_i) + \sum_{j=1, j \neq i}^m c_j(\beta_j \cdot \beta_i)$$

$$2. \omega = \alpha - \gamma : \omega \perp \beta_i$$

$$(\alpha - \gamma) \perp \beta_i$$

$$(\alpha - \gamma) \cdot \beta_i = 0$$

$$\alpha \cdot \beta_i = \gamma \cdot \beta_i$$

$$\text{Then } \gamma \cdot \beta = \alpha \cdot \beta_i = c_i(\beta_i \cdot \beta_i)$$

$$c_i = \frac{\alpha \cdot \beta_i}{\beta_i \cdot \beta_i}$$

$$\gamma = \frac{\alpha \cdot \beta_1}{\beta_1 \cdot \beta_1} + \dots + \frac{\alpha \cdot \beta_m}{\beta_m \cdot \beta_m} = \text{Proj}_\omega(\alpha)$$

$S = \{\alpha_1, \dots, \alpha_n\}$ which is linearly independent "Gram Schmidt" $\implies T = \{\beta_1, \dots, \beta_n\}$ which is mutually orthogonal


$$L(S) = L(T)$$

Def:

If V is a vector space and S is a subspace of V , then

$$\omega = \{\alpha + \beta : \alpha \in S, \beta \in S^\perp\} = V = S \oplus S^\perp$$

where S^\perp is the orthogonal complement of S



Gram-Schmidt Process

Given $S = \{\alpha_1, \dots, \alpha_n\}$ where S is linearly independent.

Find $T = \{\beta_1, \dots, \beta_n\}$ where S is mutually orthogonal and $L(s) = L(\tau)$

$$\beta_1 = \alpha_1$$


$$V_1 = \text{span}\{\alpha_1\} = \text{span}\{\beta_1\}$$

$$\beta_2 = \alpha_2 - \frac{\alpha_2 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 \text{ where } \text{Proj}_{V_1}(\alpha_2) = \frac{\alpha_2 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1$$

$$V_2 = \text{span}(\{\alpha_1, \alpha_2\}) = \text{span}(\{\beta_1, \beta_2\})$$

$$\beta_3 = \alpha_3 - \left[\frac{\alpha_3 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 + \frac{\alpha_3 \cdot \beta_2}{\beta_2 \cdot \beta_2} \beta_2 \right] \text{ where } \text{Proj}_{V_2}(\alpha_3) = \frac{\alpha_3 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 + \frac{\alpha_3 \cdot \beta_2}{\beta_2 \cdot \beta_2} \beta_2$$

$$\beta_k = \alpha_k - \left[\frac{\alpha_k \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 + \dots + \frac{\alpha_k \cdot \beta_{(k-1)}}{\beta_{(k-1)} \cdot \beta_{(k-1)}} \beta_{(k-1)} \right]$$



$$\beta_2 = \alpha_2 - \frac{\alpha_2 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1$$

$$\text{Ex. } \alpha_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \alpha_2 = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix}, \alpha_3 = \begin{bmatrix} 2 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \alpha_4 = \begin{bmatrix} 0 \\ 0 \\ 3 \\ 1 \end{bmatrix}$$

$$\beta_1 = \alpha_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}$$

$$\beta_2 = \alpha_2 - \frac{\alpha_2 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 = \begin{bmatrix} 1 \\ 2 \\ 0 \\ 1 \end{bmatrix} - \frac{4}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix}$$

$$\beta_3 = \alpha_3 - \left[\frac{\alpha_3 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 + \frac{\alpha_3 \cdot \beta_2}{\beta_2 \cdot \beta_2} \beta_2 \right] = \begin{bmatrix} 2 \\ 1 \\ 1 \\ 0 \end{bmatrix} - \frac{4}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{0}{2} \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}$$

$$\beta_4 = \alpha_4 - \left[\frac{\alpha_4 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 + \frac{\alpha_4 \cdot \beta_2}{\beta_2 \cdot \beta_2} \beta_2 + \frac{\alpha_4 \cdot \beta_3}{\beta_3 \cdot \beta_3} \beta_3 \right] = \begin{bmatrix} 0 \\ 0 \\ 3 \\ 1 \end{bmatrix} - \frac{4}{4} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{-3}{2} \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix} - \frac{-1}{2} \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} -1/2 \\ 1/2 \\ 1/2 \\ -1/2 \end{bmatrix}$$

Let V be a vector where

$$V = \left\{ \begin{bmatrix} a & 0 \\ 0 & a \end{bmatrix}, \text{ where } a \in \mathbb{R} \right\}$$

then $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right\}$ is a basis, and the dimension is 1

Week 4 Session 1 (Messed)

Outlines

- Review
- Dimension
- Rank Theorem
- Counting Theorem
- Fundamental Theorem

Review:

Theorem 1:

Interchanging rows of a matrix leaves its row rank unchanged.

Theorem 2:

If $Ax = 0$ and $Bx = 0$ have the same solution then A and B have the same column rank

[Proof: Lecture 6]

Theorem 3:

Elementary row operation does not change the column rank.

Reason: Elementary row operation preserves solution, then apply Theorem 2

$Ax = b$ where $A \in \mathbb{R}^{m \times n}$, $x \in \mathbb{R}^{n \times 1}$, $b \in \mathbb{R}^{m \times 1}$

Theorem 4: Rank Theorem:

Dimension of column space = Dimension of row space

Proof:

column rank = Dimension of column space

row rank = Dimension of row space

Let r = row rank of A

c = column rank of A

Claim 1: $c \leq r$

Proof:

$A = \begin{bmatrix} \dots & a_1 & \dots \\ \dots & \dots & \dots \\ \dots & a_r & \dots \\ \dots & a_{r+1} & \dots \\ \dots & \dots & \dots \\ \dots & a_m & \dots \end{bmatrix}$, where $\begin{bmatrix} \dots & a_1 & \dots \\ \dots & \dots & \dots \\ \dots & a_r & \dots \end{bmatrix}$ is linearly independent rows, $\begin{bmatrix} \dots & a_{r+1} & \dots \\ \dots & \dots & \dots \\ \dots & a_m & \dots \end{bmatrix}$ is linearly dependent rows

Let $B = \begin{bmatrix} \dots & a_1 & \dots \\ \dots & \dots & \dots \\ \dots & a_r & \dots \end{bmatrix}$, where $B \in \mathbb{R}^{r \times n}$, $D = \begin{bmatrix} \dots & a_{r+1} & \dots \\ \dots & \dots & \dots \\ \dots & a_m & \dots \end{bmatrix}$, where $D \in \mathbb{R}^{(m-r) \times n}$

Note: $\forall j : r+1 \leq j \leq m, \exists t_{ji}$'s such that

$a_j = \sum_{i=1}^r t_{ji} a_i$ - Linearly dependent rows

Let $T = [t_{ji}]$

$D = TB$

$$A = \begin{bmatrix} B \\ D \end{bmatrix} = \begin{bmatrix} B \\ TB \end{bmatrix}$$

$$\text{So, } Ax = \mathbf{0} \text{ means } \begin{bmatrix} B \\ TB \end{bmatrix} x = \begin{bmatrix} Bx \\ TBx \end{bmatrix} = \mathbf{0}$$

$$Ax = \mathbf{0} \text{ if and only if } Bx = \mathbf{0}$$

The column rank of $A = c$

so the column rank of $B = c$

Remember:

$$Bx = d \in \mathbb{R}^{r \times 1}$$

so the column space of $B \subset \mathbb{R}^{r \times 1}$

$$\text{Dim}(C(B)) \leq \text{Dim}(\mathbb{R}^{r \times 1}), \text{ where } C(B) \text{ is the column space}$$

$$c \leq r$$

Claim 2: $r \leq c$

Proof:

Definition of transpose

$$c = \text{row rank of } A^T$$

$$r = \text{column rank of } A^T$$

$$A = \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_c & \dots \\ \dots & \alpha_{c+1} & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_n & \dots \end{bmatrix}, \text{ where } \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_c & \dots \end{bmatrix} \text{ is linearly independent rows, } \begin{bmatrix} \dots & \alpha_{c+1} & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_n & \dots \end{bmatrix} \text{ is linearly}$$

dependent rows

$$\text{Let } E = \begin{bmatrix} \dots & \alpha_1 & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_c & \dots \end{bmatrix}, \text{ where } B \in \mathbb{R}^{c \times m}, F = \begin{bmatrix} \dots & \alpha_{r+1} & \dots \\ \dots & \dots & \dots \\ \dots & \alpha_n & \dots \end{bmatrix}, \text{ where } D \in \mathbb{R}^{(n-c) \times m}$$

Note: $\forall i : c + 1 \leq i \leq n, \exists r_{ij}$'s such that

$$\alpha_i = \sum_{j=1}^c r_{ij} \alpha_j$$

$$\text{Let } R = [r_{ij}]$$

$$\text{then } F = RE$$

$$A^T = \begin{bmatrix} E \\ F \end{bmatrix} = \begin{bmatrix} E \\ RE \end{bmatrix}$$

$$A^T y = \mathbf{0} \text{ means } \begin{bmatrix} E \\ RE \end{bmatrix} y = \begin{bmatrix} Ey \\ REy \end{bmatrix} = \mathbf{0}$$

$$A^T y = \mathbf{0} \text{ if and only if } Ey = \mathbf{0}$$

$$\text{The column rank of } A^T = r$$

$$\text{So the column rank of } E = r$$

Remember:

$$Ex = f \in \mathbb{R}^{c \times 1}$$

So the column space of $E \subset \mathbb{R}^{c \times 1}$

$$\text{Dim}(C(E)) \leq \text{Dim}(\mathbb{R}^{c \times 1}), \text{ where } C(E) \text{ is the column space of } E$$

Then $r \leq c$

Therefore $c = r$

Theorem 5: Counting Theorem

Dimension of column space + Dimension of null space = number of columns

Reason: Let R_r be the row-reduced echelon form of A

- Row space of A = row space of R_r
Because rows of R_r are linear combinations of rows of A and vice versa
- Column Space of A = column space of R_r
Because same solution for $Ax = 0$ and $R_r x = 0$
- Null space of A = null space of R_r
Because elementary row operation preserves solution

From R_r

n is the number of variables

r is the number of pivot variables

$n - r$ is the number of free variables

$$n = r + (n - r)$$

$$\text{number of columns} = \text{Dim}(C(R_r)) + \text{Dim}(n(R_r))$$

$$= \text{Dim}(C(A)) + \text{Dim}(n(A))$$

Similarly,

Dimension of row space + Dimension of left null space = Dimension of rows

Theorem 6: Fundamental Theorem

1. The row space and null space of A are orthogonal complements in \mathbb{R}^n
2. The column space and left null space of A are orthogonal complements in \mathbb{R}^m

Proof:

Definition: Let V be a vector space. Let U be a subspace of V and W be a subspace of V . u and w are orthogonal complements in V means that $\forall u \in U$ and $\forall w \in W, u \perp w (u \cdot w = 0)$

1. Row space: $C(A^T) = \{A^T y : y \in \mathbb{R}^{m \times 1}\}$
Null space: $n(A) = \{x \in \mathbb{R}^{n \times 1} : Ax = \mathbf{0}\}$
Let $A^T y_1 \in C(A^T)$ and $x_0 \in n(A)$
 $(x_0)^T A^T y_1 = ((x_0)^T A^T y_1)^T$ where it is a representation of transpose of scalar
 $= y_1 A x_0$ where $A x_0 = \mathbf{0}$
 $= \mathbf{0}$
2. Column space: $C(A) = \{Ax : x \in \mathbb{R}^{n \times 1}\}$
Left null space: $n(A^T) = \{z \in \mathbb{R}^{m \times 1} : A^T z = \mathbf{0}\}$
Let $A x_1 \in C(A)$ and $z_0 \in n(A^T)$
 $(z_0)^T A x_1 = ((z_0)^T A x_1)^T$

$$= (x_1)^T A^T z_0 \text{ where } A^T z_0 = \mathbf{0}$$

$$= \mathbf{0}$$

Summary

for $A \in \mathbb{R}^{m \times n}$

Column rank = Row rank = Rank = r

Dimension of the null space = $n - r$

Dimension of the left null space = $m - r$

Theorem 1: Existence and Uniqueness

Let $Ax = b$ be a system with n unknowns with augmented matrix $M = [A|b]$ then:

Existence

The system has at least one solution if and only if

$$\text{rank}(A) = \text{rank}(M)$$

Uniqueness

The system has a unique solution if and only if

$$\text{rank}(A) = \text{rank}(M) = n$$

Proof:

1. A has no solution if and only if there exist a degenerate row $[0, 0, \dots, 0|b]$ in the echelon form of M
 2. $\text{rank}(A) = n$ if and only if no free variable
-

Inner product and orthogonality

Real inner product space:

Let V be a vector space over \mathbb{R} . Suppose that $\forall \alpha, \beta \in V$ $\langle \alpha, \beta \rangle$ assigns a real number. Then $\langle \alpha, \beta \rangle$ is an inner product on V if

$$[I_1] \text{ Linearity: } \langle \alpha, a\beta + b\gamma \rangle = a \langle \alpha, \beta \rangle + b \langle \alpha, \gamma \rangle, \forall \alpha, \beta, \gamma \in V \text{ and } a, b \in \mathbb{R}$$

$$[I_2] \text{ Symmetry: } \langle \alpha, \beta \rangle = \langle \beta, \alpha \rangle, \forall \alpha, \beta \in V$$

$$[I_3] \text{ Positive Definite: } \langle \alpha, \alpha \rangle \geq 0 \text{ and } \langle \alpha, \alpha \rangle = 0 \text{ if and only if } \alpha = \mathbf{0}$$

Examples:

1. Euclidean \mathbb{R}^n

$$\langle u, v \rangle = u \cdot v = \sum_{i=1}^n u_i v_i$$

2. Function space $C[a, b]$ and polynomial space $P_n(t)$

$C[a, b]$ - vector space of all continuous functions on the closed interval $[a, b]$

$$\langle f, g \rangle = \int_a^b f(x)g(x)dx$$

3. Matrix space $M = \mathbb{R}^{m \times n}$

M - vector space of all real $m \times n$ matrices

$$\langle A, B \rangle = \text{Tr}(B^T A)$$

Complex Inner product Space

Vector space V : $\alpha, \beta, \gamma \in V$

The field is \mathbb{C} : $a, b \in \mathbb{C}$

$\langle u, v \rangle$ must satisfy the following:

$[I_1]$: Linearity

$$\langle \alpha, a\beta + b\gamma \rangle = a^* \langle \alpha, \beta \rangle + b^* \langle \alpha, \gamma \rangle$$

$[I_2]$: Conjugate Symmetry

$$\langle \alpha, \beta \rangle = \langle \beta, \alpha \rangle^*$$

$[I_3]$: Positive Definite:

$$\langle \alpha, \alpha \rangle \geq 0 \text{ and } \langle \alpha, \alpha \rangle = 0 \text{ if and only if } \alpha = \mathbf{0}$$

Normed Vector Spaces

Let $V = \{\alpha, \beta, \gamma, \dots\}$ be a vector space over a field F . A norm $\|\cdot\|$ of V is a function from the elements of V (vectors in V) into the non-negative real number such that:

$$[N_1]: \|\alpha\| \geq 0, \forall \alpha \in V \text{ and } \|\alpha\| = 0 \text{ if and only if } \alpha = \mathbf{0}$$

$$[N_2]: \|k\alpha\| = |k| \|\alpha\|, \forall \alpha \in V \text{ and } \forall k \in F$$

$$[N_3]: \|\alpha + \beta\| \leq \|\alpha\| + \|\beta\|, \forall \alpha, \beta \in V \text{ (triangle inequality)}$$

Example:

$$1. v = \mathbb{R}^n, \alpha \in V, \alpha = [a_1, \dots, a_n]$$

$$\|\alpha\| = \sqrt{(a_1)^2 + \dots + (a_n)^2} \text{ - Euclidean Norm}$$

$$2. v = \mathbb{C}^n \text{ Complex field}$$

$$\forall \alpha \in V, \|\alpha\| = \sqrt{(a_1)^2 + \dots + (a_n)^2}$$

Definition: A metric $M(\alpha, \beta)$ on pairs of elements $\alpha, \beta \in V$ satisfies the following:

$$[M_1]: M(\alpha, \beta) = 0 \text{ if and only if } \alpha = \beta$$

$$[M_2]: M(\alpha, \beta) = M(\beta, \alpha)$$

$$[M_3]: M(\alpha, \beta) + M(\beta, \gamma) \geq M(\alpha, \gamma), \forall \alpha, \beta, \gamma \in V$$

l^p - distance

$$l^p(x, y) = \sqrt[p]{\sum_{i=1}^n |x_i - y_i|^p}, 1 \leq p \leq \infty$$

$$P = 1$$

$$l^1(x, y) = \sum_{i=1}^n |x_i - y_i| \text{ - Absolute}$$

Let $x, y \in B^n = \{0, 1\}^n$

$$\text{consider } x = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, y = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}$$

Mid Term 1 Review

Sample 1

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}$$

Find A^{-1} if the inverse exists otherwise give sufficient reason.

Using Gaussian Jordan-Elimination

$$\begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 3 & 0 & 0 & 1 \end{bmatrix}$$

$$R_1 : R_1$$

$$R_2 : R_2 - 3R_1$$

$$R_3 : R_3 - 2R_1$$

$$\begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -4 & -8 & -3 & 1 & 0 \\ 0 & -3 & -3 & -2 & 0 & 1 \end{bmatrix}$$

$$R_3 : R_3 - 3/4R_2$$

$$\begin{bmatrix} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -4 & -8 & -3 & 1 & 0 \\ 0 & 0 & 1 & 1/12 & -1/4 & 1/3 \end{bmatrix}$$

$$R_1 : R_1 - 3R_3$$

$$R_2 : R_2 + 8R_3$$

$$\begin{bmatrix} 1 & 2 & 0 & 3/4 & 3/4 & -1 \\ 0 & -4 & 0 & -7/3 & -1 & 8/3 \\ 0 & 0 & 1 & 1/12 & -1/4 & 1/3 \end{bmatrix}$$

$$R_2 : -1/4R_2$$

$$\begin{bmatrix} 1 & 2 & 0 & 3/4 & 3/4 & -1 \\ 0 & 1 & 0 & 7/12 & 1/4 & -2/3 \\ 0 & 0 & 1 & 1/12 & -1/4 & 1/3 \end{bmatrix}$$

$$R_1 : R_1 - 2R_2$$

$$\begin{bmatrix} 1 & 0 & 0 & -5/12 & 1/4 & 1/3 \\ 0 & 1 & 0 & 7/12 & 1/4 & -2/3 \\ 0 & 0 & 1 & 1/12 & -1/4 & 1/3 \end{bmatrix}$$

Sample 2

Find the left null space of matrix A

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & -2 & -1 & 1 \\ -1 & -8 & -10 & -11 \end{bmatrix}$$

Find the dimension of $C(A)$ and $C(A^T)$

$$N(A^T) = \{y \in \mathbb{R}^{3 \times 1}, A^T y = 0\}$$

$$A^T = \begin{bmatrix} 1 & 2 & -1 \\ 2 & -2 & -8 \\ 3 & -1 & -10 \\ 4 & 1 & -11 \end{bmatrix}$$

$$R_2 : R_2 - 2R_1$$

$$R_3 : R_3 - 3R_1$$

$$R_4 : R_4 - 4R_1$$

$$\begin{bmatrix} 1 & 2 & -1 \\ 0 & -6 & -6 \\ 0 & -7 & -7 \\ 0 & -7 & -7 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 2 & -1 \\ 0 & -6 & -6 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$-6y_2 - 6y_3 = 0$$

$$y_2 = -y_3$$

$$y_1 + 2y_2 - y_3 = 0$$

$$y_1 = 3y_3$$

$$y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 3y_3 \\ -y_3 \\ y_3 \end{bmatrix} = y_3 \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}$$

$$\text{Basis}(n(A^T)) = \begin{bmatrix} 3 \\ -1 \\ 1 \end{bmatrix}$$

Null space of A ($Ax = 0, x \in \mathbb{R}^{4 \times 1}$)

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & -2 & -1 & 1 \\ -1 & -8 & -10 & -11 \end{bmatrix}$$

$$R_2 : R_2 - 2R_1$$

$$R_3 : R_3 + R_1$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & -6 & -7 & -7 \\ 0 & -6 & -7 & -7 \end{bmatrix}$$

$$R_3 : R_3 - R_2$$

$$\begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & -6 & -7 & -7 \\ 0 & 0 & 0 & 0 \end{bmatrix} \implies \text{dimension is 2}$$

According to the counting theorem

$$D(C(A)) + D(N(A)) = \text{number of columns} = 4$$

$$D(N(A)) = 2$$

$$D(C(A^T)) + D(N(A^T)) = \text{number of rows} = 3$$

$$D(N(A^T)) = 2$$

Sample 3

Use the Gram-Schmidt procedure to construct a set of orthonormal set of $\{[-1, 1, 0], [-1, 0, 1], [0, 1, 1]\}$

Express $v = [2 \ 3 \ 5]$ as linear combination of such orthonormal vectors.

$$\alpha_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \alpha_2 = \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}, \alpha_3 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

$$\beta_1 = \alpha_1 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

$$\|\beta_1\|^2 = 2$$

$$\beta_2 = \alpha_2 - \frac{\alpha_2 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1$$

$$= \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} -1/2 \\ -1/2 \\ 1 \end{bmatrix}$$

$$\|\beta_2\|^2 = 3/2$$

$$\beta_3 = \alpha_3 - \left[\frac{\alpha_3 \cdot \beta_1}{\beta_1 \cdot \beta_1} \beta_1 + \frac{\alpha_3 \cdot \beta_2}{\beta_2 \cdot \beta_2} \beta_2 \right]$$

$$= \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} - -\frac{1}{2} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} - \frac{1/2}{3/2} \begin{bmatrix} -1/2 \\ -1/2 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} 2/3 \\ 2/3 \\ 2/3 \end{bmatrix}$$

$$\|\beta_3\|^2 = 12/9$$

The orthonormal basis are:

$$\gamma_1 = \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$$

$$\gamma_2 = \sqrt{2/3} \begin{bmatrix} -1/2 \\ -1/2 \\ 1 \end{bmatrix} = \sqrt{1/6} \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix}$$

$$\gamma_3 = \sqrt{9/12} \begin{bmatrix} 2/3 \\ 2/3 \\ 2/3 \end{bmatrix} = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$v = [2 \quad 3 \quad 5]$$

$$v = (v \cdot \gamma_1)\gamma_1 + (v \cdot \gamma_2)\gamma_2 + (v \cdot \gamma_3)\gamma_3$$

$$v \cdot \gamma_1 = \frac{1}{\sqrt{2}}$$

$$v \cdot \gamma_2 = \frac{5}{\sqrt{6}}$$

$$v \cdot \gamma_3 = \frac{10}{\sqrt{3}}$$

$$v = \frac{1}{\sqrt{2}}\gamma_1 + \frac{5}{\sqrt{6}}\gamma_2 + \frac{10}{\sqrt{3}}\gamma_3$$

$$= 1/2 \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix} + 5/6 \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix} + 10/3 \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Week 5 Session 1 (Only 1)

Outlines

Linear Transformation (mappings)

Groups

Symmetric Group

Determinants

A is $n \times n$

Linear mapping

$$Ax = b$$

Non empty sets A and B

$$f : A \rightarrow B$$

f assigns a unique to $a \in A$ in B

$A \equiv \text{Domain of } f, B \equiv \text{Codomain}$

$$A' \subset A, f(A') = \{f(a) : a \in A'\}$$

$$B' \subset B, f^{-1}(B') = \{a \in A : f(a) = b, b \in B'\}$$

Matrix Mapping

Let $A \in K^{m \times n}$ (field K)

F_A is the transformation determined by A

$$F_A : K^m \rightarrow K^n$$

$$\text{For } \alpha \in K^m, F_A(\alpha) = A\alpha$$

Composition of Mappings:

$$f : A \rightarrow B \text{ and } g : B \rightarrow C$$

$$g \cdot f : A \rightarrow C \text{ or } g \circ f : A \rightarrow C$$

$$(g \circ f)(\alpha) = g(f(\alpha)) \quad \alpha \in A$$

Let $f : A \rightarrow B$

1. f is injective (one to one) if

$$f(\alpha) = f(\alpha')$$

$$\implies \alpha = \alpha'$$

2. f is surjective (onto) if

$$\forall \beta \in B, \exists \alpha \in A : f(\alpha) = \beta$$

3. f is bijective (one to one correspondence) means that

f is injective and surjective

Identify mapping:

$$f : A \rightarrow A$$

$$\mathbb{1}_A : \mathbb{1}_A(\alpha) = \alpha$$

Inverse mapping:

$$f : A \rightarrow B \text{ and } g : B \rightarrow A, g = f^{-1} \text{ if}$$

$$f \circ g = \mathbb{1}_B \text{ and } g \circ f = \mathbb{1}_A$$

Linear Mapping

Let v and u are vector spaces over field K and

$F : v \rightarrow u$ then F is a linear mapping if :

1. For any $\alpha, \beta \in v$, $F(\alpha + \beta) = F(\alpha) + F(\beta)$
2. For any $k \in K, \alpha \in v$, $F(k\alpha) = kF(\alpha)$

Note:

$F : v \rightarrow u$ is a linear mapping if

For any $a, b \in k$ and $\alpha, \beta \in v$

$$F(a\alpha + b\beta) = aF(\alpha) + bF(\beta)$$

Example

Let $F : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be a projection onto the xy plane, where $F(x, y, z) = (x, y)$. Is F a linear mapping?

$$\text{Let } \alpha = (x_1, y_1, z_1), \beta = (x_2, y_2, z_2)$$

$$F(\alpha + \beta) = F(x_1 + x_2, y_1 + y_2, z_1 + z_2)$$

$$= (x_1 + x_2, y_1 + y_2)$$

$$= (x_1, y_1) + (x_2, y_2)$$

$$= F(\alpha) + F(\beta)$$

$$F(k\alpha) = F(kx_1, ky_1, kz_1)$$

$$= (kx_1, ky_1)$$

$$= k(x_1, y_1)$$

$$= kF(\alpha)$$

Example

$$G : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$G(x, y) = (x + 1, y + 2)$. Is G a linear mapping?

$$\text{Let } \alpha = (x_1, y_1), \beta = (x_2, y_2)$$

$$G(\alpha) = (x_1 + 1, y_1 + 2)$$

$$G(\beta) = (x_2 + 1, y_2 + 2)$$

$$G(\alpha + \beta) = G(x_1 + x_2, y_1 + y_2)$$

$$= (x_1 + x_2 + 1, y_1 + y_2 + 2)$$

$$G(\alpha + \beta) - G(\alpha) = (x_2, y_2) \neq G(\beta)$$

G is not a linear mapping

Example

$$J : v \Rightarrow \mathbb{R}$$

$$J(f(t)) = \int_0^1 f(t) dt$$

$$J(af(t) + bg(t)) = \int_0^1 af(t) + bg(t) dt$$

$$= \int_0^1 af(t) dt + \int_0^1 bg(t) dt$$

$$= a \int_0^1 f(t) dt + b \int_0^1 g(t) dt$$

$$= aJ(f(t)) + bJ(g(t))$$

Thm: Let v and u be vector space over field K and $S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$ be a basis of v , then there exists a linear mapping $F : V \rightarrow U$ such that any $\beta_1, \beta_2, \dots, \beta_n \in U$ is a unique representation with respect to F such that $F(\alpha_i) = \beta_i$

$$F : V \rightarrow U \text{ where } V \rightarrow S = \{\alpha_1, \alpha_2, \dots, \alpha_n\}, U \rightarrow \{\beta_1, \beta_2, \dots, \beta_n\}$$

Proof:

1. Define F
2. F is a linear mapping
3. F is unique

Claim 1:

$$\gamma \in V$$

$$\gamma = a_1\alpha_1 + \dots + a_n\alpha_n \quad - a_1 \text{ is unique}$$

$$F(\gamma) = a_1\beta_1 + \dots + a_n\beta_n$$

$$F(\alpha_1) = F(1\alpha_1 + 0\alpha_2 + \dots + 0\alpha_n) = 1\beta_1$$

$$F(\alpha_i) = 1\beta_i$$

Claim 2: F is a linear mapping

$$\text{Let } v, w \in V$$

$$v = a_1\alpha_1 + \dots + a_n\alpha_n$$

$$w = b_1\alpha_1 + \dots + b_n\alpha_n$$

$$F(v) = \sum_{j=1}^n a_j\beta_j$$

$$F(w) = \sum_{j=1}^n b_j \beta_j$$

$$F(v + w) = F((a_1 + b_1)\alpha_1 + \dots + (a_n + b_n)\alpha_n)$$

$$= \sum_{j=1}^n (a_j + b_j) \beta_j$$

$$= \sum_{j=1}^n a_j \beta_j + \sum_{j=1}^n b_j \beta_j$$

$$= F(v) + F(w)$$

$$F(kv) = F(k(a_1\alpha_1 + \dots + a_n\alpha_n))$$

$$= F(ka_1\alpha_1 + \dots + ka_n\alpha_n)$$

$$= \sum_{j=1}^n ka_j \beta_j$$

$$= k \sum_{j=1}^n a_j \beta_j$$

$$= kF(v)$$

F is a linear mapping

Claim 3:

$G : v \rightarrow u$ is a linear mapping and $G(\alpha_i) = \beta_i$

$$G(a_1\alpha_1 + \dots + a_n\alpha_n) = \sum_{j=1}^n G(a_j\alpha_j) \quad - G \text{ is a linear mapping}$$

$$= \sum_{j=1}^n G(a_j\alpha_j)$$

$$= \sum_{j=1}^n a_j G(\alpha_j)$$

$$= \sum_{j=1}^n a_j \beta_j$$

$$= F(v)$$

$$= F(a_1\alpha_1 + \dots + a_n\alpha_n)$$

Isomorphism

Definition

Two vector space v and s over field K are isomorphic if there exist $F : v \rightarrow u$ such that

1. F is bijective
2. F is a linear mapping

Example

vector space v and $s = \{\alpha_1, \dots, \alpha_n\}$ is a basis of v

$Proj_s \alpha$ for all α in v is an isomorphism between v and K^n

Kernel and image of a linear mapping

$$F : v \rightarrow u$$

Kernel of $F : Ker(F)$

$$Ker(F) = \{\alpha \in v : F(\alpha) = 0\}$$

Image of $F : Im(F)$

$$Im(F) = \{\beta \in u : \exists \alpha \in v, F(\alpha) = \beta\}$$

Nullity of $F : Dim(Ker(F))$

Rank of $F : Dim(Im(F))$

$$\text{Dim}(v) = \text{Dim}(\text{Ker}(F)) + \text{Dim}(\text{Im}(F))$$

Note:

$\text{Ker}(F)$ is a subspace of v

$\text{Im}(F)$ is a subspace of u

Thm:

Suppose $\{\alpha_1, \dots, \alpha_n\}$ spans V and $F : v \rightarrow u$ is a linear mapping, then $F(\alpha_1), \dots, F(\alpha_n)$ spans the image of F ($\text{Im}(F)$)

Proof:

If $\gamma \in v$ then $\gamma = a_1\alpha_1 + \dots + a_n\alpha_n$ - $a'_i s$

$$F(\gamma) = F(a_1\alpha_1 + \dots + a_n\alpha_n)$$

$$= \sum_{j=1}^n F(a_j\alpha_j)$$

$$= \sum_{j=1}^n a_j F(\alpha_j)$$

$$= a_1 F(\alpha_1) + a_2 F(\alpha_2) + \dots + a_n F(\alpha_n)$$

Singularity

$$F : v \rightarrow u$$

$$\text{Ker}(F) = \{0\}$$

If $\exists \alpha \in v : \alpha \neq 0$ and $F(\alpha) = 0$, then F is singular

$$A \in n \times n$$

$$\text{Det}(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_1 \sigma^1 a_2 \sigma^2 \dots a_n \sigma^n \text{ where } \text{sgn}(\sigma) \text{ is the parity}$$

Group

A collection of objects and a binary operation such that:

1. Closure: $\forall a, b \in G, a * b \in G$
2. Associativity: $\forall a, b, c \in G, a * (b * c) = (a * b) * c$
3. Identity: $\exists e \in G : \forall a \in G, a * e = e * a = a$
4. Inverse: $\forall a \in G, \exists a' \in G : a * a' = a' * a = e$

Semi-group operation $*$: A collection of objects and a binary such that it satisfies 1, 2, 3

Abelian Group (commutative group):

$(G, *)$ is an Abelian group if $(G, *)$ is a group and $\forall a, b \in G, a * b = b * a$

Ring

A collection of items and two operations (usually addition $+$ and multiplication \times) such that

1. $(R, +)$ is a commutative group
2. (R, \times)
 1. $\forall a, b \in R, a \times b \in R$
 2. $\forall a, b, c \in R, a \times (b + c) = (a \times b) + (a \times c)$
 3. $\forall a, b, c \in R, (a + b) \times c = (a \times c) + (b \times c)$
 4. $\forall a, b, c \in R, a \times (b \times c) = (a \times b) \times c$

$$5. \exists 0 \in R : 0 \times a = 0 \forall a \in R$$

Field

A field F is a ring where $F' =$ all the elements in F without the 0-element and (F', \times) is commutative group

Symmetric Group S_x on x

Group $G : |G|$ is the order of G = number of elements in G

$S_x = \{\sigma : x \rightarrow x \text{ such that } \sigma \text{ is bijective}\}$

σ – shuffles x

σ is permutation of $x \in X$

$$|S_n| = n! = n(n-1) \dots 2 \cdot 1$$

σ, τ

$\sigma \circ \tau$ or $\tau \circ \sigma$

$\sigma \circ \tau \in S_x, \tau \circ \sigma \in S_x$

Example

Let $X = \{1, 2, 3\}$

$$|S_x| = 3! = 6$$

$S_x = \{e, \sigma_1, \sigma_2, \tau_1, \tau_2, \tau_3\}$

$$e = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 1 & 2 & 3 \end{bmatrix}$$

$$\sigma_1 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 2 & 3 & 1 \end{bmatrix}$$

$$\sigma_2 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 3 & 1 & 2 \end{bmatrix}$$

$$\tau_1 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 1 & 3 & 2 \end{bmatrix}$$

$$\tau_2 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 3 & 2 & 1 \end{bmatrix}$$

$$\tau_3 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 2 & 1 & 3 \end{bmatrix}$$

Is $\alpha_1 \circ \tau_1 = \tau_1 \circ \alpha_1$

$$\alpha_1 \circ \tau_1 = \tau_1 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 1 & 3 & 2 \\ \downarrow & \downarrow & \downarrow \\ 3 & 2 & 1 \end{bmatrix} = \tau_2$$

$$\tau_1 \circ \sigma_1 = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 2 & 3 & 1 \\ \downarrow & \downarrow & \downarrow \\ 2 & 1 & 3 \end{bmatrix} = \tau_3$$

sign or parity $sgn(\sigma)$

$$x = \{1, 2, 3\}, \sigma \in S_x$$

$$\sigma = \{\sigma^1, \sigma^2, \sigma^3\}$$

$$\text{Example: } \sigma = \sigma^1, \sigma_1^1 = 2, \sigma_1^2 = 1, \sigma_1^3 = 3$$

Even and odd parity

$$\sigma_1 = \{2, 1, 3\}$$

number of i and $k : i < k$ but $\sigma^i > \sigma^k$

$$\tau_2 = \{3, 2, 1\}$$

$$sgn(\alpha) = \prod_{i < k} \frac{\sigma^k - \sigma^i}{k - i}$$

$$\text{For } S_3: sgn(\sigma) = \frac{\sigma^3 - \sigma^2}{3 - 2} \frac{\sigma^2 - \sigma^1}{2 - 1} \frac{\sigma^3 - \sigma^1}{3 - 1}$$

$$= \frac{1-2}{3-2} \frac{2-3}{2-1} \frac{1-3}{3-1}$$

$$= -1$$

Week 6 Session 1

Outline

Determinants

Eigenvalues and Eigenvectors

$$f: \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$x_{t+1} = f(x_t)$$

Fixed point: $f(\hat{x}) = \hat{x}$

$$A\hat{x} = \hat{x}$$

$$A \in n \times n$$

$$\text{Det}(A) = \sum_{\sigma \in S_n} sgn(\sigma) a_1 \sigma^1 \dots a_n \sigma^n$$

$$sgn(\sigma) = \prod_{i > k} \frac{\sigma^i - \sigma^k}{i - k}$$

$$sgn(\sigma) = \begin{cases} +1 & \text{if } \sigma \text{ is even} \\ -1 & \text{if } \sigma \text{ is odd} \end{cases}$$

$$sgn(\sigma) = (-1)^{N(\sigma)}$$

$N(\sigma) \equiv$ number of (i, k) such that $(i > k)$ but $\sigma^i < \sigma^k$

$$\sigma = \begin{bmatrix} 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow \\ 2 & 3 & 1 \end{bmatrix}$$

$$i = \begin{bmatrix} 3 \\ 3 \\ 2 \end{bmatrix}, k = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}$$

$$N(\sigma) = 2$$

$$\text{sgn}(\sigma) = (-1)^2 = 1$$

Facts:

1. Let $g(x_1, \dots, x_n) = \prod_{i > k} (x_i - x_k)$
 2. Let $\sigma(g) = \prod_{i > k} (x_{\sigma^i} - x_{\sigma^k})$

$$\sigma(g) = \begin{cases} +g & \text{if } \sigma \text{ is even} \\ -g & \text{if } \sigma \text{ is odd} \end{cases}$$

$$\sigma(g) = \text{sgn}(\sigma)g$$
 3. Let $\sigma, \tau \in S_n$, $\text{sgn}(\sigma \circ \tau) = \text{sgn}(\sigma)\text{sgn}(\tau)$

$$\begin{aligned} \text{sgn}(\sigma \circ \tau)g &= (\sigma \circ \tau)g \\ &= \sigma(\tau(g)) \\ &= \text{sgn}(\sigma)\tau(g) \\ &= \text{sgn}(\sigma)\text{sgn}(\tau)g \\ \text{sgn}(\sigma \circ \tau) &= \text{sgn}(\sigma)\text{sgn}(\tau) \end{aligned}$$
 4. $\text{sgn}(\sigma) = \text{sgn}(\sigma^{-1})$

$$\sigma \circ \sigma^{-1} = \epsilon$$

$$\text{sgn}(\sigma \circ \sigma^{-1}) = \text{sgn}(\sigma)\text{sgn}(\sigma^{-1}) = \text{sgn}(\epsilon) = 1$$

check the two cases of $\text{sgn}(\sigma) \in \{+1, -1\}$
 5. Let $\sigma = j_1 j_2 \dots j_n$ for scalar a_{ij} and $a_{j_1 1} a_{j_2 2} \dots a_{j_n n} = a_{1 k_1} a_{2 k_2} \dots a_{n k_n}$

$$\sigma(k_i) = i$$

Let us assume that $\tau = k_1 k_2 \dots k_n$

$$\begin{aligned} \tau(j_i) &= i \\ \tau(j_i) &= \tau(\sigma(i)) = i \\ (\tau \circ \sigma)i &= i \\ \tau &= \sigma^{-1} \end{aligned}$$
-

Thm: If σ^i and σ^j are interchanged in $\sigma = (\sigma^1, \dots, \sigma^n)$ to given $\hat{\sigma}$, then $\text{sgn}(\hat{\sigma}) = -\text{sgn}(\sigma)$

Proof: $\prod_{i > k} \frac{\sigma^i - \sigma^j}{i - j}$

Example

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$\text{Det}(A) = \sum_{\sigma \in S_2} \text{sgn}(\sigma) a_{1\sigma^1} a_{2\sigma^2}$$

$$S_2 = \{\epsilon, \sigma\}$$

$$\epsilon = \begin{bmatrix} 1 & 2 \\ \downarrow & \downarrow \\ 1 & 2 \end{bmatrix}$$

$$\sigma = \begin{bmatrix} 1 & 2 \\ \downarrow & \downarrow \\ 2 & 1 \end{bmatrix}$$

$$\text{sgn}(\epsilon) = +1$$

$$\text{sgn}(\sigma) = -1$$

$$\text{Det}(A) = (+1)a_{11}a_{22} + (-1)a_{12}a_{21}$$

$$= a_{11}a_{22} - a_{12}a_{21}$$

Properties of Determinants

1. $\text{Det}(A) = \text{Det}(A^T)$

$$A = [a_{ij}]$$

$$A^T = B = [b_{ij}] \text{ where } b_{ij} = a_{ji}$$

$$\text{Det}(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma^1} \dots a_{n\sigma^n}$$

$$\text{Det}(A^T) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) b_{1\sigma^1} \dots b_{n\sigma^n}$$

$$= \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{\sigma^1 1} \dots a_{\sigma^n n} \quad - A^T = B$$

$$= \sum_{\sigma \in S_n, \tau = \sigma^{-1}} \text{sgn}(\sigma) a_{1\tau^1} \dots a_{n\tau^n} \quad - \text{Fact 5}$$

$$= \sum_{\tau \in S_n} \text{sgn}(\tau) a_{1\tau^1} \dots a_{n\tau^n} \quad - \text{Fact 4}$$

$$= \text{Det}(A)$$

2. If A is a square matrix and two rows (or columns) are interchanged to form B , then

$$\text{Det}(A) = -\text{Det}(B)$$

3. If A is a square matrix with a zero row (or zero column), then

$$\text{Det}(A) = 0$$

$$\sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma^1} \dots a_{n\sigma^n}$$

4. If A has two identical rows (or two identical columns), then the determinant

$$\text{Det}(A) = 0$$

$$A = \begin{bmatrix} \dots & R_i & \dots \\ \dots & R_j & \dots \end{bmatrix} \quad R_i = R_j$$

$$A' = \begin{bmatrix} \dots & R_i & \dots \\ \dots & R_j & \dots \end{bmatrix}$$

$$\text{Det}(A) = -\text{Det}(A')$$

$$\text{Det}(A) = -\text{Det}(A') = -\text{Det}(A) \implies \text{Det}(A) = 0$$

5. If scaling a row (or a column) by k transforms a square A to B , then

$$\text{Det}(B) = k\text{Det}(A)$$

$$A = [\dots \quad R_i \quad \dots]$$

$$\sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma^1} \dots a_{n\sigma^n}$$

$$B = [\dots \quad kR_i \quad \dots]$$

$$\sum_{\sigma \in S_n} \text{sgn}(\sigma) b_{1\sigma^1} \dots b_{n\sigma^n}$$

$$= \sum_{\sigma \in S_n} \text{sgn}(\sigma) k(a_{1\sigma^1} \dots a_{n\sigma^n})$$

$$\text{Det}(B) = k\text{Det}(A)$$

6. $R_i : R_i + kR_j$

If adding a scalar multiple of a row (or a column) to another transforms a square matrix A to B , then

$$\text{Det}(B) = \text{Det}(A)$$

$$\text{Det}(B) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma^1} \dots a_{i\sigma^i} + ka_{j\sigma^i} \dots a_{n\sigma^n}$$

$$= \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1\sigma^1} \dots a_{i\sigma^i} \dots a_{n\sigma^n} + k \sum_{\tau \in S_n} \text{sgn}(\tau) (a_{1\tau^1} \dots a_{j\tau^i} \dots a_{n\tau^n})$$

$$\text{where } a_{1\tau^1} \dots a_{j\tau^i} \dots a_{n\tau^n} = 0$$

$$= \text{Det}(A)$$

7. If E is an elementary matrix and A is a square matrix, then

$$\text{Det}(EA) = \text{Det}(E)\text{Det}(A)$$

8. $\text{Det}(AB) = \text{Det}(A)\text{Det}(B)$

9. If A is a diagonal matrix

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & a_{nn} \end{bmatrix}$$

then $\text{Det}(A) = \prod_{i=1}^n a_{ii}$

10. If A is a triangular matrix, then

$$\text{Det}(A) = \prod_{i=1}^n a_{ii}$$

11. $\text{Det}(A^{-1}) = (\text{Det}(A))^{-1}$ if $\text{Det}(A) \neq 0$

Thm:

The following statements are equivalent

1. A is invertible - $M = [A|I] \sim [I|A^{-1}]$
2. $Ax = 0$ has only the zero solution
3. $\text{Det}(A) \neq 0$

$$A = E_n E_{n-1} \dots E_1 I$$

$$\text{Det}(A) = \text{Det}(E_n) \dots \text{Det}(E_1) \text{Det}(I) \neq 0$$

$$\text{Det}(E_n) \neq 0, \text{Det}(E_1) \neq 0, \text{Det}(I) = 1$$

Block matrix

$$M = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

A_{11} is $r \times r$

A_{22} is $s \times s$

$$M_1 = \begin{bmatrix} I & 0 \\ A_{21}A_{11}^{-1} & I \end{bmatrix}, M_2 = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} - A_{21}A_{11}^{-1}A_{12} \end{bmatrix}$$

$$M = M_1 M_2 = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

$$\text{If } B = \begin{bmatrix} B_{11} & 0 & \dots & 0 \\ B_{21} & B_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ B_{n1} & B_{n2} & \dots & B_{nn} \end{bmatrix}$$

$$\text{Det}(B) = \prod_{i=1}^n \text{Det}(B_{ii})$$

$$\text{Det}(M) = \text{Det}(M_1) \text{Det}(M_2)$$

$$= 1 \times \text{Det}(A_{11}) \text{Det}(A_{22} - A_{21}A_{11}^{-1}A_{12})$$

$$= \text{Det}(A_{11}) \text{Det}(A_{22} - A_{21}A_{11}^{-1}A_{12})$$

Determinants and volume

Let $u_1, \dots, u_n \in \mathbb{R}^n$

$$V^T = \begin{bmatrix} \dots & \dots & \dots & \dots \\ v_1 & v_2 & \dots & v_n \\ \dots & \dots & \dots & \dots \end{bmatrix}$$

$$V = \begin{bmatrix} \dots & v_1 & \dots \\ \dots & v_2 & \dots \\ \dots & \dots & \dots \\ \dots & v_n & \dots \end{bmatrix}$$


volume enclosed by v_1, \dots, v_n is $|Det(v)|$

Cofactors and minors

$$A \equiv [a_{ij}]$$

$M \equiv$ Delete row i and column j from A

$$M_{ij} \in (n-1) \times (n-1)$$



$$\text{say } A = \begin{bmatrix} 4 & 5 & 7 \\ -2 & 1 & 0 \\ 3 & 8 & 7 \end{bmatrix}$$

$$\text{then } M_{12} = \begin{bmatrix} -2 & 0 \\ 3 & 7 \end{bmatrix}$$

$$\text{Minor: } m_{ij} = Det(M_{ij})$$

$$\text{Cofactor: } c_{ij} = (-1)^{i+j} m_{ij}$$

$$Det(A) = \sum_{j=1}^n a_{ij} c_{ij} = \sum_{i=1}^n a_{ij} c_{ij} \quad - \text{Laplace Expansion}$$

$$\text{Adjoint: } A = [a_{ij}]$$

$$\tilde{A} = [c_{ij}]$$

$$Adj(A) = \tilde{A}^T$$

$$A^{-1} = \frac{Adj(A)}{Det(A)}$$

Eigenvalues and Eigenvectors

Fixed point:

$$x_{t+1} = f(x_t)$$

Fixed point \hat{x} is such that $f(\hat{x}) = \hat{x}$

Example

$$y = f(x) = x^2$$

$$f: \mathbb{R} \rightarrow \mathbb{R}$$

$$x = x^2$$

$$\hat{x} \in \{0, 1\}$$

$$y = f(x) = x^3$$

$$f : \mathbb{R} \rightarrow \mathbb{R}$$

$$\hat{x} \in \{-1, 0, 1\}$$

$$Df = \frac{df(x)}{dx} = f'(x)$$

$$f(x) = e^x \text{ is the eigen function}$$

$$\frac{d}{dx} e^{cx} = ce^{cx} \text{ where } c \text{ is the eigen value}$$

Week 6 Session 2

Outline

Eigenvalues and Eigenvectors

Linearly Independent Eigenvectors (LIE)

Eigenvectors generalizes fixed points

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n$$

$$\hat{x} \text{ is a fixed point for } f \text{ is } f(\hat{x}) = \hat{x}$$

Example

$$y = f(x) = x^2; \hat{x} \in \{0, 1\}$$

$$y = f(x) = x^3; \hat{x} \in \{-1, 0, 1\}$$

$$D : f \rightarrow g$$

$$Df = \frac{df(x)}{dx}$$

$$\hat{f} = e^x \text{ because } \frac{d\hat{f}}{dx} = e^x = \hat{f}$$

$$\frac{d}{dx} e^{cx} = ce^{cx} \text{ where } c \text{ is the eigen value and } e^{cx} \text{ is the eigenfunction}$$

Matrix

$$A \in \mathbb{C}^{n \times n}$$

$$A : \mathbb{C}^n \rightarrow \mathbb{C}^n$$

$$\hat{x} \text{ is a fixed point for } A \text{ if } A\hat{x} = \hat{x}$$

Definition:

Suppose $x \neq 0$ and $\lambda \in \mathbb{C}$ if $Ax = \lambda x$ for $A \in \mathbb{C}^{n \times n}$ then


λ is the eigenvalue and x is the corresponding eigenvector

$$Ax = \lambda x$$

Example

$$A = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$$

$$x_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, Ax_1 = \begin{bmatrix} 3 \\ 7 \end{bmatrix}$$



$$x_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, Ax_2 = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Eigenvectors are not unique

$$Ax = \lambda x$$

$$A(cx) = \lambda(cx) \quad - c \in \mathbb{C}$$

$$x_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, x_3 = \begin{bmatrix} -2 \\ 2 \end{bmatrix}$$

$$Ax_3 = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} -2 \\ 2 \end{bmatrix} = \begin{bmatrix} -2 \\ 2 \end{bmatrix}$$

Recall: The following statements are equivalent for a square matrix A

1. A is invertible
2. $Ax = 0$ has the zero-vector as the only solution
3. $\text{Det}(A) \neq 0$

Thm: $\lambda \in \mathbb{C}$ is an eigenvalue of A if and only if $\text{Det}(A - \lambda I) = 0$

Proof: $Ax = \lambda x$ and $x \neq 0$ (Definition of eigenvector and eigenvalue)

$$Ax = \lambda Ix$$

$$Ax - \lambda Ix = \mathbf{0}$$

$$(A - \lambda I)x = \mathbf{0} \text{ and } x \neq 0$$

$$\text{Det}(A - \lambda I) = 0$$

Characteristic Polynomial

$$P_A(\lambda) = \text{Det}(A - \lambda I)$$

$$P_A(\lambda) = 0$$

Finding eigenvalues and eigenvectors

1. Define the characteristic polynomial $P_A(\lambda)$
2. Solve $P_A(\lambda) = 0$
3. For each λ , solve for x in $(A - \lambda I)x = 0$

Example

$$A = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$$

Find its eigenvalues and eigenvectors

Step 1: $P_A(\lambda)$

$$P_A(\lambda) = \text{Det}(A - \lambda I)$$

$$= \begin{bmatrix} 2 - \lambda & 1 \\ 3 & 4 - \lambda \end{bmatrix}$$

$$= (2 - \lambda)(4 - \lambda) - 3$$

$$= \lambda^2 - 6\lambda + 5$$

Step 2: Solve $P_A(\lambda) = 0$

$$\lambda_1 = 1, \lambda_2 = 5$$

Step 3:

For $\lambda = 1$

$$(A - \lambda I)x = \mathbf{0}$$

$$\begin{bmatrix} 2 - 1 & 1 \\ 3 & 4 - 1 \end{bmatrix} x = \mathbf{0}$$

$$\begin{bmatrix} 1 & 1 \\ 3 & 3 \end{bmatrix} x = \mathbf{0}$$

$$x = \begin{bmatrix} z \\ -z \end{bmatrix} \text{ for } z \in \mathbb{C}, z \neq 0$$

$$x = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ is the corresponding eigenvector for } \lambda = 1$$

For $\lambda = 5$

$$(A - \lambda I)x = \mathbf{0}$$

$$\begin{bmatrix} 2 - 5 & 1 \\ 3 & 4 - 5 \end{bmatrix} x = \mathbf{0}$$

$$\begin{bmatrix} -3 & 1 \\ 3 & -1 \end{bmatrix} x = \mathbf{0}$$

$$x = \begin{bmatrix} z \\ 3z \end{bmatrix} \text{ for } z \in \mathbb{C}, z \neq 0$$

$$x = \begin{bmatrix} 1 \\ 3 \end{bmatrix} \text{ is the corresponding eigenvector for } \lambda = 5$$

Eigenspace

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\lambda_1 = 1, \lambda_2 = 1$$

$$Ax = \lambda x$$

$$Ix = 1x$$

$$P_A(\lambda) = \text{Det}(A - \lambda I) = \prod_{i=1}^n (\lambda_i - \lambda)$$

$$P_A(\lambda) = (\lambda_1 - \lambda)(\lambda_2 - \lambda) = (1 - \lambda)(5 - \lambda) = \lambda^2 - 6\lambda + 5$$

$$\text{Det}(A - \lambda I) = \prod_{i=1}^n (\lambda_i - \lambda)$$

$$\text{Det}(A - \lambda I) = \prod_{i=1}^n (\lambda_i - \lambda)^{m_j} \text{ where } k \leq n$$

$$\sum_{j=1}^k m_j = n$$

Let A be a diagonal matrix

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & a_{nn} \end{bmatrix}$$

$$A - \lambda I = \begin{bmatrix} a_{11} - \lambda & 0 & \dots & 0 \\ 0 & a_{22} - \lambda & \dots & 0 \\ \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & a_{nn} - \lambda \end{bmatrix}$$

$$\text{Det}(A - \lambda I) = \prod_{i=1}^n (a_{ii} - \lambda) = P_A(\lambda) \text{ where } a_{ii} = \lambda_i$$

Let A be a triangular matrix

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ a_{21} & a_{22} & \dots & 0 \\ \dots & \dots & \dots & 0 \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\text{Det}(A - \lambda I) = \prod_{i=1}^n (a_{ii} - \lambda) \text{ where } a_{ii} = \lambda_i$$

$$\text{Thm: } \text{Det}(A) = \prod_{i=1}^n \lambda_i$$

$$\text{Proof: } \text{Det}(A - I\lambda) = \prod_{i=1}^n (\lambda_i - \lambda) \text{ where } \lambda = 0$$

$$\text{Det}(A) = \prod_{i=1}^n \lambda_i$$

LIE (Linearly independent eigenvectors)

Distinct Eigenvalues \implies LIE (Linearly independent eigenvectors)

LIE (Linearly independent eigenvectors) $\not\Rightarrow$ Distinct Eigenvalues

Thm: If $A \in \mathbb{C}^{n \times n}$ has n distinct eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$ Then A has n linearly independent eigenvectors $\alpha_1, \alpha_2, \dots, \alpha_n$

Proof: Mathematical Induction

Basis step:

Show that A with a distinct eigenvalue has linearly independent eigenvector set

$$Ax = \lambda x \text{ where } A, x \in 1 \times 1$$

$$cx = \lambda x, x \neq 0$$

$\{x\}$ is linearly independent

Induction step:

Induction hypothesis: Assume that $\lambda_1, \lambda_2, \dots, \lambda_n$ are unique (distinct) and $\alpha_1, \alpha_2, \dots, \alpha_n$ are linearly independent

If $\lambda_1, \lambda_2, \dots, \lambda_{n+1}$ are distinct eigenvalues then $\alpha_1, \alpha_2, \dots, \alpha_{n+1}$ are linearly independent for $A\alpha_i = \lambda_i\alpha_i$

$P \rightarrow Q$

Proof: By contradiction ($\sim Q \rightarrow \sim P$)

$\sim Q : \exists c'_j$ such that $\alpha_{n+1} = \sum_{j=1}^n c_j \alpha_j$

$$A\alpha_{n+1} = \lambda_{n+1}\alpha_{n+1}$$

$$A\alpha_{n+1} = A \sum_{j=1}^n c_j \alpha_j = \sum_{j=1}^n c_j A\alpha_j = \sum_{j=1}^n c_j \lambda_j \alpha_j \quad Eq. 3$$

$$A\alpha_{n+1} = \lambda_{n+1}\alpha_{n+1} = \lambda_{n+1} \sum_{j=1}^n c_j \alpha_j = \sum_{j=1}^n c_j \lambda_{n+1} \alpha_j \quad Eq. 4$$

Eq. 3 – Eq. 4

$$\sum_{j=1}^n c_j (\lambda_j - \lambda_{n+1}) \alpha_j = 0 \text{ where } \alpha_j \neq 0$$

$$\exists j : \lambda_j - \lambda_{n+1} = 0$$

$$\lambda_j = \lambda_{n+1}$$

Contradiction $\sim P$

Thm: Projection matrix $p : p^2 = p$ and $p = p^T$, then $A_i = 0$ or $A_i = 1$

Proof: $px = \lambda x$

$$ppx = p\lambda x = \lambda px = \lambda \lambda x = \lambda^2 x$$

$$ppx = px = \lambda x$$

$$\lambda^2 x = \lambda x$$

$$\lambda \in \{0, 1\}$$

Thm: A and A^T have the same eigenvalues

Proof: $\text{Det}(A^T - \lambda I) = \text{Det}(A^T - \lambda I^T)$

$$= \text{Det}((A - \lambda I)^T)$$

$$= \text{Det}(A - \lambda I) \text{ where } \text{Det}(A) = \text{Det}(A^T)$$

$$\begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 2 - \lambda & 1 \\ 0 & -\lambda \end{bmatrix}$$

Thm: $\text{Tr}(A) = \sum_{i=1}^n a_{ii} = \sum_{i=1}^n \lambda_i$

Proof: $\text{Det}(A - \lambda I) = \prod_{i=1}^n (\lambda_i - \lambda) = P_A(\lambda)$

$$(x + a_1)(x + a_2) \dots (x + a_n) = x^n + x^{n-1}(a_1 + a_2 + \dots + a_n) + x^{n-2}(a_1 a_2 + a_1 a_3 + \dots + a_{n-1} a_n) + \dots + (a_1 a_2 \dots a_n)$$

$$\text{RHS: } \prod_{i=1}^n (\lambda_i - \lambda) = (-\lambda)^n + (-\lambda)^{n-1}(\lambda_1 + \lambda_2 + \dots + \lambda_n) + (-\lambda)^{n-2}(\lambda_1 \lambda_2 + \dots + \lambda_{n-1} \lambda_n)$$

$$\text{LHS: } \text{Det}(A - \lambda I)$$

$$C = A - \lambda I = \begin{bmatrix} a_{11} - \lambda & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} - \lambda & \dots & a_{2n} \\ \dots & \dots & \dots & \dots \\ a_{n1} & a_{n2} & \dots & a_{nn} - \lambda \end{bmatrix}$$

$$\text{Det}(A - \lambda I) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) c_1 \sigma^1 \dots c_n \sigma^n$$

If look at the term of $(-\lambda)^{n-1}$

$$(a_{11} - \lambda)(a_{22} - \lambda) \dots (a_{nn} - \lambda)$$

$$(-\lambda)^n + (-\lambda)^{n-1}(a_{11} + \dots + a_{nn})$$

$$\sum_{i=1}^n \lambda_i = \sum_{i=1}^b a_{ii}$$

coefficient of $(-\lambda)^{n-1}$