

MSDS650 Week 2 ANOVA Assignment - Nathan Worsham

Following the ANOVA exercises from <http://www.r-tutor.com/category/statistical-concept/anova>, I created the text files the exercises requested.

0.1 Completely Randomized Design

```
> setwd("/Users/worshamn/Dropbox/Documents/Regis/MSDS650/week2/")
> df1 = read.table("fastfood-1.txt", header=TRUE); df1
  Item1 Item2 Item3
1    22    52    16
2    42    33    24
3    44     8    19
4    52    47    18
5    45    43    34
6    37    32    39
> r = c(t(df1)); r
[1] 22 52 16 42 33 24 44  8 19 52 47 18 45 43 34 37 32 39
> f = c("Item1", "Item2", "Item3")
> k = 3
> n = 6
> tm = gl(k, 1, n*k, factor(f)); tm
[1] Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3
  Item1 Item2 Item3 Item1 Item2 Item3
Levels: Item1 Item2 Item3
> av = aov(r ~ tm)
> summary(av)
              Df Sum Sq Mean Sq F value Pr(>F)
tm              2  745.4   372.7    2.541  0.112
Residuals      15 2200.2   146.7
> #vertical
> rAsC = cbind(c(t(df1))); rAsC
      [,1]
[1,]    22
[2,]    52
[3,]    16
[4,]    42
[5,]    33
[6,]    24
[7,]    44
[8,]     8
[9,]    19
[10,]   52
[11,]   47
[12,]   18
[13,]   45
[14,]   43
[15,]   34
[16,]   37
```

```

[17,] 32
[18,] 39
> f = c("Item1", "Item2", "Item3")
> k = 3
> n = 6
> tm = gl(k, 1, n*k, factor(f)); tm
[1] Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3
Item1 Item2 Item3 Item1 Item2 Item3
Levels: Item1 Item2 Item3
> av = aov(rAsC ~ tm)
> summary(av)

```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
tm	2	745.4	372.7	2.541	0.112
Residuals	15	2200.2	146.7		

In this instance the P value is 0.112 which is well above 0.05, so the null hypothesis that the 3 menu items have the same mean sales is not rejected.

0.2 Randomized Block Design

```

> df2 = read.table("fastfood-2.txt", header=TRUE); df2
Item1 Item2 Item3
1 31 27 24
2 31 28 31
3 45 29 46
4 21 18 48
5 42 36 46
6 32 17 40
> r = c(t(df2)); r
[1] 31 27 24 31 28 31 45 29 46 21 18 48 42 36 46 32 17 40
> f = c("Item1", "Item2", "Item3")
> k = 3
> n = 6
> tm = gl(k, 1, n*k, factor(f)); tm
[1] Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3
Item1 Item2 Item3 Item1 Item2 Item3
Levels: Item1 Item2 Item3
> blk = gl(n, k, k*n); blk
[1] 1 1 1 2 2 2 3 3 3 4 4 4 5 5 5 6 6 6
Levels: 1 2 3 4 5 6
> av = aov(r ~ tm + blk)
> summary(av)

```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)
tm	2	538.8	269.39	4.959	0.0319 *
blk	5	559.8	111.96	2.061	0.1547
Residuals	10	543.2	54.32		

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

> #vertical
> rAsC = cbind(c(t(df2))); rAsC
      [,1]
 [1,]  31
 [2,]  27
 [3,]  24
 [4,]  31
 [5,]  28
 [6,]  31
 [7,]  45
 [8,]  29
 [9,]  46
[10,]  21
[11,]  18
[12,]  48
[13,]  42
[14,]  36
[15,]  46
[16,]  32
[17,]  17
[18,]  40
> f = c("Item1", "Item2", "Item3")
> k = 3
> n = 6
> tm = gl(k, 1, n*k, factor(f)); tm
 [1] Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3
      Item1 Item2 Item3 Item1 Item2 Item3
Levels: Item1 Item2 Item3
> blk = gl(n, k, k*n); blk
 [1] 1 1 1 2 2 2 3 3 3 4 4 4 5 5 5 6 6 6
Levels: 1 2 3 4 5 6
> av = aov(rAsC ~ tm + blk)
> summary(av)
              Df Sum Sq Mean Sq F value Pr(>F)
tm              2  538.8   269.39    4.959 0.0319 *
blk              5  559.8   111.96    2.061 0.1547
Residuals      10   543.2    54.32
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

0.3 Randomized Block Design

```

> df3 = read.csv("fastfood-3.csv")
> r = c(t(df3))
> r
 [1] 25 39 36 36 42 24 31 39 28 26 35 29 51 43 42 47 39 36 47 53 32 52 46 33
> f1 = c("Item1", "Item2", "Item3")
> f2 = c("East", "West")

```

```

> k1 = length(f1)
> k2 = length(f2)
> n = 4
> tm1 = gl(k1, 1, n*k1*k2, factor(f1))
> tm1
[1] Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3
     Item1 Item2 Item3 Item1 Item2 Item3 Item1
[20] Item2 Item3 Item1 Item2 Item3
Levels: Item1 Item2 Item3
> tm2 = gl(k2, n*k1, n*k1*k2, factor(f2))
> tm2
[1] East East East East East East East East East East East East West West
     West West West West West West West West
[23] West West
Levels: East West
> av = aov(r ~ tm1 * tm2)
> summary(av)
          Df Sum Sq Mean Sq F value    Pr(>F)
tm1         2  385.1   192.5    9.554 0.00149 **
tm2         1   715.0   715.0   35.481 1.23e-05 ***
tm1:tm2      2   234.1   117.0    5.808 0.01132 *
Residuals   18   362.8    20.2
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
> #column exercise
> rAsC = cbind(c(t(df3)))
> rAsC
      [,1]
[1,]    25
[2,]    39
[3,]    36
[4,]    36
[5,]    42
[6,]    24
[7,]    31
[8,]    39
[9,]    28
[10,]   26
[11,]   35
[12,]   29
[13,]   51
[14,]   43
[15,]   42
[16,]   47
[17,]   39
[18,]   36
[19,]   47
[20,]   53

```

```

[21,] 32
[22,] 52
[23,] 46
[24,] 33
> f1 = c("Item1", "Item2", "Item3")
> f2 = c("East", "West")
> k1 = length(f1)
> k2 = length(f2)
> n = 4
> tm1 = gl(k1, 1, n*k1*k2, factor(f1))
> tm1
[1] Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3 Item1 Item2 Item3
     Item1 Item2 Item3 Item1 Item2 Item3 Item1
[20] Item2 Item3 Item1 Item2 Item3
Levels: Item1 Item2 Item3
> tm2 = gl(k2, n*k1, n*k1*k2, factor(f2))
> tm2
[1] East East East East East East East East East East East East West West
     West West West West West West West West
[23] West West
Levels: East West
> av = aov(rAsC ~ tm1 * tm2)
> summary(av)

```

	Df	Sum Sq	Mean Sq	F value	Pr(>F)	
tm1	2	385.1	192.5	9.554	0.00149	**
tm2	1	715.0	715.0	35.481	1.23e-05	***
tm1:tm2	2	234.1	117.0	5.808	0.01132	*
Residuals	18	362.8	20.2			

```

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

In all three cases the extra "exercise" of changing from horizontal rows to vertical columns did not make much sense to me as all I had to do was used the `cbind` command to change it vertical but after that the code remained the exact same.

After running through all of the examples, I was still confused on how to use ANOVA but it is at least clear in these examples that the p value was the line that divides. David Lane from onlinestatbook.com (n.d.) gives some helpful plain english on what ANOVA does which is that it ?is used to test general rather than specific differences among means? and then he goes on to say that it tests the null hypothesis—which the examples spoke often about—that all the means are equal. When the null hypothesis is rejected, the conclusion is that at least one mean is different from at least one other mean but does not reveal which means are different from which (Lane, n.d.) and in these examples when the p-value is less than 0.05 they are recommending to reject the null hypothesis. So it would seem that ANOVA is used as a generic test for comparing the difference or variation between the means of some variable between two or more groups (Boslaugh, 2014). David Lane points out—at first it seems odd that it is not called ?Analysis of Means? but variance is what is analyzed to make inferences (Lane, n.d.)

Since it is used to analyze many types of experimental design, it would seem then that

the types of questions it answers are

- In the experiment, can variation (assuming there is any) be explained by the "grouping introduced by the classification factor(s)?" (r-bloggers.com, 2010) - Or on average do any statistically significant (not due to chance) differences exist between the treatment and control groups?

So wanting the experiment to "determine the effect and interplay of factors" on the treatment group, the design of the experiment is important to limit "the impact of variability?" (Krzywinski, 2014). Put another way, experimental design—when done properly—allows us to see both if the treatment causes the outcome and if the lack of treatment does not cause the outcome (Trochim, 2006). The design chosen can reduce the amount of trials participants have to take but possibly need a larger sample or allow a smaller number of participants with likely more trials required (Boslaugh, 2014). Either of these could translate to costs associated with the experiment. However depending on the circumstances, the design that is best may not be feasible or ethical, likely there will always be a compromise between what is ideal and what is feasible, but the design should be directed by what is most important to the question at hand (Boslaugh, 2014).

References

Lane, David. n.d. Retrieved from http://onlinestatbook.com/2/analysis_of_variance/intro.html, http://onlinestatbook.com/2/logic_of_hypothesis_testing/significance.html R-bloggers.com, 2010. Retrieved from <http://www.r-bloggers.com/one-way-analysis-of-variance-anova/> Krzywinski, Martin. 2014. Retrieved from <http://www.nature.com/nmeth/journal/v11/n6/full/nmeth.2974.html> Trochim, William. 2006. Retrieved from <http://www.socialresearchmethods.net/kb/desexper.php> Boslaugh, Sarah. 2014. Statistics in a Nutshell. Chapter 8 - Introduction to Regression and ANOVA. Chapter 18 - Research Design.