



STANDARD CANDLES

Rujuta Mokal, Orion Yeung, Michael Wozniak

Contents

Overview

The Distance Ladder

Different Types of Standard Candles

Broader use in Astrophysics and Cosmology

— Standard Candles

- An astrophysical object whose absolute magnitude (M) is known
- Apparent magnitude (m) can be measured
- Used to measure distance (d) through
$$\mu = m - M = 5 \log(d/10 \text{ pc})$$
- Distances of objects around standard candles can be found
- Commonly known: Cepheids, Type Ia supernovae



Cartoon of Standard Candles


— Applications of Standard Candles



Cartoon of Standard Candles, happy birthday Orion!

- Distance measurements
 - The Cosmic Distance Ladder
- Calculating the value of the Hubble constant (H_0)

— The Cosmic Distance Ladder



Cartoon of Cosmic Distance Ladder

- Approach to measuring larger scales
- Rungs of ladder → Standard Candles
- Shorter distances (0-100 pc): Parallax
- Intermediate distances (100 - 10^7 pc): Variable stars
- Large distances (10^6 - 10^{10} pc): Type 1A supernovae, Gravitationally lensed quasars

— Cepheid Variables, RR Lyrae $(100-10^6 \text{ pc})$

At the Harvard Observatory

Cepheid Variables


- Discovered by Henrietta Leavitt
- Found in Small and Large Magellanic clouds
- Luminosity (L) - Period (P) relationship discovered by Leavitt

RR Lyrae

- Discovered by Wilhelmina Flemming
- Found in Globular Clusters



Henrietta Leavitt



Movie of Cepheid over 1 month




Wilhelmina Flemming


— Cepheid Variables, RR Lyrae (100-10⁶ pc)

Finding distance:

- Photometric observations → Apparent magnitude (m)
- Light curve of m vs pulsation period (P)
- M from P
- $\mu = m - M = 5 \log(d/10 \text{ pc})$
- Their brightness is limited, only suitable out to $\leq 10^5 \text{ pc}$
 - Farther?



Luminosity-period light curve from LMC and Milky Way



Oscillating brightness of Cepheid




Picture of red giant

—Tip of Red Giant Branch

(100-10⁶ pc)

- Upper boundary of red giant branch: H fusion stops, max star size
- He flash: He fusion commences
- For Mass < ~2 M_{sun}, He flash → specific M
- Photometric measurement infrared band insensitive to star composition
- m_i can be found
- $M, m_i \rightarrow d$




HR diagram highlighting RGB

—Type Ia supernovae

($10^6 - 10^{10}$ pc)

- Occurs in binary systems with white dwarfs
- Mass accretes onto white dwarf from companion star
 - Reaches the Chandrasekhar mass
- Runaway fusion reaction causes massive explosion
- Fixed critical mass; give similar peak luminosities



Movie of Type IA SN

— Gravitationally Lensed Quasars

($10^9 - 10^{10}$ pc)

- Hot (10^{12} K) accretion disk around supermassive black hole
- Intrinsically variable luminosities
- Luminosity variations \rightarrow Time delay between gravitationally lensed quasar images
- Estimate distance from time-delay distance and redshift



Lensed Quasar smiling at us





Diagram of how lensing informs path lengths



Picture of Spiral Galaxy

– Spiral Galaxies




Graph of Tully-Fisher Relation

- Tully Fisher Relation: Correlation between mass and rotational velocity of galaxy (v)
- v measured through width of emission lines
- Constant mass-luminosity ratio; determine luminosity from mass
- Obtain M from luminosity


— Planetary Nebulae ($10^6 - 10^8$ pc)

- Glowing, ionized gas ejected from red giants
- Measure [O III] $\lambda 5007$ forbidden lines in a galaxy; identify planetary nebulae
- Find distribution of planetary nebula magnitudes at this wavelength
- Distribution will have a cutoff observed at m_c , corresponds to $M_c = -4.48$
- $\mu = m_c - M_c$, giving d



Distribution of m for PN in Andromeda

— Sunyaev-Zeldovich Effect $(10^6 - 10^8 \text{ pc})$




Cartoon of light seen via Sunyaev-Zeldovich effect

- In a galaxy cluster, high energy electrons scatter with CMB photons, giving photons an energy boost
- Distorts radiation spectrum of CMB; magnitude of effect is proportional to gas density (ρ) and size (S)
- Hot intergalactic gas emits X-rays with luminosity that is proportional to ρ^2 and S
- Use X-ray luminosity and variations in microwave observations to determine S
- Measure angular size (θ) of cluster directly
- Calculate distance, $d = S / \theta$

— The Hubble Constant (H_0)

- Hubble's Law
 - Expanding universe
 - Astrophysical objects moving away from each other
 - Rate of recession \propto Distance from earth
 - Recession velocity, $v = cz = H_0 d$
- Ways to measure
 - Through distance measurements
 - Cosmic Microwave Background




Cartoon of Hubble's Law

— Measuring the Hubble constant


For $z \ll 1$ (When effects of accelerated expansion are less significant)

1. Identify standard candles and measure luminosity (L), redshift (z) and flux (f)
2. Compute distance, d
3. Plot $v = cz$ vs d (for $z \ll 1$)
4. Slope gives H_0

For higher redshift, deviations from the straight line of cz vs d indicates towards nature of expansion



Plot of Hubble's Law demonstrating cosmological expansion



How measuring H_0 changed over past 2 decades

— Limits of Standard Candles

- Degree of variation between absolute magnitudes of standard candles
- Require identification of specific objects
- Standard candles may not be as standard for all objects
 - Distant type 1a supernovae
- Interstellar dust lowers observed luminosity
- Candle-specific challenges
 - Metallicity effects on period-luminosity relation in cepheids
 - Short duration of type 1a supernovae

— Summary

- Standard candles are objects with identifiable absolute magnitudes (M)
- Variety of types depending on distances to be measured
- Applications in understanding structure and expansion of the universe
- Limitations due to nature of objects and precision of measurement

