Moral Hazard Heterogeneity:
Genes and Health Insurance Influence Smoking after a Health Shock

Pietro Biroli and Laura Zwyssig

thanks to GEIGHEI team, Regina Seibel, and Pia Arce

University of Zürich

University of Connecticut
March 8th, 2021

Outline

Genes, health insurance, smoking choices

What:

Agenda : Use genes to assess individual level heterogeneity in economic parameters

Paper : Does reaction to a negative health shock depend on health insurance and genetic differences?

Genes, health insurance, smoking choices

What:

Agenda : Use genes to assess individual level heterogeneity in economic parameters

Paper : Does reaction to a negative health shock depend on health insurance and genetic differences?

How:

- Focus on smoking behavior following a cardio-vascular health shock
- Identification: US adults who receive free health insurance coverage after 65 (Medicare)
- Compare high and low genetic predisposition to smoking (G×E)

Genes, health insurance, smoking choices

What:

Agenda : Use genes to assess individual level heterogeneity in economic parameters

Paper : Does reaction to a negative health shock depend on health insurance and genetic differences?

How:

- Focus on smoking behavior following a cardio-vascular health shock
- Identification: US adults who receive free health insurance coverage after 65 (Medicare)
- Compare high and low genetic predisposition to smoking (G×E)

Why:

- Interplay between financial and biological constraints:
    - Health insurance buffers financial consequences of health shocks
    - Genetic predispositions influence behavioral responses
- Understand **heterogeneity in moral hazard**

Preview

- Identification assumptions:
    - Probability of health shock increases with age ...
    - ... but <u>no jump</u> at 65

- Results:
    - Health shock when uninsured $\Rightarrow$ less smoking...
    - ... but *only* for low PGS.
    - G$\times$E effect size: 27.9 pp

Why should we care?

- Genes cannot be changed ...
- ... but **environment** can!
- Interplay between genes (bio) and environment (econ) is essentially everywhere [Rutter, 2006]

- Understand differential effects of genetic predisposition based on environment (G×E)
  – shed light on pathways and mechanisms
  – provide measure of essential heterogeneity
  – revisit old econ concept with new lens

- Empirics: genes are cheaper to measure and more and more available ▶
- Theory: biologically founded model
- Policy: understand sources of heterogeneity and discuss *fairness*
- Individual: choose individualized intervention

Previous work:

- Smoking increases diseases and health-care costs (400k deaths, $300 billion per year in US)
  [United States Department of Health and Human Services, 2014, Goodchild et al., 2017, Ma et al., 2018, Xu et al., 2015]

- Severe health shocks reduce smoking
  [Clark and Etilé, 2002, Falba, 2005, Khwaja et al., 2006, Keenan, 2009, Sundmacher, 2012]

- Health insurance plays a big role (moral hazard)
  [Richards and Marti, 2014, Marti and Richards, 2017, Einav et al., 2013, Einav and Finkelstein, 2018]

- G × E: Use molecular genetics and measures of the environment and investigate their interplay.
  See [Caspi et al., 2002, Barcellos et al., 2018, Belsky et al., 2018, Rimfeld et al., 2018,
  Schmitz and Conley, 2017a, Schmitz and Conley, 2017b, Rosenquist et al., 2015]

- Genoeconomics: Find genetic determinants of economic behaviors: risk aversion, time and social pref., addiction.
  See [Benjamin et al., 2007, Cesarini et al., 2009, Benjamin et al., 2016]

Previous work:

- Smoking increases diseases and health-care costs (400k deaths, $300 billion per year in US)
  [United States Department of Health and Human Services, 2014, Goodchild et al., 2017, Ma et al., 2018, Xu et al., 2015]

- Severe health shocks reduce smoking
  [Clark and Etilé, 2002, Falba, 2005, Khwaja et al., 2006, Keenan, 2009, Sundmacher, 2012]

- Health insurance plays a big role (moral hazard)
  [Richards and Marti, 2014, Marti and Richards, 2017, Einav et al., 2013, Einav and Finkelstein, 2018]

- G × E: Use molecular genetics and measures of the environment and investigate their interplay.
  See [Caspi et al., 2002, Barcellos et al., 2018, Belsky et al., 2018, Rimfeld et al., 2018, Schmitz and Conley, 2017a, Schmitz and Conley, 2017b, Rosenquist et al., 2015]

- Genoeconomics: Find genetic determinants of economic behaviors: risk aversion, time and social pref., addiction.
  See [Benjamin et al., 2007, Cesarini et al., 2009, Benjamin et al., 2016]

↪ **Contribution**: leverage genetic variants to measure heterogeneity in moral hazard

Outline

Genetics for social scientists

- Human genome: series of 3 billion letter pairs (A,G,T,C)
- Genetic variants: one-letter changes across individuals (single nucleotide polymorphisms, SNPs)
- About $\approx$ 10m SNPs
  [The 1000 Genomes Project Consortium et al., 2015]
- Genome-wide association studies (GWAS) have identified genome-wide significant relationships between specific SNPs and health behaviors
- We use SNPs identified from large, replicated GWAS to create summarized genetic scores to study gene-by-SES interplay ▸ hits

PolyGenic Scores (PGS)

Discovery stage:

GWAS
[Liu et al., 2019]

Source: Liu et al. (2019)

Prediction stage:

$$PGS_i = \sum_{j=1}^{J} W_j G_{ij},$$

where G_{ij} is the genotype for individual i at SNP j, and the weight W_j is the OLS association between SNP j and outcome. Normalized to have mean zero and standard deviation one.

Polygenic Scores

In our setting
Data = Health and Retirement Study
Y = Pr(Smoking)
W_j = taken from [Liu et al., 2019]

Outline

Outline

Data

- Use HRS data: US-representative survey of $50+$ (1992-2016)
- Final sample size $= 26,022$ obs from $5,854$ individuals
    - Ever smoked ≥ 100 cigarettes (at baseline)
    - Ages: 60-70
    - Observed at least 2 waves
    - European descendants
   - Non-missing smoking, PGS, insurance, health shock

Variables

- Outcome $Y =$ current smoking status $\in \{0, 1\}$
  - Self reported

- High PGS $g =$ above 33^{rd} in PGS for smoking initiation
  - Use [Liu et al., 2019] for weights

- Health shock $=$ first diagnosis of acute cardiovascular condition
  - Heart problem: heart attack, coronary heart disease, angina, congestive heart failure, or other heart problems
  - or Stroke: transient ischemic attack

- Uninsured: self-reported coverage
  - Pre 65 uninsured: never report being covered by medical insurance
  - Post 65 everyone insured: eligible for Medicare
  - Who are the uninsured? ▶

Identification

Diff in response to health shock before and after 65
Main assumption: **timing** of health shock is exogenous
[Marti and Richards, 2017, Card et al., 2009]

Summary statistics

Table: Descriptive Statistics for Full Analytic Sample and Stratified by Genetic Group

	All	**Low PGS**	**High PGS**
	Mean (SD)	Mean (SD)	Mean (SD)
Age (baseline)	61.17 (1.93)	61.18 (1.96)	61.16 (1.92)
Smoking PGS	0.11 (0.99)	-0.96 (0.51)	0.64 (0.71)
No. waves present	4.44 (1.38)	4.46 (1.37)	4.43 (1.39)
	%	%	%
Female	50.42	46.97	52.14
Smoking (baseline)	29.55	27.06	30.79
Persistently uninsured	5.85	5.39	6.08
CV health shock	12.44	11.82	12.74
No. of individuals	5813	1929	3884
No. Person-year observations	25800	8602	17198

▸ Sumstats2   ▸ Uninsured

Smoking rates decrease with age

Share of smokers over different ages, split by PGS

Outline

Regression analysis

$$Y_{it} = \beta\, shock_{it} + \gamma\, post65_{it}$$
$$+ \lambda_1 \left(shock_{it} \times post65_{it}\right)$$
$$+ \lambda_2 \left(shock_{it} \times uninsured_i\right)$$
$$+ \lambda_3 \left(post65_{it} \times uninsured_i\right)$$
$$+ \lambda_4 \left(shock_{it} \times g_i\right)$$
$$+ \lambda_5 \left(post65_{it} \times g_i\right)$$
$$+ \delta_1 \left(shock_{it} \times post65_{it} \times uninsured_i\right)$$
$$+ \delta_2 \left(shock_{it} \times uninsured_i \times g_i\right)$$
$$+ \delta_3 \left(shock_{it} \times post65_{it} \times g_i\right)$$
$$+ \delta_4 \left(post65_{it} \times uninsured_i \times g_i\right)$$
$$+ \zeta \left(shock_{it} \times post65_{it} \times uninsured_i \times g_i\right)$$
$$+ \sum_{a=1}^{3} \phi_a\, age_{it}^a + \eta_i + \tau_t + \varepsilon_{it}$$

Current smoking status (Y) regressed on the full set of interactions between the indicators for the health shock (*shock*), being uninsured pre-65 (*uninsured*), Medicare eligibility (*post*65), and high polygenic risk for smoking (g).

Controlling for age + individual and time F.E.

Effect of the shock on the outcomes

The derivative of the outcome with respect to shock is:

$$\frac{\partial Y_{it}}{\partial shock_{it}} = \beta + \lambda_1 post65_{it} + \lambda_2 uninsured_i + \lambda_4 g_i + \delta_1(post65 \times uninsured_i) + \delta_2(uninsured_i \times g_i) \tag{1}$$

$$+ \delta_3(post65_{it} \times g_i) + \xi(post65_{it} \times uninsured_i \times g_i)$$

Looking at the effect for the four different types (before-after 65, high-low g):

$$E\left[\frac{\partial Y_{it}}{\partial shock_{it}}|post65_{it} = 0, g_i = 0, uninsured_i = 1\right] = \beta + \lambda_2 \tag{2}$$

$$E\left[\frac{\partial Y_{it}}{\partial shock_{it}}|post65_{it} = 1, g_i = 0, uninsured_i = 1\right] = \beta + \lambda_1 + \lambda_2 + \delta_1 \tag{3}$$

$$E\left[\frac{\partial Y_{it}}{\partial shock_{it}}|post65_{it} = 0, g_i = 1, uninsured_i = 1\right] = \beta + \lambda_2 + \lambda_4 + \delta_2 \tag{4}$$

$$E\left[\frac{\partial Y_{it}}{\partial shock_{it}}|post65_{it} = 1, g_i = 1, , uninsured_i = 1\right] = \beta + \lambda_1 + \lambda_2 + \lambda_4 + \delta_1 + \delta_2 + \delta_3 + \xi \tag{5}$$

Calculating the first two differences as above:

$$(10) - (9) = \lambda_1 + \delta_1 \tag{6}$$

$$(12) - (11) = \lambda_1 + \delta_1 + \delta_3 + \xi \tag{7}$$

And the diff-in-diff (G×E):

$$(14) - (13) = \delta_3 + \xi \tag{8}$$

# Regression results

Controlling for individual and time F.E. + age:

Evidence of $G \times E$

Table: Effect of the shock by timing and PGS

Effect of health shock on smoking probability		
	Low PGS	High PGS
Pre 65	-0.167**	-0.108
	(0.069)	(0.083)
Post 65	0.089***	-0.115
	(0.024)	(0.093)
Effect of health insurance on effect of health shock		
	Low PGS	High PGS
Post 65 - Pre 65	0.256***	-0.008
	(0.079)	(0.124)
Differential effect of health insurance by genetic group		
	High PGS	- low PGS
Post 65 - Pre 65	-0.264*	
	(0.146)	

Notes: Summary of the effect of the shock on smoking for those who were uninsured before 65, stratified by timing of the shock (before vs. after 65) and genetic group (high vs. low PGS)

Different PGS cutoffs

Outline

Outline

Potential interpretation problems

What else can drive this relation?

- Something that jumps at 65 (besides medicare)
   - Retirement and income: no sharp change at age 65 [Card et al., 2008, Card et al., 2009]

Potential Confounders

Check for differential response to shock on possible confounders [Pei et al., 2018]

- Retirement ▸
- Individual income ▸
- Household income ▸
- Out of pocket med. expenditure ▸
- Having a partner ▸
- Partner Smoking ▸
- Mortality ▸ 2y ▸ 5y

Retirement: not likely a Confounder

Coefficient plot of the effect of the shock on being retired.

Wage: not likely a Confounder

Coefficient plot of the effect of the shock on log reported earnings.

Household Income: not likely a Confounder

Coefficient plot of the effect of the shock on household income.

Medical expenditure: not likely a Confounder

Coefficient plot of the effect of the shock on out of poket medical enxpenditure.

Having a partner: not likely a Confounder

Coefficient plot of the effect of the shock on having a partner.

Partner smoking: not likely a Confounder

Coefficient plot of the effect of the shock on having a partner.

2 year mortality: not likely a Confounder

Coefficient plot of the probability of dying within 2 years of the shock.

5 year mortality: not likely a Confounder

Coefficient plot of the probability of dying within 5 years of the shock.

Outline

Robustness checks

Robustness checks:

- Different cutoffs for high-PGS `▸ PGS`
    - Still holds for lower quartile, not for highest
    - Using continuous PGS, G×E effect = -0.13 (0.11)

- Using the publicly available PGS (older GWAS from [The Tobacco and Genetics Consortium et al., 2010]) `▸ PGS`
    - Similar pattern

- Different cutoffs for uninsured `▸ uninsured`
    - Noisy results for uninsured only 1/3 of the times

- Different cutoffs for age `▸ age`
   - Noisy results after age 72

Robustness: high PGS if above 25^{th} percentile

Coefficient plot of the main regression.

Robustness: high PGS if above 50^{th} percentile

Coefficient plot of the main regression.

Robustness: high PGS if above 75^{th} percentile

Coefficient plot of the main regression.

Robustness: using publicly available PGS (older GWAS)

Coefficient plot of the main regression.

Outline

Is it really genes?

Are there other characteristics that might be driving this relationship?

- Try to cut the data according to other dimensions:
  - Cognitive skills ▶
  - Conscientiousness ▶
  - Risk aversion ▶
  - Gender ▶
  - Education ▶
  - Income ▶
- Or according to other PGS:
  - Cognition PGS ▶
  - Risk aversion PGS ▶

Split by cognitive ability

Split by conscientiousness

Split by risk aversion

Split by gender

▶ back

Split by education

Split by former smoking behaviour

Outline

Outline

Limitations

Limitations:

- Small sample size of those hit by a shock (replication needed) ▶ Sumstats2
- Short-run smoking response
- Self-reported information
- Unobserved selection into DNA-sample

- External validity:
    - U.S. insurance system
    - Results only for old, uninsured people ▶ sumstats

Outline

Summary of Results

- Health shock when uninsured $\Rightarrow$ less smoking...
- ... but *only* for low PGS.
- Effect size is quite sizable (27.9 pp)

- Interaction between financial and "biological" constraints:
  - Health insurance buffers financial consequences of health shocks
  - Genetic predisposition to smoking mutes this effect (lower elasticity)

- Biological foundation of heterogeneity in **moral hazard** [Einav et al., 2013]

Conclusions

What does this tell us?

- Environment <u>and</u> genes jointly influence healthy behaviors

- Biological predispositions can tell a story about choices and economic fundamentals

Thank you

Outline

Outline

Smoking and lung cancer have same genetic hits

US Cigarette Use vs. Lung Cancer Deaths, 1900 - 2011

Death rates sources: Public-use data files, National Vital Statistics System, National Center for Health Statistics, Centers for Disease Control and Prevention; and Jemal et al., CA Cancer J Clin, 2010. Cigarette consumption sources: Tobacco Outlook Report, Economic Research Service, US Department of Agriculture; and Alcohol and Tobacco Trade and Tax Bureau, US Department of Treasury.

Smoking and Obesity Manhattan Plots

GSCAN: Cigarettes per Day

Nicotine Dependence:
Chromosome 15:
nAChR Gene Cluster
(Nicotinic Acetylcholine
Receptor)

Source: GSCAN (2019)

BMI:
More poligenic

Source: Speliotes et al., 2010

The biological pathways of the nicotine receptor gene

15q25: nAChR Gene Cluster

Nicotinic Acetylcholine Receptor

Improgo, et al., *Prog Neurobiol.* 2010 Oct;92(2):212-26

Outline

Summary statistics, by health shock timing

	Low PGS	High PGS	P value
Shock at ages 60-64			
	Mean (SD)	Mean (SD)	
Age (baseline)	60.49 (0.57)	60.47 (0.64)	0.78
Smoking PGS	-0.97 (0.54)	0.63 (0.68)	0.00
Years of education	12.2 (3.41)	12.11 (3.17)	0.81
Income (nominal $ 1000)	19.79 (27.82)	18.95 (30.14)	0.79
No. waves present	4.65 (1.32)	4.59 (1.32)	0.65
	%	%	
Female	48.7	45.05	0.51
Smoking (baseline)	31.3	37	0.28
Persistently uninsured	4.35	6.59	0.36
Avg. cessation rate (baseline smokers)	12.05	12.17	0.97
No. of individuals	115	273	
No. of Person-year individuals	535	1252	
Shock at ages 67-70			
	Mean (SD)	Mean (SD)	
Age (baseline)	61.33 (2.15)	61.04 (1.4)	0.19
Smoking PGS	-0.94 (0.48)	0.75 (0.79)	0.00
Years of education	12.68 (3.13)	12.41 (3.09)	0.46
Income (nominal $ 1000)	17.81 (27.44)	15.74 (20.15)	0.48
No. waves present	5.06 (1.15)	5.12 (0.84)	0.65
	%	%	
Female	40.71	49.1	0.14
Smoking (baseline)	30.09	35.59	0.31
Persistently uninsured	8.85	6.31	0.42
Avg. cessation rate (baseline smokers)	10.92	11.03	0.97
No. of individuals	113	222	
No. of Person-year individuals	572	1136	

▸ back   ▸ limitations

Who are the uninsured?

Table: Descriptive Statistics for Full Analytic Sample Stratified by Insurance Status

	All	Uninsured	Insured	P value
	Mean (SD)	Mean (SD)	Mean (SD)	
Age (baseline)	61.17 (1.93)	60.85 (0.94)	61.19 (1.97)	0.00
Smoking PGS	0.11 (0.99)	0.19 (0.92)	0.1 (1)	0.07
Years of education	12.48 (3.1)	10.44 (3.55)	12.61 (3.02)	0.00
Income (nominal $ 1000)	20.37 (34.95)	8.66 (13.29)	21.1 (35.74)	0.00
No. waves present	4.44 (1.38)	4.36 (1.28)	4.44 (1.39)	0.26
	%	%	%	
Female	50.42	57.06	50.01	0.01
Smoking (baseline)	29.55	46.76	28.49	0.00
Persistently uninsured	5.85	100	0	-
CV health shock	12.44	13.82	12.35	0.45
Avg. cessation rate (baseline smokers)	10.35	8.54	10.53	0.08
No. of individuals	5813	340	5473	
No. of person-year individuals	25800	24317	1483	

► back ► back

Outline

Regression results

Coefficients from Estimating the Linear Probability Model in Equation (2) Using OLS ▸ back

| | | | *Dependent variable:* | | |
| | | | Smoking status | | |
	(1)	(2)	(3)	(4)	(5)
Health Shock	−0.026	−0.026	−0.026	−0.050**	−0.050**
	(0.040)	(0.040)	(0.040)	(0.023)	(0.023)
Post-65	−0.066***	−0.021*	−0.021*	−0.011	−0.011
	(0.008)	(0.013)	(0.013)	(0.008)	(0.008)
Uninsured	0.171***	0.171***	0.171***		
	(0.027)	(0.027)	(0.027)		
High PGS	0.032**	0.032**	0.032**		
	(0.012)	(0.013)	(0.013)		
Shock × Post-65	0.014	0.023	0.022	0.022	0.022
	(0.055)	(0.056)	(0.056)	(0.032)	(0.032)
Shock × Uninsured	−0.198	−0.192	−0.193	−0.115	−0.117
	(0.184)	(0.184)	(0.184)	(0.073)	(0.073)
Post-65 × Uninsured	0.067	0.065	0.065	−0.052*	−0.053*
	(0.048)	(0.048)	(0.048)	(0.032)	(0.032)
Shock × High PGS	0.024	0.024	0.022	0.020	0.020
	(0.049)	(0.049)	(0.049)	(0.028)	(0.029)
Post-65 × High PGS	0.001	0.001	0.001	−0.003	−0.003
	(0.010)	(0.010)	(0.010)	(0.008)	(0.008)
Shock × Post-65 × Uninsured	0.286	0.280	0.280	0.233***	0.235***
	(0.250)	(0.251)	(0.250)	(0.086)	(0.085)
Shock × Uninsured × High PGS	0.189	0.188	0.189	0.037	0.039
	(0.219)	(0.220)	(0.219)	(0.112)	(0.111)
Shock × Post-65 × High PGS	−0.048	−0.048	−0.046	−0.077*	−0.076*
	(0.068)	(0.068)	(0.068)	(0.042)	(0.042)
Post-65 × Uninsured × High PGS	−0.117*	−0.117*	−0.117*	0.043	0.043
	(0.062)	(0.062)	(0.062)	(0.037)	(0.037)
Shock × Post-65 × Uninsured × High PGS	−0.151	−0.153	−0.154	−0.185	−0.188
	(0.309)	(0.310)	(0.309)	(0.153)	(0.152)
Age FE		Yes	Yes	Yes	Yes
Year FE			Yes		Yes
Individual FE				Yes	Yes
Observations	25.800	25.800	25.800	25.800	25.800

Meaning of OLS coefficients

From estimating equation 1 we get the following: (▸ back)

$$E\left[Y_{it}|post65_{it}=0, g_i=0, shock_{it}=1, uninsured_i=1\right] = \beta + \lambda_2 \tag{9}$$

$$E\left[Y_{it}|post65_{it}=1, g_i=0, shock_{it}=1, uninsured_i=1\right] = \beta + \gamma + \lambda_1 + \lambda_2 \\ + \lambda_3 + \delta_1 \tag{10}$$

$$E\left[Y_{it}|post65_{it}=0, g_i=1, shock_{it}=1, uninsured_i=1\right] = \beta + \lambda_2 + \lambda_4 + \delta_2 \tag{11}$$

$$E\left[Y_{it}|post65_{it}=1, g_i=1, shock_{it}=1, uninsured_i=1\right] = \beta + \gamma + \lambda_1 + \lambda_2 \\ + \lambda_3 + \lambda_4 + \lambda_5 + \delta_1 + \delta_2 \\ + \delta_3 + \delta_4 + \xi \tag{12}$$

Then the first two differences yield:

$$(2) - (1) = \gamma + \lambda_1 + \lambda_3 + \delta_1 \tag{13}$$

$$(4) - (3) = \gamma + \lambda_1 + \lambda_3 + \lambda_5 + \delta_1 + \delta_3 + \delta_4 + \xi \tag{14}$$

And finally the diff-in-diff (G×E) is identified by:

$$(6) - (5) = \lambda_5 + \delta_3 + \delta_4 + \xi \tag{15}$$

Effect of the shock on the outcomes

The derivative of the outcome with respect to shock is: `▸ back`

$$
\begin{aligned}
\frac{\partial Y_{it}}{\partial shock_{it}} = \beta &+ \lambda_1 post65_{it} + \lambda_2 uninsured_i + \lambda_4 g_i \\
&+ \delta_1(post65 \times uninsured_i) + \delta_2(uninsured_i \times g_i) + \delta_3(post65_{it} \times g_i) \\
&+ \xi(post65_{it} \times uninsured_i \times g_i)
\end{aligned}
\tag{16}
$$

Again, we can look at the decomposition:

$$
E\left[\frac{\partial Y_{it}}{\partial shock_{it}} | post65_{it}=0, g_i=0, uninsured_i=1\right] = \beta + \lambda_2
\tag{17}
$$

$$
E\left[\frac{\partial Y_{it}}{\partial shock_{it}} | post65_{it}=1, g_i=0, uninsured_i=1\right] = \beta + \lambda_1 + \lambda_2 + \delta_1
\tag{18}
$$

$$
E\left[\frac{\partial Y_{it}}{\partial shock_{it}} | post65_{it}=0, g_i=1, uninsured_i=1\right] = \beta + \lambda_2 + \lambda_4 + \delta_2
\tag{19}
$$

$$
\begin{aligned}
E\left[\frac{\partial Y_{it}}{\partial shock_{it}} | post65_{it}=1, g_i=1, , uninsured_i=1\right] = \beta &+ \lambda_1 + \lambda_2 + \lambda_4 \\
&+ \delta_1 + \delta_2 + \delta_3 + \xi
\end{aligned}
\tag{20}
$$

Calculating the first two differences as above:

$$
(10) - (9) = \lambda_1 + \delta_1
\tag{21}
$$

$$
(12) - (11) = \lambda_1 + \delta_1 + \delta_3 + \xi
\tag{22}
$$

Outline

Robustness: age range 59-71

Coefficient plot of the main regression.

Robustness: age range 58-72

Coefficient plot of the main regression.

Robustness: age range 55-70

Coefficient plot of the main regression.

Robustness: uninsured only 2/3 of the time

Coefficient plot of the main regression.

Robustness: uninsured only 1/3 of the time

Coefficient plot of the main regression.

Split by cognitive ability PGS

Split by risk-seeking PGS

DNA Sequencing Cost

Cost of sequencing the DNA has been falling rapidly (NIH)

Cost per participant: ≈ 50 \$ (with SNP imputation)

▶ back

References I

Barcellos, S. H., Carvalho, L. S., and Turley, P. (2018).
Education can reduce health differences related to genetic risk of obesity.
Proceedings of the National Academy of Sciences, page 201802909.

Belsky, D. W., Domingue, B. W., Wedow, R., Arseneault, L., and Boardman, J. D. (2018).
Genetic analysis of social-class mobility in five longitudinal studies.
Proceedings of the National Academy of Sciences, pages 1–10.

Benjamin, D. J., Caplin, A., Cesarini, D., Thom, K., and Turley, P. (2016).
Smoking, Genes, and Health.
Working Paper, pages 1–27.

Benjamin, D. J., Chabris, C. F., Glaeser, E. L., Gudnason, V., Harris, T. B., Laibson, D. I., Launer, L. J., and Purcell, S. M. (2007).
Genoeconomics.
In Weinstein, M., Vaupel, J. W., and Wachter, K. W., editors, Biosocial Surveys, chapter 15, pages 304–335. The National Academies Press, Washington, D.C.

Card, D., Dobkin, C., and Maestas, N. (2008).
The impact of nearly universal insurance coverage on health care utilization: evidence from medicare.
American Economic Review, 98(5):2242–58.

Card, D., Dobkin, C., and Maestas, N. (2009).
Does Medicare Save Lives?
Quarterly Journal of Economics, 124(2):597–636.

Caspi, A., McClay, J., Moffitt, T. E., Mill, J., Martin, J., Craig, I. W., Taylor, A., and Poulton, R. (2002).
Role of genotype in the cycle of violence in maltreated children.
Science, 297(5582):851–4.

References II

Cesarini, D., Dawes, C. T., Johannesson, M., Lichtenstein, P., and Wallace, B. (2009).
Genetic Variation in Preferences for Giving and Risk Taking.
Quarterly Journal of Economics, 124(2):809–842.

Clark, A. and Etilé, F. (2002).
Do health changes affect smoking? Evidence from British panel data.
Journal of health economics, 21(4):533–62.

Einav, L. and Finkelstein, A. (2018).
Moral Hazard in Health Insurance: What We Know and How We Know It.
Journal of the European Economic Association, 16(4):957–982.

Einav, L., Finkelstein, A., Ryan, S. P., Schrimpf, P., and Cullen, M. R. (2013).
Selection on Moral Hazard in Health Insurance.
American Economic Review, 103(1):178–219.

Falba, T. (2005).
Health events and the smoking cessation of middle aged Americans.
Journal of behavioral medicine, 28(1):21–33.

Goodchild, M., Nargis, N., and d'Espaignet, E. (2017).
Global economic cost of smoking-attributable diseases.
Tobacco Control.

Keenan, P. S. (2009).
Smoking and Weight Change After New Health Diagnoses in Older Adults.
Archives of Internal Medicine, 169(3):237.

References III

Khwaja, A., Sloan, F., and Chung, S. (2006).
Learning about individual risk and the decision to smoke.
International Journal of Industrial Organization, 24(4):683–699.

Liu, M., Jiang, Y., Wedow, R., Li, Y., Brazel, D. M., Chen, F., Datta, G., Davila-Velderrain, J., McGuire, D., Tian, C., Zhan, X., Choquet, H., Docherty, A. R., Faul, J. D., Foerster, J. R., Fritsche, L. G., Gabrielsen, M. E., Gordon, S. D., Haessler, J., Hottenga, J.-J., Huang, H., Jang, S.-K., Jansen, P. R., Ling, Y., Mägi, R., Matoba, N., McMahon, G., Mulas, A., Orrù, V., Palviainen, T., Pandit, A., Reginsson, G. W., Skogholt, A. H., Smith, J. A., Taylor, A. E., Turman, C., Willemsen, G., Young, H., Young, K. A., Zajac, G. J. M., Zhao, W., Zhou, W., Bjornsdottir, G., Boardman, J. D., Boehnke, M., Boomsma, D. I., Chen, C., Cucca, F., Davies, G. E., Eaton, C. B., Ehringer, M. A., Esko, T., Fiorillo, E., Gillespie, N. A., Gudbjartsson, D. F., Haller, T., Harris, K. M., Heath, A. C., Hewitt, J. K., Hickie, I. B., Hokanson, J. E., Hopfer, C. J., Hunter, D. J., Iacono, W. G., Johnson, E. O., Kamatani, Y., Kardia, S. L. R., Keller, M. C., Kellis, M., Kooperberg, C., Kraft, P., Krauter, K. S., Laakso, M., Lind, P. A., Loukola, A., Lutz, S. M., Madden, P. A. F., Martin, N. G., McGue, M., McQueen, M. B., Medland, S. E., Metspalu, A., Mohlke, K. L., Nielsen, J. B., Okada, Y., Peters, U., Polderman, T. J. C., Posthuma, D., Reiner, A. P., Rice, J. P., Rimm, E., Rose, R. J., Runarsdottir, V., Stallings, M. C., Stančáková, A., Stefansson, H., Thai, K. K., Tindle, H. A., Tyrfingsson, T., Wall, T. L., Weir, D. R., Weisner, C., Whitfield, J. B., Winsvold, B. S., Yin, J., Zuccolo, L., Bierut, L. J., Hveem, K., Lee, J. J., Munafò, M. R., Saccone, N. L., Willer, C. J., Cornelis, M. C., David, S. P., Hinds, D. A., Jorgenson, E., Kaprio, J., Stitzel, J. A., Stefansson, K., Thorgeirsson, T. E., Abecasis, G., Liu, D. J., and Vrieze, S. (2019).
Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use.
Nature Genetics.

Ma, J., Siegel, R. L., Jacobs, E. J., and Jemal, A. (2018).
Smoking-attributable Mortality by State in 2014, U.S.
American Journal of Preventive Medicine, 54(5):661–670.

Marti, J. and Richards, M. R. (2017).
Smoking Response to Health and Medical Spending Changes and the Role of Insurance.
Health Economics, 26:305–320.

Pei, Z., Pischke, J. S., and Schwandt, H. (2018).
Poorly Measured Confounders are More Useful on the Left than on the Right.
Journal of Business and Economic Statistics, pages 1–12.

References IV

Richards, M. R. and Marti, J. (2014).
Heterogeneity in the smoking response to health shocks by out-of-pocket spending risk.
Health Economics, Policy and Law, 9(04):343–357.

Rimfeld, K., Krapohl, E., Trzaskowski, M., Coleman, J. R. I., Selzam, S., Dale, P. S., Esko, T., Metspalu, A., and Plomin, R. (2018).
Genetic influence on social outcomes during and after the Soviet era in Estonia.
Nature Human Behaviour, 2(4):269.

Rosenquist, J. N., Lehrer, S. F., O'Malley, A. J., Zaslavsky, A. M., Smoller, J. W., and Christakis, N. A. (2015).
Cohort of birth modifies the association between FTO genotype and BMI.
Proceedings of the National Academy of Sciences, 112(2):354–9.

Rutter, M. (2006).
Genes and Behavior: Nature-Nurture Interplay Explained.
Blackwell Publishers, Oxford, UK.

Schmitz, L. L. and Conley, D. C. (2017a).
Modeling Gene-Environment Interactions With Quasi-Natural Experiments.
Journal of Personality, 85(1):10–21.

Schmitz, L. L. and Conley, D. C. (2017b).
The effect of Vietnam-era conscription and genetic potential for educational attainment on schooling outcomes.
Economics of Education Review, 61:85–97.

Sundmacher, L. (2012).
The effect of health shocks on smoking and obesity.
The European Journal of Health Economics, 13(4):451–460.

References V

The 1000 Genomes Project Consortium, Auton, A., Abecasis, G. R., ..., Dunn, M., and Schloss, J. A. (2015).
A global reference for human genetic variation.
Nature, 526(7571):68–74.

The Tobacco and Genetics Consortium, Furberg, H., Kim, Y., Dackor, J., Boerwinkle, E., Franceschini, N., Ardissino, D., Bernardinelli, L., Mannucci, P. M., Mauri, F., Merlini, P. A., Absher, D., Assimes, T. L., Fortmann, S. P., Iribarren, C., Knowles, J. W., Quertermous, T., Ferrucci, L., Tanaka, T., Bis, J. C., Furberg, C. D., Haritunians, T., McKnight, B., Psaty, B. M., Taylor, K. D., Thacker, E. L., Almgren, P., Groop, L. C., Ladenvall, C., Boehnke, M., Jackson, A. U., Mohlke, K. L., Stringham, H. M., Tuomilehto, J., Benjamin, E. J., Hwang, S.-J., Levy, D., Preis, S. R., Vasan, R. S., Duan, J., Gejman, P. V., Levinson, D. F., Sanders, A. R., Shi, J., Lips, E. H., McKay, J. D., Agudo, A., Barzan, L., Bencko, V., Benhamou, S., Castellsagué, X., Canova, C., Conway, D. I., Fabianova, E., Foretova, L., Janout, V., Healy, C. M., Holcátová, I., Kjaerheim, K., Lagiou, P., Lissowska, J., Lowry, R., Macfarlane, T. V., Mates, D., Richiardi, L., Rudnai, P., Szeszenia-Dabrowska, N., Zaridze, D., Znaor, A., Lathrop, M., Brennan, P., Bandinelli, S., Frayling, T. M., Guralnik, J. M., Milaneschi, Y., Perry, J. R. B., Altshuler, D., Elosua, R., Kathiresan, S., Lucas, G., Melander, O., O'Donnell, C. J., Salomaa, V., Schwartz, S. M., Voight, B. F., Penninx, B. W. J. H., Smit, J. H., Vogelzangs, N., Boomsma, D. I., de Geus, E. J. C., Vink, J. M., Willemsen, G., Chanock, S. J., Gu, F., Hankinson, S. E., Hunter, D. J., Hofman, A., Tiemeier, H., Uitterlinden, A. G., van Duijn, C. M., Walter, S., Chasman, D. I., Everett, B. M., Paré, G., Ridker, P. M., Li, M. D., Maes, H. H., Audrain-McGovern, J., Posthuma, D., Thornton, L. M., Lerman, C., Kaprio, J., Rose, J. E., Ioannidis, J. P. A., Kraft, P., Lin, D.-Y., and Sullivan, P. F. (2010).
Genome-wide meta-analyses identify multiple loci associated with smoking behavior.
Nature Genetics, 42(5):441–447.

United States Department of Health and Human Services (2014).
The Health Consequences of Smoking–50 Years of Progress A Report of the Surgeon General.
The Health Consequences of Smoking-50 Years of Progress A Report of The Surgeon General.

Xu, X., Bishop, E. E., Kennedy, S. M., Simpson, S. A., and Pechacek, T. F. (2015).
Annual Healthcare Spending Attributable to Cigarette Smoking.
American Journal of Preventive Medicine, 48(3):326–333.