

# Collapsing Earth

Mike Geipel

April 29, 2021

## 1 What if the Earth collapses to a black hole, instantaneously?

### 1.1 Premise

You're standing on the Earth, when it suddenly collapses to a black hole. Do you fall into the resulting black hole, the size of a peanut?

### 1.2 Assumptions

You don't die immediately from this mysterious event, so you can experience your fate. This means:

- This is a non-violent event: no gamma ray burst or accretion disk spewing sterilizing x-rays.
- Ergo, the atmosphere and any resulting debris are inside the black hole. No accretion disk.
- You're wearing a space suit. (Since there's no atmosphere.)
- The resulting black hole has the same mass and angular momentum as the Earth had had.

Also, assume that you're on the equator at the time, since values like radius and rotational speed are easy to find. (And this is the particular scenario described in a podcast.)

"r" is distance between the astronaut and the black hole.

Equation for speed, given semimajor axis ("a"):

$$v = \sqrt{\mu \left( \frac{2}{r} - \frac{1}{a} \right)}$$

Apogee initial conditions (r,v):

where r = Earth radius at equator, and v = Earth rotation speed at equator.

Now, solve for the length of semimajor axis:

$$a = \frac{\mu r}{2\mu - rv^2}$$

Equation for period of an orbit, given semimajor axis:

$$T = 2\pi \sqrt{\frac{a^3}{\mu}}$$

Schwarzschild radius:

$$R_g = \frac{2\mu}{c^2}$$

semimajor axis(a):	3194597 meters	(or 3194.597 km)
orbital period(T):	1797 secs	(or 29.95 mins)
perigee (closest):	11058 meters	(or 11.058 km)

perigee (fastest):            268272 m/s            (or 268.272 km/s)  
blackhole radius :            0.0089 meters        (or 8.9 millimeters)

### 1.3 Tidal Forces

Assume that you are two meters tall in your space suit. The worst case is if you're oriented along the semimajor axis at perigee.

$$\alpha = \frac{\mu}{r_2^2} - \frac{\mu}{r_1^2}$$

tidal acceleration:            1179 m/s/s            (or 120 g)  
This is the difference between head and foot; each end experiences half that.

### 1.4 Conclusion

You'll not just fall into the peanut-sized black hole. The closest you'll get is about 11 kilometers. But – you'd be moving at almost 270 km/s, while turning the corner very, VERY, quickly. This is a highly eccentric orbit.



So, assuming you're 2m tall: the tidal difference between head and foot could be about 120g. Ouch! However!!! In the worst case scenario, you would experience these extreme accelerations for only a fraction of a second. It might be survivable, like experiencing a shock wave on Earth.

In fact, it's so quick that there wouldn't be time for the gravity gradient to align you with "down". So, if your space suit has a jetpack, your best use of it would be to make sure you were not oriented along the semimajor axis. (Even so, you're likely to come out of each close pass spinning.) Unfortunately, you would need to deal with this every 30 mins or so – until you get rescued.

## 2 Diagrams of the Highly Eccentric Orbit

$$b = \sqrt{a^2 - c_f^2}$$

semiminor axis(b):            265570 meters        (or 265.570 km)



### 3 Author's Notes

#### 3.1 The astronaut is a pendulum.

My intuition tells me that our astronaut is likely to be spun around by tidal forces every time she encounters perigee, depending on orientation.

But I'm not yet sure how to evaluate the **likely** vs **worst-case** spin scenarios.

#### 3.2 GR effects were not considered.

I have not yet looked into how to evaluate the effect of the following on subsequent orbits:

- GR orbital precession
- frame-dragging (due to Earth's angular momentum)

This would be really cool to investigate. I'd need to do some research for follow-up.

Given the magnitude for change in size, the black hole would likely be spinning very fast, indeed.

#### 3.3 The Moon's effect may not be negligible.

Everything described above only considers the **Earth!!** But the Earth and the Moon actually **both** orbit about the barycenter (combined center of mass).

If all of the Earth's mass collapsed into its equivalent-mass black hole, this orbital dynamic would not change. The resulting black hole would trace out a circle every 28 days with 4700 km radius.

In the 15 minutes that it would take for our astronaut to fall towards the black hole, the black hole will have been displaced by almost **11 km!!**

This would not likely change the orbit analysis by much since most of the acceleration takes place over a tiny time frame, but it would complicate any calculation of orbital precession, particularly if we consider latitudes away from the equator.

#### 3.4 Final surprise: black hole spin

So: how fast could the black hole be spinning? (Our previous analysis did not need this info.)

According to [https://en.wikipedia.org/wiki/Moment\\_of\\_inertia\\_factor](https://en.wikipedia.org/wiki/Moment_of_inertia_factor), the Earth's moment of inertia factor is about 0.3307. I have no idea what this factor would be for a black hole; it's just a constant for different mass distributions.

angular moment of inertia:  $I = (0.3307) M R^2$ , (for Earth)

angular momentum:  $L = I \omega$ , (where  $\omega$  is angular velocity)

So roughly:  $\omega \propto \frac{1}{R^2}$ , ignoring relativistic effects

As the Earth's radius ( $R$ ) shrinks:  $6.378\text{e}+06 \mapsto 0.0089$  meters,

the corresponding tangent velocity grows:  $465.1 \mapsto 2.405\text{e}+20$  m/s.

That's orders of magnitude faster than the speed of light ( $c$ ):  $2.998\text{e}+08$  m/s!

### 3.4.1 Getting the answer

I suppose such calculations could be handled with relativistic equations to get a precise figure, where momentum (mass) must rise as the speed gets asymptotically closer to the speed of light.

But for our purposes, we don't need this: we already have a good enough answer for such extreme values: **our black hole is spinning at *nearly* the speed of light.**

If we don't relax the assumption (on conservation of Earth's rotational angular momentum), the Ergosphere and frame-dragging effects could be very interesting.

### 3.4.2 Follow-up questions:

- How big is the Ergosphere?
- How could frame-dragging affect the tidal force felt by our astronaut?
- Would frame-dragging mean our astronaut could be kicked into a significantly higher orbit? If so, she could use her jetpack (thrusting away from the black hole) to gain orbital angular momentum at perigee, and then wait for the next apogee (now farther out) to improve orbit eccentricity, using her remaining  $\Delta v$  more efficiently.