

Outline

- 1 Markov chain Monte Carlo
- 2 Metropolis-Hastings algorithms
- 3 Gibbs sampling

Markov chain Monte Carlo

Markov chain Monte Carlo

- Assume now simulating from $\pi(\mathbf{x})$ is difficult directly
 - $\pi(\cdot)$ complicated
 - \mathbf{x} high-dimensional

Markov chain Monte Carlo

- Assume now simulating from $\pi(\mathbf{x})$ is difficult directly
 - $\pi(\cdot)$ complicated
 - \mathbf{x} high-dimensional
- Markov chain Monte Carlo:
 - Generates $\{\mathbf{x}^{(t)}\}$ **sequentially**
 - Markov structure: $\mathbf{x}^{(t)} \sim P(\cdot | \mathbf{x}^{(t-1)})$

Markov chain Monte Carlo

- Assume now simulating from $\pi(\mathbf{x})$ is difficult directly
 - $\pi(\cdot)$ complicated
 - \mathbf{x} high-dimensional
- Markov chain Monte Carlo:
 - Generates $\{\mathbf{x}^{(t)}\}$ **sequentially**
 - Markov structure: $\mathbf{x}^{(t)} \sim P(\cdot | \mathbf{x}^{(t-1)})$
- Aim now:
 - The distribution of $\mathbf{x}^{(t)}$ **converges** to $\pi(\cdot)$ as t increases
 - $\hat{\mu}_{MCMC} = N^{-1} \sum_{t=1}^N h(\mathbf{x}^{(t)})$ **converges** towards $\mu = E^{\pi}[h(\mathbf{x})]$ as t increases
- References:
 - **Givens and Hoeting (2012)**: *Computational statistics*
 - **Robert and Casella (1999)**: *Monte Carlo statistical methods*

Markov chain theory - discrete case

- Assume $\{X^{(t)}\}$ is a **Markov chain** where $X^{(t)}$ is a **discrete** random variable

$$\Pr(X^{(t)} = \mathbf{x}^* | X^{(t-1)} = \mathbf{x}) = P(\mathbf{x}^* | \mathbf{x}) \quad \text{transition probabilities}$$

Markov chain theory - discrete case

- Assume $\{X^{(t)}\}$ is a **Markov chain** where $X^{(t)}$ is a **discrete** random variable

$$\Pr(X^{(t)} = \mathbf{x}^* | X^{(t-1)} = \mathbf{x}) = P(\mathbf{x}^* | \mathbf{x}) \quad \text{transition probabilities}$$

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any state infinitely often.
 - aperiodic**: Does not go in cycles

Markov chain theory - discrete case

- Assume $\{X^{(t)}\}$ is a **Markov chain** where $X^{(t)}$ is a **discrete** random variable

$$\Pr(X^{(t)} = \mathbf{x}^* | X^{(t-1)} = \mathbf{x}) = P(\mathbf{x}^* | \mathbf{x}) \quad \text{transition probabilities}$$

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any state infinitely often.
 - aperiodic**: Does not go in cycles
- Then there exists a **unique** distribution $\pi(\mathbf{x})$ such that

$$\lim_{T \rightarrow \infty} \Pr(X^{(T)} = \mathbf{x}^* | X^{(0)} = \mathbf{x}) = \pi(\mathbf{x}^*)$$

$$\lim_{T \rightarrow \infty} \hat{\mu}_{MCMC} = \mu = E^{\pi}[\mathbf{h}(X)]$$

$$\hat{\mu}_{MCMC} = \frac{1}{T} \sum_{t=1}^T \mathbf{h}(\mathbf{x}^{(t)})$$

Markov chain theory - discrete case

- Assume $\{X^{(t)}\}$ is a **Markov chain** where $X^{(t)}$ is a **discrete** random variable

$$\Pr(X^{(t)} = \mathbf{x}^* | X^{(t-1)} = \mathbf{x}) = P(\mathbf{x}^* | \mathbf{x}) \quad \text{transition probabilities}$$

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any state infinitely often.
 - aperiodic**: Does not go in cycles
- Then there exists a **unique** distribution $\pi(\mathbf{x})$ such that

$$\lim_{T \rightarrow \infty} \Pr(X^{(T)} = \mathbf{x}^* | X^{(0)} = \mathbf{x}) = \pi(\mathbf{x}^*)$$

$$\lim_{T \rightarrow \infty} \hat{\mu}_{MCMC} = \mu = E^\pi[h(\mathbf{X})] \quad \hat{\mu}_{MCMC} = \frac{1}{T} \sum_{t=1}^T h(\mathbf{x}^{(t)})$$

- How to find $\pi(\cdot)$ (the **stationary** distribution): Solve

$$\pi(\mathbf{x}^*) = \sum_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^* | \mathbf{x})$$

Markov chain theory - discrete case

- Assume $\{X^{(t)}\}$ is a **Markov chain** where $X^{(t)}$ is a **discrete** random variable

$$\Pr(X^{(t)} = \mathbf{x}^* | X^{(t-1)} = \mathbf{x}) = P(\mathbf{x}^* | \mathbf{x}) \quad \text{transition probabilities}$$

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - recurrent**: The chain will visit any state infinitely often.
 - aperiodic**: Does not go in cycles
- Then there exists a **unique** distribution $\pi(\mathbf{x})$ such that

$$\lim_{T \rightarrow \infty} \Pr(X^{(T)} = \mathbf{x}^* | X^{(0)} = \mathbf{x}) = \pi(\mathbf{x}^*)$$

$$\lim_{T \rightarrow \infty} \hat{\mu}_{MCMC} = \mu = E^\pi[h(\mathbf{X})] \quad \hat{\mu}_{MCMC} = \frac{1}{T} \sum_{t=1}^T h(\mathbf{x}^{(t)})$$

- How to find $\pi(\cdot)$ (the **stationary** distribution): Solve

$$\pi(\mathbf{x}^*) = \sum_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^* | \mathbf{x})$$

- Our situation**: We have $\pi(\mathbf{x})$, want to find $P(\mathbf{x}^* | \mathbf{x})$
 - Note: **Many** possible $P(\mathbf{x}^* | \mathbf{x})$!

Markov chain theory - general setting

- Assume $\{\mathbf{x}^{(t)}\}$ is a **Markov chain** where $\mathbf{x}^{(t)} \in S$

$$\Pr(\mathbf{x}^{(t)} \in A | \mathbf{x}^{(t-1)} = \mathbf{x}) = P(\mathbf{x}, A) = \int_{\mathbf{x}^* \in A} P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}^*$$

giving the **transition densities**

Markov chain theory - general setting

- Assume $\{\mathbf{x}^{(t)}\}$ is a **Markov chain** where $\mathbf{x}^{(t)} \in S$

$$\Pr(\mathbf{x}^{(t)} \in A | \mathbf{x}^{(t-1)} = \mathbf{x}) = P(\mathbf{x}, A) = \int_{\mathbf{x}^* \in A} P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}^*$$

giving the **transition densities**

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any $A \subset S$ infinitely often.
 - aperiodic**: Do not go in cycles

Markov chain theory - general setting

- Assume $\{\mathbf{x}^{(t)}\}$ is a **Markov chain** where $\mathbf{x}^{(t)} \in S$

$$\Pr(\mathbf{x}^{(t)} \in A | \mathbf{x}^{(t-1)} = \mathbf{x}) = P(\mathbf{x}, A) = \int_{\mathbf{x}^* \in A} P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}^*$$

giving the **transition densities**

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any $A \subset S$ infinitely often.
 - aperiodic**: Do not go in cycles
- Then there exists a distribution $\pi(\mathbf{x})$ such that

$$\lim_{T \rightarrow \infty} \Pr(\mathbf{x}^{(T)} \in A | \mathbf{x}^{(0)} = \mathbf{x}) = \int_A \pi(\mathbf{x}^*) d\mathbf{x}^*$$

$$\hat{\mu}_{MCMC} \rightarrow \mu$$

Markov chain theory - general setting

- Assume $\{\mathbf{x}^{(t)}\}$ is a **Markov chain** where $\mathbf{x}^{(t)} \in S$

$$\Pr(\mathbf{x}^{(t)} \in A | \mathbf{x}^{(t-1)} = \mathbf{x}) = P(\mathbf{x}, A) = \int_{\mathbf{x}^* \in A} P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}^*$$

giving the **transition densities**

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any $A \subset S$ infinitely often.
 - aperiodic**: Do not go in cycles
- Then there exists a distribution $\pi(\mathbf{x})$ such that

$$\lim_{T \rightarrow \infty} \Pr(\mathbf{x}^{(T)} \in A | \mathbf{x}^{(0)} = \mathbf{x}) = \int_A \pi(\mathbf{x}^*) d\mathbf{x}^*$$

$$\hat{\mu}_{MCMC} \rightarrow \mu$$

- How to find $p(\cdot)$ (the **stationary** distribution): Solve

$$\pi(\mathbf{x}^*) = \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}$$

Markov chain theory - general setting

- Assume $\{\mathbf{x}^{(t)}\}$ is a **Markov chain** where $\mathbf{x}^{(t)} \in S$

$$\Pr(\mathbf{x}^{(t)} \in A | \mathbf{x}^{(t-1)} = \mathbf{x}) = P(\mathbf{x}, A) = \int_{\mathbf{x}^* \in A} P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}^*$$

giving the **transition densities**

- Assume the chain is
 - irreducible**: Possible to move from any \mathbf{x} to any \mathbf{x}^* in a finite number of steps
 - reccurent**: The chain will visit any $A \subset S$ infinitely often.
 - aperiodic**: Do not go in cycles
- Then there exists a distribution $\pi(\mathbf{x})$ such that

$$\lim_{T \rightarrow \infty} \Pr(\mathbf{x}^{(T)} \in A | \mathbf{x}^{(0)} = \mathbf{x}) = \int_A \pi(\mathbf{x}^*) d\mathbf{x}^*$$

$$\hat{\mu}_{MCMC} \rightarrow \mu$$

- How to find $p(\cdot)$ (the **stationary** distribution): Solve

$$\pi(\mathbf{x}^*) = \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^* | \mathbf{x}) d\mathbf{x}$$

- Our situation**: We have $\pi(\cdot)$, want to find $P(\mathbf{x}^* | \mathbf{x})$

Detailed balance

- The task: Find a transition probability/density $P(\mathbf{x}^*|\mathbf{x})$ satisfying

$$\pi(\mathbf{x}^*) = \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) d\mathbf{x}$$

Can in general be a difficult criterion to check

Detailed balance

- The task: Find a transition probability/density $P(\mathbf{x}^*|\mathbf{x})$ satisfying

$$\pi(\mathbf{x}^*) = \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) d\mathbf{x}$$

Can in general be a difficult criterion to check

- **Sufficient** criterion:

$$\pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) = \pi(\mathbf{x}^*) P(\mathbf{x}|\mathbf{x}^*) \quad \text{Detailed balance}$$

Detailed balance

- The task: Find a transition probability/density $P(\mathbf{x}^*|\mathbf{x})$ satisfying

$$\pi(\mathbf{x}^*) = \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) d\mathbf{x}$$

Can in general be a difficult criterion to check

- **Sufficient** criterion:

$$\pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) = \pi(\mathbf{x}^*) P(\mathbf{x}|\mathbf{x}^*) \quad \text{Detailed balance}$$

We then have

$$\begin{aligned} \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) d\mathbf{x} &= \int_{\mathbf{x}} \pi(\mathbf{x}^*) P(\mathbf{x}|\mathbf{x}^*) d\mathbf{x} \\ &= \pi(\mathbf{x}^*) \int_{\mathbf{x}} P(\mathbf{x}|\mathbf{x}^*) d\mathbf{x} = \pi(\mathbf{x}^*) \end{aligned}$$

since $P(\mathbf{x}|\mathbf{x}^*)$ is, for any given \mathbf{x}^* , a density wrt \mathbf{x} .

Detailed balance

- The task: Find a transition probability/density $P(\mathbf{x}^*|\mathbf{x})$ satisfying

$$\pi(\mathbf{x}^*) = \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) d\mathbf{x}$$

Can in general be a difficult criterion to check

- **Sufficient** criterion:

$$\pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) = \pi(\mathbf{x}^*) P(\mathbf{x}|\mathbf{x}^*) \quad \text{Detailed balance}$$

We then have

$$\begin{aligned} \int_{\mathbf{x}} \pi(\mathbf{x}) P(\mathbf{x}^*|\mathbf{x}) d\mathbf{x} &= \int_{\mathbf{x}} \pi(\mathbf{x}^*) P(\mathbf{x}|\mathbf{x}^*) d\mathbf{x} \\ &= \pi(\mathbf{x}^*) \int_{\mathbf{x}} P(\mathbf{x}|\mathbf{x}^*) d\mathbf{x} = \pi(\mathbf{x}^*) \end{aligned}$$

since $P(\mathbf{x}|\mathbf{x}^*)$ is, for any given \mathbf{x}^* , a density wrt \mathbf{x} .

- **Question:** What would be the ideal choice of $P(\mathbf{x}^*|\mathbf{x})$?
- **Question:** Why only necessary to check for $\mathbf{x}^* \neq \mathbf{x}$.
- **Question:** Why do you think it is called detailed **balance**?

Metropolis-Hastings algorithms

Metropolis-Hastings algorithms

- $P(\mathbf{x}^*|\mathbf{x})$ defined through an algorithm:

- 1 Sample a candidate value \mathbf{x}^* from a **proposal distribution** $g(\cdot|\mathbf{x})$.
- 2 Compute the Metropolis-Hastings ratio

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})}$$

- 3 Put

$$\mathbf{Y} = \begin{cases} \mathbf{x}^* & \text{with probability } \min\{1, R(\mathbf{x}, \mathbf{x}^*)\} \\ \mathbf{x} & \text{otherwise} \end{cases}$$

Metropolis-Hastings algorithms

- $P(\mathbf{x}^*|\mathbf{x})$ defined through an algorithm:

- 1 Sample a candidate value \mathbf{x}^* from a **proposal distribution** $g(\cdot|\mathbf{x})$.
- 2 Compute the Metropolis-Hastings ratio

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})}$$

- 3 Put

$$\mathbf{Y} = \begin{cases} \mathbf{x}^* & \text{with probability } \min\{1, R(\mathbf{x}, \mathbf{x}^*)\} \\ \mathbf{x} & \text{otherwise} \end{cases}$$

- For $\mathbf{x}^* \neq \mathbf{x}$:

$$P(\mathbf{x}^*|\mathbf{x}) = g(\mathbf{x}^*|\mathbf{x}) \min \left\{ 1, \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} \right\}$$

- Note: $P(\mathbf{x}|\mathbf{x})$ somewhat difficult to evaluate in this case.
- Detailed balance fulfilled?

$$\begin{aligned} \pi(\mathbf{x})P(\mathbf{x}^*|\mathbf{x}) &= \pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x}) \min \left\{ 1, \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} \right\} \\ &= \min\{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x}), \pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)\} \\ &= \pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*) \min \left\{ \frac{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})}{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}, 1 \right\} = \pi(\mathbf{x}^*)P(\mathbf{x}|\mathbf{x}^*) \end{aligned}$$

Random walk chains

- Popular choice of proposal distribution:

$$\mathbf{x}^* = \mathbf{x} + \boldsymbol{\epsilon}$$

- $g(\mathbf{x}^*|\mathbf{x}) = h(\mathbf{x}^* - \mathbf{x})$
- Popular choices: Uniform, Gaussian, t -distribution
- Note: If $g(\cdot)$ is symmetric, $g(\mathbf{x}^*|\mathbf{x}) = g(\mathbf{x}|\mathbf{x}^*)$ and

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} = \frac{\pi(\mathbf{x}^*)}{\pi(\mathbf{x})}$$

- Results in the original Metropolis algorithm ([Metropolis et al., 1953](#))
- Extended by [Hastings \(1970\)](#) for non-symmetric (\cdot) to the now-called Metropolis-Hastings algorithm

Independence chains

- Assume $g(\mathbf{x}^*|\mathbf{x}) = g(\mathbf{x}^*)$. Then

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x})}{\pi(\mathbf{x})g(\mathbf{x}^*)} = \frac{\frac{\pi(\mathbf{x}^*)}{g(\mathbf{x}^*)}}{\frac{\pi(\mathbf{x})}{g(\mathbf{x})}},$$

fraction of **importance weights**!

- Behave very much like importance sampling and SIR
- Difficult to specify $g(\mathbf{x})$ for high-dimensional problems
- Theoretical properties easier to evaluate than for random walk versions.




Question: For independence chains, acceptance rates should be as large as possible, for random walk it should be in the range [0.2,0.7]. Why this difference?

Example

- Assume $p(x) \propto \exp(-|x|^3/3)$
- Proposal distribution $N(x, \sigma_p^2)$
- `Example_MH_cubic.R`
- $N = 10\,000$ samples, varying values of σ_p^2




Example - $\sigma_p = 10$

Acceptance rate = 0.13






Example - $\sigma_p = 1$

Acceptance rate = 0.72



Example - $\sigma_p = 4$

Acceptance rate = 0.26



M-H and unknown constant

- Assume now $\pi(\mathbf{x}) = c \cdot q(\mathbf{x})$ with c unknown.

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} = \frac{c \cdot q(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{c \cdot q(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} = \frac{q(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{q(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})}$$

- Do not depend on c !

M-H and multivariate settings

- $\mathbf{x} = (X_1, \dots, X_p)$
- Typical in this case: Only change **one** or a few components at a time.
 - 1 Choose index j (randomly)
 - 2 Sample $X_j^* \sim g_j(\cdot|\mathbf{x})$, put $X_k^* = X_k$ for $k \neq j$
 - 3 Compute

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} = \frac{\pi(x_j^*|\mathbf{x}_{-j})g_j(x_j|\mathbf{x}^*)}{\pi(x_j|\mathbf{x}_{-j})g_j(x_j^*|\mathbf{x})}$$

- 4 Put

$$\mathbf{y} = \begin{cases} \mathbf{x}^* & \text{with probability } \min\{1, R(\mathbf{x}, \mathbf{x}^*)\} \\ \mathbf{x} & \text{otherwise} \end{cases}$$

M-H and multivariate settings

- $\mathbf{x} = (X_1, \dots, X_p)$
- Typical in this case: Only change **one** or a few components at a time.
 - 1 Choose index j (randomly)
 - 2 Sample $X_j^* \sim g_j(\cdot|\mathbf{x})$, put $X_k^* = X_k$ for $k \neq j$
 - 3 Compute

$$R(\mathbf{x}, \mathbf{x}^*) = \frac{\pi(\mathbf{x}^*)g(\mathbf{x}|\mathbf{x}^*)}{\pi(\mathbf{x})g(\mathbf{x}^*|\mathbf{x})} = \frac{\pi(x_j^*|\mathbf{x}_{-j})g_j(x_j|\mathbf{x}^*)}{\pi(x_j|\mathbf{x}_{-j})g_j(x_j^*|\mathbf{x})}$$

- 4 Put

$$\mathbf{y} = \begin{cases} \mathbf{x}^* & \text{with probability } \min\{1, R(\mathbf{x}, \mathbf{x}^*)\} \\ \mathbf{x} & \text{otherwise} \end{cases}$$

- Can show that this version also satisfies detailed balance
- Can even go through indexes systematic
 - Should then consider the whole loop through all components as one iteration

Example

- Assume $\pi(\mathbf{x}) \propto \exp(-\|\mathbf{x}\|^3/3) = \exp(-[\|\mathbf{x}\|^2]^{3/2}/3) =$
- Algorithm given current value \mathbf{x}^t :



- 1: Sample $j \in \{1, \dots, p\}$
- 2: Put

$$x_k^* = \begin{cases} x_k^t & \text{if } k \neq j \\ N(x_k^t, \sigma_k^2) & \text{if } k = j \end{cases}$$

- 3: Calculate $R(\mathbf{x}^t, \mathbf{x}^*) = \pi(\mathbf{x}^*)/\pi(\mathbf{x}^t)$
- 4: Generate $u \sim \text{Uniform}[0, 1]$
- 5: **if** $u < R(\mathbf{x}, \mathbf{x}^*)$ **then**
- 6: Put $\mathbf{x}^{t+1} = \mathbf{x}^*$
- 7: **else**
- 8: Put $\mathbf{x}^{t+1} = \mathbf{x}^t$
- 9: **end if**

Example_MH_cubic_multivariate.R

Results multivariate cubic



Acceptance rate: 0.30

Gibbs sampling

- Assume $\mathbf{x} = (X_1, \dots, X_p)$
- Aim: Simulate $\mathbf{x} \sim \pi(\mathbf{x})$
- Gibbs sampling:
 - 1 Select starting values $\mathbf{x}^{(0)}$ and set $t = 0$
 - 2 Generate, in turn

$$X_1^{(t+1)} \sim p(x_1 | x_2^{(t)}, x_3^{(t)}, \dots, x_p^{(t)})$$

$$X_2^{(t+1)} \sim p(x_2 | x_1^{(t+1)}, x_3^{(t)}, \dots, x_p^{(t)})$$

$$\vdots$$

$$X_{p-1}^{(t+1)} \sim p(x_{p-1} | x_1^{(t+1)}, \dots, x_{p-2}^{(t+1)}, x_p^{(t)})$$

$$X_p^{(t+1)} \sim p(x_p | x_1^{(t+1)}, \dots, x_{p-1}^{(t+1)})$$

Gibbs sampling

- Assume $\mathbf{x} = (X_1, \dots, X_p)$
- Aim: Simulate $\mathbf{x} \sim \pi(\mathbf{x})$
- Gibbs sampling:
 - 1 Select starting values $\mathbf{x}^{(0)}$ and set $t = 0$
 - 2 Generate, in turn

$$X_1^{(t+1)} \sim p(x_1 | x_2^{(t)}, x_3^{(t)}, \dots, x_p^{(t)})$$

$$X_2^{(t+1)} \sim p(x_2 | x_1^{(t+1)}, x_3^{(t)}, \dots, x_p^{(t)})$$

$$\vdots$$

$$X_{p-1}^{(t+1)} \sim p(x_{p-1} | x_1^{(t+1)}, \dots, x_{p-2}^{(t+1)}, x_p^{(t)})$$


$$X_p^{(t+1)} \sim p(x_p | x_1^{(t+1)}, \dots, x_{p-1}^{(t+1)})$$

- 3 Increment t and go to step 2.
- Completion of step 2 is called a **cycle**

Example - Ising model

- Ising model: $x_i \in \{-1, 1\}$, i lattice

$$p(\mathbf{x}) = \frac{1}{Z} e^{\beta \sum_{i \sim j} I(x_i = x_j)}$$





- Conditional distributions:

$$\Pr(x_i = 1 | \mathbf{x}_{-i}) = \frac{e^{-\beta \sum_{j:j \sim i} I(x_j = 1)}}{e^{-\beta \sum_{j:j \sim i} I(x_j = 1)} + e^{-\beta \sum_{j:j \sim i} I(x_j = -1)}}$$

- Simulations: $\beta = 0.7$, $ss = \sum_{i \sim j} I(x_i = x_j)$
- R-script: `Ising.R`

Example - Ising model



Properties of Gibbs sampler (random scan)

- Gibbs sampling (random scan):
 - 1 Select starting values $\mathbf{x}^{(0)}$ and set $t = 0$
 - 2 Sample $j \sim \text{Uniform}\{1, \dots, p\}$
 - 3 Sample $X_j^{(t+1)} \sim p(x_j | \mathbf{x}_{-j}^{(t)})$
 - 4 Put $X_k^{(t+1)} = X_k^{(t)}$ for $k \neq j$

Properties of Gibbs sampler (random scan)

- Gibbs sampling (random scan):
 - 1 Select starting values $\mathbf{x}^{(0)}$ and set $t = 0$
 - 2 Sample $j \sim \text{Uniform}\{1, \dots, p\}$
 - 3 Sample $X_j^{(t+1)} \sim p(x_j | \mathbf{x}_{-j}^{(t)})$
 - 4 Put $X_k^{(t+1)} = X_k^{(t)}$ for $k \neq j$
- The chain $\{\mathbf{x}^{(t)}\}$ is Markov
- Detailed balance:
 - Consider \mathbf{x}, \mathbf{x}^* where $x_j \neq x_j^*$ while $x_k = x_k^*$ for $k \neq j$

Properties of Gibbs sampler (random scan)

- Gibbs sampling (random scan):
 - 1 Select starting values $\mathbf{x}^{(0)}$ and set $t = 0$
 - 2 Sample $j \sim \text{Uniform}\{1, \dots, p\}$
 - 3 Sample $X_j^{(t+1)} \sim p(x_j | \mathbf{x}_{-j}^{(t)})$
 - 4 Put $X_k^{(t+1)} = X_k^{(t)}$ for $k \neq j$
- The chain $\{\mathbf{x}^{(t)}\}$ is Markov
- Detailed balance:
 - Consider \mathbf{x}, \mathbf{x}^* where $x_j \neq x_j^*$ while $x_k = x_k^*$ for $k \neq j$

$$\begin{aligned}\pi(\mathbf{x})P(\mathbf{x}^*|\mathbf{x}) &= \pi(\mathbf{x})p^{-1}p(x_j^*|\mathbf{x}_{-j}) \\ &= \pi(\mathbf{x}_{-j})p(x_j|\mathbf{x}_{-j})p^{-1}p(x_j^*|\mathbf{x}_{-j}) \\ &= \pi(\mathbf{x}_{-j}^*)p(x_j|\mathbf{x}_{-j}^*)p^{-1}p(x_j^*|\mathbf{x}_{-j}^*) \\ &= \pi(\mathbf{x}^*)p^{-1}p(x_j|\mathbf{x}_{-j}^*) \\ &= \pi(\mathbf{x}^*)P(\mathbf{x}|\mathbf{x}^*)\end{aligned}$$

Tuning the Gibbs sampler

- Random or deterministic scan?
 - Deterministic scan most common (?)
 - When high correlation, random scan can be more efficient

Tuning the Gibbs sampler

- Random or deterministic scan?
 - Deterministic scan most common (?)
 - When high correlation, random scan can be more efficient
- Blocking:
 - When dividing $\mathbf{x} = (X_1, \dots, X_p)$, each X_j **can be vectors**
 - Making each X_j as large as possible will typically improve convergence
 - Especially beneficial when high correlation between single components
- Hybrid Gibbs sampling
 - If $p(x_j | \mathbf{x}_{-j})$ is difficult to sample from, use an Metropolis-Hastings step for this component
 - Example ($p = 5$)
 - 1 Sample $X_1^{(t+1)} \sim p(x_1 | \mathbf{x}_{-1}^{(t)})$
 - 2 Sample $(X_2^{(t+1)}, X_3^{(t+1)})$ through an M-H step
 - 3 Sample $X_4^{(t+1)}$ through another M.H step
 - 4 Sample $X_5^{(t+1)} \sim p(x_5 | \mathbf{x}_{-5}^{(t+1)})$

Lemmings data

- Model

$$\mathbf{y}_t \sim \text{Binom} \left(1, \frac{\exp(\mathbf{x}_t)}{1 + \exp(\mathbf{x}_t)} \right)$$

$$\mathbf{x}_t = a\mathbf{x}_{t-1} + \varepsilon_t, \quad \varepsilon_t \sim N(0, \sigma^2)$$

$$a \sim \text{Uniform}[0, 1]$$

- Simulation of a given \mathbf{x} :

$$a|\mathbf{x} \sim N \left(\frac{\sum_{i=2}^T x_i x_{i-1}}{\sum_{i=2}^T x_{i-1}^2} \middle| 0 < a < 1 \right)$$


- Simulation of x_i :

- 1 Simulate $x_i^* | \mathbf{x}_{-i} = N \left(\frac{a(x_{i-1} + x_{i+1})}{1 + a^2}, \sigma \frac{1 - a^2}{1 - a^4} \right)$



- Calculate $r = \frac{p(y_i | x_i^*)}{p(y_i | x_i)}$

- Accept x_i^* with probability $\min\{1, r\}$

Lemmings - results



Lemmings - results



- G. H. Givens and J. A. Hoeting. *Computational statistics*, volume 710. John Wiley & Sons, 2012.
- W. K. Hastings. Monte carlo sampling methods using markov chains and their applications. *Biometrika*, 57(1):97–109, 1970.
- N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, and E. Teller. Equation of state calculations by fast computing machines. *The journal of chemical physics*, 21(6):1087–1092, 1953.
- C. P. Robert and G. Casella. *Monte Carlo statistical methods*, volume 2. Springer, 1999.