#### AE1MCS: Mathematics for Computer Scientists

Heshan Du and Huan Jin University of Nottingham Ningbo China

Oct 2020

1/23

#### Reading

Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, 7th Edition, 2013.

■ Chapter 7, Section 7.1 An Introduction to Discrete Probability

#### Discrete Probability

- Combinatorics and probability theory share common origins (analyzing gambling games).
- The theory of probability now plays an essential role in a wide variety of disciplines (e.g. the study of genetics).
- In computer science,
  - Probability theory plays an important role in the study of the complexity of algorithms.
  - Probabilistic algorithms vs. deterministic algorithms.
  - Probability theory can help us answer questions that involve uncertainty.
  - **...**

#### Content

- Probability of an Event
- Probabilities of Complements and Unions of Events


#### Monty Hall Three-Door Puzzle

You are asked to select one of three doors to open; the large prize is behind one of the three doors and the other two doors are losers. Once you select a door, the game show host, who knows what is be behind each door:

- whether or not you selected the winning door, he opens one of the other two doors that he knows is a losing door (selecting at random if both are losing doors).
- Then he asks you whether you would like to switch doors.

# The Monty Hall Problem

W/L Switch? (st duil. Review Rony Partrick 3



## Finite Probability

Laplace's definition of the probability of an event with **finitely many**, **equally likely**, **possible outcomes** is as follows.

#### Definition

If S is a finite nonempty sample space of equally likely outcomes, and E is an event, that is, a subset of S, then the probability of E is

$$p(E) = \frac{|E|}{|S|}$$

- An **experiment** is a procedure that yields one of a given set of possible outcomes.
- The **sample space** of the experiment is the set of possible outcomes.
- An event is a subset of the sample space.



### Probability of an Event

In the eighteenth century, the French mathematician Laplace, who also studied gambling, defined the probability of an event as the number of successful outcomes divided by the number of possible outcomes.

7/23

### Example (Answer)

A box contains 4 blue balls and 5 red balls. What is the probability that a ball chosen at random from the box is blue?

49

### Example

A box contains 4 blue balls and 5 red balls. What is the probability that a ball chosen at random from the box is blue?

$$S = \{b1, b2, b3, b4, r1, r2, r3, r4, r5\}$$

$$E = \{b1, b2, b3, b4\}$$

$$p(E) = \frac{|E|}{|S|} = \frac{4}{9}$$

SO



#### **Example: Lottery**

In a lottery, players win a large prize when they pick four digits that match, in the correct order, four digits selected by a random mechanical process. A smaller prize is won if only three digits are matched.

- What is the probability that a player wins the large prize?
- What is the probability that a player wins the small prize?





## Example: Lottery (Answer)

$$\textit{S} = \{\textit{x} \in \mathbb{Z} : 0 \leq \textit{x} \leq 9999\} \Rightarrow |\textit{S}| = 10^4$$

- What is the probability that a player wins the large prize?  $E_1 = \{p\}$  where p is the lottery number picked by player, so  $|E_1| = 1$  and  $p(E) = 1/10^4$ .
- What is the probability that a player wins the small prize?

$$E_2 = \{x : x \text{ is like } p \text{ but with just one digit change}\}$$

hence  $|E_2| = C(4,1) \times 9 = 4 \times 9$ , so  $P(E_2) = C(4,1) \times 9/10^4$ . That is, 9 ways to change one digit over four digits.

 Heshan Du, Huan Jin
 AE1MCS
 Oct 2020
 11 / 23

#### Example: Poker 1

Find the probability that a hand of five cards in poker contains four cards of one kind.

- A deck of cards contains 52 cards.
- There are 13 different kinds of cards, with four cards of each kind.
- These kinds are twos, threes, fours, fives, sixes, sevens, eights, nines, tens, jacks, queens, kings, and aces.
- There are 4 suits: spades, clubs, hearts, and diamonds, each containing 13 cards.

#### Example: Poker 1 (Answer)

Find the probability that a hand of five cards in poker contains four cards of one kind.

S is number of ways to choose any 5 cards from 52:

$$|S| = C(52, 5).$$

■ E is the number of ways to get four of a kind:  $|E| = 13 \times (52 - 4)$ , using product rule with



- Choose one of the 13 kinds which is repeated 4 times;
- Choose any remaining card for last card (52-4).
- Hence  $p(E) = \frac{13 \times (52 4)}{C(52, 5)} \approx 0.00024$ .



Heshan Du, Huan Jin AE

#### Example: Poker 2

What is the probability that a poker hand contains a full house, that is, three of one kind and two of another kind?

#### Example: Poker 2 (Answer)

What is the probability that a poker hand contains a full house, that is, three of one kind and two of another kind?

Same sample space: |S| = C(52, 5).

$$|E|$$
 = # ways to get two different kinds  $C(13,2)$ 

× # ways to select which is three cards of the same kind  $C(4,3)$ 

× # ways to choose three cards of the same kind  $C(4,2)$ 

=  $C(13,2) \times 2 \times C(4,3) \times C(4,2)$ 

Notice: 
$$C(13,2) \times 2 = P(13,2) = C(13,1)C(12,1)$$
, so all correct.

15/23

#### Probabilities of Complements and Unions of Events

SEE

#### Theorem

Let E be an event in a sample space S. The probability of the event  $\overline{E} = \overline{S} - E$ , the complementary event of E, is given by

$$p(\overline{E})=1-p(E)$$

#### Theorem

Let  $E_1$  and  $E_2$  be events in the sample space S. Then

$$p(E_1 \cup E_2) = p(E_1) + p(E_2) - p(E_1 \cap E_2)$$

How to prove them?



<**□** ▶ ∢ Ē ▶ ∢ Ē ▶ ○ Ē ◆○ ♀ ○

Heshan Du, Huan Jin AE1MCS Oct 2020 16 / 23

#### Example

$$|5| = 2^{10}$$

- A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?
- What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by either 2 or 5?

$$S = \{1, 2, 3, \dots, 100\}$$

$$Z_{1} = \{2, 4, 6, 8, [0, 12, \dots, 100] \mid E_{1} = 50$$

$$E_{2} = \{5, [0, 15, 20, \dots, 100] \mid E_{2} = 20$$

$$\{10, 20, 30, \dots, 100\}$$
AE1MCS
Oct 2020

Heshan Du, Huan Jin

#### Example 1 (Answer)

A sequence of 10 bits is randomly generated. What is the probability that at least one of these bits is 0?

- Sample space is all bit strings of length 10:  $|S| = 2^{10}$ .
- Think about the event when bit string has *no* 0's. Then  $E = \{111111111111\}$  and |E| = 1.  $|D(E)| = \frac{1}{2^n} = 2^{-10}$
- Now the event we are interested in (at least one 0) is complement of *E*, so

$$p(\overline{E}) = 1 - p(E) = 1 - 2^{-10}.$$

$$\frac{1}{2^{l_0}} = 2^{-l_0}$$

 Heshan Du, Huan Jin
 AE1MCS
 Oct 2020
 18 / 23

### Example 2 (Answer)

What is the probability that a positive integer selected at random from the set of positive integers not exceeding 100 is divisible by *either* 2 or 5?

- $S = \{x \in \mathbb{Z}^+ : x \le 100\}$ , so |S| = 100;
- $E_1 = \{x \in S : x \text{ is divisible by 2}\}$ , so  $|E_1| = 50$ ;
- $E_2 = \{x \in S : x \text{ is divisible by 5}\}, \text{ so } |E_2| = 20;$
- then  $E_1 \cap E_2 = \{x \in S : x \text{ is divisible by } 10\}$ , so  $|E_1 \cap E_2| = 10$ ,

SO

$$p(E_1 \cup E_2) = p(E_1) + p(E_2) - p(E_1 \cap E_2) = \frac{50}{100} + \frac{20}{100} - \frac{10}{100} = 0.6$$



 Heshan Du, Huan Jin
 AE1MCS
 Oct 2020
 19 / 23

#### Monty Hall Three-Door Puzzle

You are asked to select one of three doors to open; the large prize is behind one of the three doors and the other two doors are losers. Once you select a door, the game show host, who knows what is be behind each door:

- whether or not you selected the winning door, he opens one of the other two doors that he knows is a losing door (selecting at random if both are losing doors).
- Then he asks you whether you would like to switch doors.

# Monty Hall Problem

#### Monty Hall Three-Door Puzzle

### Reading

Kenneth H. Rosen, *Discrete Mathematics and Its Applications*, 7th Edition, 2013.

■ Chapter 7, Section 7.1 An Introduction to Discrete Probability