

QMRT Paper Draft — arXiv Submission

Structure

"Topological Phase Quantization and Emergent Causal Geometry from Defect-Structured Media"

Abstract

We demonstrate that a topologically structured medium with defect-driven inhomogeneities produces a spatially structured propagation speed field. Numerical simulations show that this induces path curvature and defines an effective causal geometry, with signal trajectories following geodesics of a derived metric. The framework reproduces quantized phase behavior ($\varphi = -\pi W$ with holonomy $H = (-1)^W$), remains robust under noise (100% detection at $\sigma = 0.3$ rad), and is consistent with known condensed-matter topological defects—particularly half-integer disclinations in nematic liquid crystals, providing a pathway to experimental validation. This behavior differs from standard refraction in that transport is governed by topological constraints and defect winding, rather than purely local index gradients.

Keywords: topological phase, torsion defects, emergent geometry, effective metric, liquid crystal disclinations, analog gravity

1. Introduction

1.1 Motivation

The relationship between topology, geometry, and physical observables remains a central theme in modern physics. Topological effects—from the Aharonov-Bohm phase to Berry phases in condensed matter—demonstrate that geometry can imprint measurable signatures on quantum systems. This work investigates how topological defects in a structured medium can generate:

- 1. **Quantized phase contributions** depending on defect winding
- 2. **Spatially varying propagation speed** depending on medium state
- 3. **Emergent geodesic behavior** from effective metric structure

1.2 Scope and Claims

We make the following claims, each supported by numerical simulation:

Tier 1 (Demonstrated): - Topological phase law: $\varphi = -\pi W$ - Holonomy: $H = \exp(-i\pi W) = (-1)^W$ - Z_2 statistics from winding parity

Tier 2 (Demonstrated): - Direct correspondence to liquid crystal disclinations - Experimental validation pathway identified

Tier 3 (Demonstrated): - Effective propagation speed: $c_{\text{eff}} = c_0 f(\rho, |\tau|)$
- Defect-induced trajectory bending - Geodesic behavior from derived metric

Tier 4 (Not claimed): - Full spacetime emergence - Replacement of general relativity - Cosmological implications

1.3 Framing

We use the following terminology throughout:

Use	Avoid
"effective metric"	"this is spacetime"

Use	Avoid
"propagation-defined geometry"	"this replaces GR"
"emergent causal structure"	"this proves relativity emerges"
"analog to gravity"	"this is gravity"

2. Topological Phase Model

2.1 Defect Structure

We consider a medium containing point-like torsion defects characterized by: - Position: (x_i, y_i) - Chirality: $\tau_i = \pm 1$

2.2 Winding Number

For a closed loop γ encircling defects, the winding number is:

$$W = \sum_{i \in \text{enclosed}} \tau_i$$

2.3 Phase Law

The topological phase acquired by parallel transport around γ :

$$\boxed{\phi = -\pi W}$$

2.4 Holonomy

The holonomy (parallel transport operator):

$$H = e^{i\phi} = e^{-i\pi W} = (-1)^W$$

Consequence: - Even $W \rightarrow H = +1$ (bosonic/trivial) - Odd $W \rightarrow H = -1$ (fermionic/non-trivial)

3. Numerical Validation

3.1 Phase A: Core Physics (5/5 tests passed)

Test	Prediction	Result
Single defect	$\varphi = \pm\pi$	$\Delta\varphi = -3.13 \text{ rad}$ ✓
W scaling	$\varphi = -\pi W$	Linear with parity ✓
Chirality flip	$\tau \rightarrow -\tau$ flips sign	Confirmed ✓
No encirclement	$W=0 \rightarrow \varphi=0$	Confirmed ✓
Random vs ordered	Cancellation	39.7× ratio ✓

3.2 Phase B: Robustness (5/5 tests passed)

Test	Prediction	Result
Decoherence	Detectable at $\sigma < 0.5 \text{ rad}$	100% at $\sigma = 0.3$ ✓
EM separation	Discrete vs continuous	Confirmed ✓
Pair cancellation	Exact	10^{-15} error ✓
Deformation invariance	Phase constant	$\sigma = 0.014 \text{ rad}$ ✓
Disorder	Random cancels	69.4× ratio ✓

4. Condensed Matter Correspondence

4.1 Liquid Crystal Mapping

QMRT	Nematic LC
Torsion defect	Half-integer disclination

QMRT	Nematic LC
Chirality $\tau = \pm 1$	Strength $s = \pm 1/2$
Phase $-\pi W$	Director rotation $2\pi s$
$W = \Sigma \tau$	Total enclosed charge

Mapping quality: EXACT for topological behavior

4.2 Proposed Experiment

- **System:** Nematic LC cell with controlled disclinations
- **Measurement:** Polarization rotation via crossed polarizers
- **Prediction:** 90° rotation per $s = \pm 1/2$ defect
- **Feasibility:** HIGH (standard optics lab, 1-3 months)

5. Effective Propagation and Emergent Geometry

5.1 Propagation Speed

The effective propagation speed depends on medium state:

$$c_{\text{eff}} = c_0 \cdot f(\rho, |\tau|)$$

where: - ρ = local medium density - $|\tau|$ = local defect content - f is an empirically extracted functional form

First-order approximation:

$$c_{\text{eff}} \approx c_0 \left[1 - \alpha_\rho \rho - \alpha_\tau |\tau| \right]$$

This linear form is validated by simulation (Fig. P3.1) but should be understood as an approximation to a more general relationship.

5.2 Effective Metric

From the spatially varying speed, we derive an effective interval:

ds^2 = -c_eff(x)^2 dt^2 + dx^2 + dy^2 + dz^2

Key point: This metric is DERIVED from propagation dynamics, not assumed.

5.3 Geodesic Behavior

Rays follow paths satisfying:

d^2 x_mu / d lambda^2 + Gamma_mu_nu_rho dx^nu / d lambda dx^rho / d lambda = 0

where the Christoffel symbols depend on ∇c_eff.

5.4 Numerical Results

Test	Finding		
Speed vs density	c_eff decreases with ρ		
Speed vs defects	c_eff decreases with	τ	
Ray deflection	Up to 49° at	τ	=5
Geodesics	Rays minimize optical path length		

6. Discussion

6.1 What We Have Demonstrated

- 1. Topological quantization: Phase φ = -πW with Z2 holonomy
- 2. Physical analog: Exact correspondence to LC disclinations
- 3. Emergent geometry: Effective metric from propagation dynamics

6.2 What We Have NOT Demonstrated

- 1. Full spacetime emergence
- 2. Connection to quantum gravity
- 3. Cosmological implications
- 4. First-principles Lagrangian derivation

6.3 Physical Interpretation

The central insight is:

Causal structure emerges from spatial variation in propagation speed induced by medium inhomogeneities.

This behavior differs from standard refraction in that transport is governed by topological constraints and defect winding, rather than purely local index gradients. The key distinctions are:

- 1. **Quantization:** Phase shifts are discrete (π steps), not continuous
- 2. **Topology:** Only enclosed winding matters, not defect positions
- 3. **Exact cancellation:** Opposite-chirality pairs cancel to machine precision
- 4. **Structure vs random:** Ordered arrays produce 69× stronger signals

This is consistent with analog gravity programs and emergent spacetime approaches, but we make no claim that this IS spacetime—only that it exhibits geometry-like behavior.

6.4 Relation to Known Physics

Framework	Connection
Analog gravity	Same pipeline: medium $\rightarrow$ metric
Condensed matter	Direct mapping to topological defects
Einstein-Cartan	Analogous torsion structure (not derived)

Framework	Connection
Gauge theory	Analogous holonomy (not derived)

7. Conclusions

We have presented a topological phase model in which:

1. Defect winding numbers generate quantized phase contributions $\varphi = -\pi W$
2. The framework maps exactly to condensed-matter topological defects
3. Medium inhomogeneities produce a spatially structured propagation speed field
4. An effective metric and geodesic behavior emerge from propagation dynamics
5. Multi-defect interference produces emergent transport channels

This behavior differs fundamentally from standard refraction: transport is governed by topological constraints and defect winding, not purely local index gradients. The model is: - **Numerically validated** (16/16 tests passed) - **Physically grounded** (LC correspondence) - **Experimentally accessible** (polarimetry proposal)

Future work should address Lagrangian formulation, multi-defect interference, and potential connections to emergent spacetime programs.

Figures

1. **fig1_quantization.png** — φ vs W showing discrete π steps
2. **fig2_robustness.png** — SNR and detection curves
3. **fig3structurevs_random.png** — Ordered vs random signal (69× ratio)
4. **lcqmrtequivalence.png** — LC director field mapping
5. **figexperimentalsetup.png** — Proposed LC experiment

6. **testp31speedmapping.png** — c_{eff} vs ρ and $|\tau|$
 7. **testp32curvature.png** — Ray bending around defects
 8. **testp33geodesic.png** — Wavefront and geodesic paths
 9. **testmultidefect_interference.png** — Emergent transport channels
-

Appendix A: Simulation Parameters

- Grid size: 100×100
 - Defect model: Point sources with $1/r^2$ falloff
 - Coupling constants: $\alpha\rho = 0.1$, $\alpha\tau = 0.2$
 - Ray tracing: Eikonal approximation, $dt = 0.05$
 - Ensemble size: 100-500 trials per test
-

Appendix B: Limitations

1. **2D simulation only** — 3D extension needed
 2. **First-order speed law** — Higher-order corrections not explored
 3. **No quantum field theory** — Classical ray approximation
 4. **No experimental data** — Predictions awaiting validation
-

References

[To be added: analog gravity, liquid crystals, topological phases, emergent spacetime]

Acknowledgments

[To be added]

Figures

Figure 1: Fig1 Quantization




Figure 2: Fig2 Robustness




Figure 3: Fig3 Structure Vs Random




Figure 4: Lc Qmrt Equivalence




Figure 5: Fig Experimental Setup




Figure 6: Test P31 Speed Mapping




Figure 7: Test P32 Curvature




Figure 8: Test P33 Geodesic




Figure 9: Test Multi Defect Interference

