

Mathematics-3
Tutorial-3
Discussion on Friday, 23rd August
Topic: Discrete Random Variables

1. Letting X denote the random variable that is defined as the sum of two fair dice; then find the PMF. Check whether it is a valid PMF. Draw it.
2. Suppose that an airplane engine will fail, when in flight, with probability $1 - p$ independently from engine to engine; suppose that the airplane will make a successful flight if at least 50 percent of its engines remain operative. For what values of p is a four-engine plane preferable to a two-engine plane?
- 3.

Uniform Distribution:

A r.v. X is called a *uniform* r.v. over (a, b) if its pdf is given by

$$f_X(x) = \begin{cases} \frac{1}{b-a} & a < x < b \\ 0 & \text{otherwise} \end{cases}$$

Determine the CDF, mean and variance of this r.v. Plot the PDF and CDF.

4.

An information source generates symbols at random from a four-letter alphabet $\{a, b, c, d\}$ with probabilities $P(a) = \frac{1}{2}$, $P(b) = \frac{1}{4}$, and $P(c) = P(d) = \frac{1}{8}$. A coding scheme encodes these symbols into binary codes as follows:

a	0
b	10
c	110
d	111

Let X be the r.v. denoting the length of the code, that is, the number of binary symbols (bits).

- (a) What is the range of X ?
- (b) Assuming that the generations of symbols are independent, find the probabilities $P(X = 1)$, $P(X = 2)$, $P(X = 3)$, and $P(X > 3)$.

5.

Let X denote the number of heads obtained in the flipping of a fair coin twice.

- (a) Find the pmf of X .
- (b) Compute the mean and the variance of X .

6.

Consider the function given by

$$p(x) = \begin{cases} \frac{k}{x^2} & x = 1, 2, 3, \dots \\ 0 & \text{otherwise} \end{cases}$$

where k is a constant. Find the value of k such that $p(x)$ can be the pmf of a discrete r.v. X .

7.

A digital transmission system has an error probability of 10^{-6} per digit. Find the probability of three or more errors in 10^6 digits by using the Poisson distribution approximation.

8.

Suppose that independent trials, each having probability $p, 0 < p < 1$, of being a success are performed until a total of r successes is accumulated. If we let X equal the number of trials required, then

Prove that

$$P\{X = n\} = \binom{n-1}{r-1} p^r (1-p)^{n-r} \quad n = r, r+1, \dots$$

Find the mean and variance of this RV.