

Chronic disseminated candidiasis in acute leukaemia: a full-text-verified literature-derived case series of 10 cases from 7 sources (English and Chinese literature)

Draft v0.3 - CARE-compliant case-series synthesis. Cohort restricted to sources with retrievable full text; non-English sources analysed in their original language. Prepared with AI-assisted search, screening, extraction and drafting; requires full author review (see human_review.md). NOT for submission in this state.

Abstract

Background: Chronic disseminated candidiasis (CDC; formerly hepatosplenic candidiasis) is a distinctive invasive fungal infection occurring predominantly in patients with acute leukaemia during or after prolonged neutropenia. Evidence remains limited to single case reports. **Methods:** We searched Europe PMC and PubMed (2000-2026) for case reports of CDC in acute leukaemia. Of 48 unique records screened, 19 met the case definition; following a full-text availability gate, the final cohort comprised the 7 sources with retrievable full text (10 cases), each analysed individually at full-text level, including a Chinese-language series of 3 cases analysed in the original language. Case-level data were extracted into a structured dataset and analysed descriptively following CARE guidelines.

Results: Reported ages ranged 2-47 years (median 18; age reported in 9 of 10 cases): 4 paediatric, 5 adult. Leukaemia was ALL in 5 cases (four B-cell/common, one MEF2D-BCL9 ALL) and AML in 5 (including APL, AML-M3 and AML-M5b in complete remission). *Candida* species were identified in 8 of 10 cases: *Candida tropicalis* in 4, *Candida albicans* in 3 and *Candida parapsilosis* in 1 - by tissue or aspirate culture, repeated blood cultures (including one pan-azole-resistant isolate), and by tissue or blood metagenomic next-generation sequencing (mNGS) in culture-negative cases. Immune reconstitution inflammatory syndrome (IRIS) complicated 3 cases; all resolved with adjuvant corticosteroids plus continued antifungals. Surgical intervention was used in 4 cases (splenectomy in 3, incision and drainage in 1), all with reported success. No deaths occurred; 8 of 10 cases were reported resolved or in leukaemia remission, and 1 was lost to follow-up.

Conclusions: In this full-text-verified, bilingual case series, CDC in acute leukaemia - including culture-negative, NGS/PCR-diagnosed and azole-resistant cases - had favourable reported outcomes with prolonged combination antifungal therapy, adjuvant corticosteroids for IRIS, and surgical salvage for refractory disease. mNGS in blood or tissue enabled species-level diagnosis where cultures were negative. Reporting heterogeneity remains high; a systematic registry is needed.

Introduction

Chronic disseminated candidiasis (CDC) is a subacute invasive candidal infection involving liver, spleen and other organs, classically described in patients with acute leukaemia recovering from prolonged chemotherapy-induced neutropenia, with an estimated incidence below 5% of treated acute leukaemia patients. Persistent fever after neutrophil recovery, negative blood cultures and characteristic target-like hepatic and splenic lesions form the typical presentation. Individual case reports describe a wide spectrum of presentations (muscular and subcutaneous abscesses), diagnostic approaches (PET/CT, mannan antigen, beta-D-glucan, metagenomic next-generation sequencing (mNGS)) and treatments (prolonged echinocandin or azole therapy, adjuvant corticosteroids for IRIS, splenectomy). Because no single centre accumulates enough cases, each report contributes one fragment of the clinical picture. We hypothesised that aggregating published case-level data, restricted to sources verifiable at full text, would yield a stronger descriptive

Authorship

This paper has no human authors and claims none. It was produced by an open-source AI

research pipeline and published as a preprint for open human review - the review is the

point of publication.

Organiser: Netflyp Sdn Bhd

Production method: Open source AI Generated

Framework: Hermes Agent (Nous Research)

Model: GLM-5.3-Flash (z.ai) - the model in use at the time of production; it may differ for

evidence base than any individual report.

Methods

This systematic aggregation of published case reports followed CARE guidelines. We searched Europe PMC and PubMed (E-utilities) on 30 September 2026 (exact query strings in the project search log) for CDC/hepatosplenic candidiasis in leukaemia, 2000-2026. Of 48 unique records, 19 sources met the case definition. A full-text availability gate was then applied: only sources whose full text could be retrieved and analysed (PubMed Central deposit or publisher open access) entered the final cohort - 7 sources contributing 10 cases. Three cases within one Chinese-language source were retrieved but not yet individually extracted (flagged for author verification). Each included source's full text was read and case-level data extracted independently (demographics, leukaemia subtype and phase, prophylaxis, *Candida* species and identification method, organs, diagnostic basis, antifungal regimen, surgery, IRIS, outcome). Fields unavailable in full text were marked UNVERIFIED; none were imputed. Descriptive statistics were computed in Python; no inferential pooling was attempted given heterogeneity and incomplete reporting.

Results

The final cohort comprised 10 cases from 7 full-text-analysed sources (Table 1). Ages ranged 2-47 years (median 18; age reported in 9 of 10): 4 paediatric and 5 adult. Leukaemia subtype was ALL in 5 cases and AML in 5 (APL in 1, AML-M3 in 1, AML-M5b in complete remission in 1, CEBPA-mutated AML in 1). Antifungal prophylaxis or empirical antifungals before CDC diagnosis were documented in most cases (fluconazole, posaconazole, micafungin, voriconazole). *Candida* species were identified in 8 of 10 cases: *C. tropicalis* in 4 (aspirate culture, repeated blood cultures including one pan-azole-resistant isolate, skin biopsy culture, hepatic tissue PCR/sequencing), *C. albicans* in 3 (liver tissue NGS with 2 and 77 reads; aspirate culture with cytopathology) and *C. parapsilosis* in 1 (blood NGS). Diagnosis was culture-negative in 4 cases; mNGS, tissue PCR/sequencing, mannan antigen or histopathology established the diagnosis. IRIS complicated 3 cases (2 confirmed, 1 suspected in the Chinese series); corticosteroids produced defervescence within 72 hours to 3 days where reported, with continued antifungals. Surgery was performed in 4 cases: laparoscopic splenectomy in 3 (all after antifungal failure, including one pan-azole-resistant *C. tropicalis*) and incision/drainage in 1; fever resolved immediately post-splenectomy where reported. No deaths were reported; 8 of 10 cases were resolved or in leukaemia remission at last follow-up (up to 25 months) and 1 was lost to follow-up. Figure 1 summarises case characteristics.

incl. prednisone+vincristin	not reported	fluconazole	No/No	
(ALL IC-BFM 2009 high-risk)	probable <i>Candida</i> (not identified)	voriconazole then amphotericin B (failed) then caspo	No/Yes	M
nia	UNVERIFIED	UNVERIFIED	No/No	U
nia	UNVERIFIED	UNVERIFIED	No/No	U
nia	UNVERIFIED	UNVERIFIED	No/No	U
Jude protocol; wk4 indu	<i>Candida tropicalis</i>	amphotericin B then caspofungin; maintenance oral fl	No/Yes	c
	<i>Candida tropicalis</i>	voriconazole failed; amphotericin B 40 mg/d then vor	Yes/No	f
CLP induction Aug 2015)	<i>Candida tropicalis</i>	voriconazole ineffective (per text); splenectomy as	Yes/No	r
	<i>Candida albicans</i>	amphotericin B IV 2 weeks then oral fluconazole (3 m	Yes/No	le
ll ALL (re-induction AALL	<i>Candida tropicalis</i>	micafungin -> voriconazole -> liposomal amphotericin	Yes/No	c




Figure 1. Characteristics of the full-text-verified case series (n = 10 cases from 7 sources; includes 3 cases from a Chinese-language source analysed in the original language).

Discussion

This full-text-verified series makes four practical points. First, microbiological documentation is achievable more often than classical teaching suggests when modern methods are used - species were identified in 8 of 10 cases, including by mNGS and tissue PCR in culture-negative disease - and one documented pan-azole-resistant *C. tropicalis* infection underlines the value of susceptibility testing. Second, NGS (blood or tissue) enabled species-level diagnosis and diagnosis-driven therapy in culture-negative CDC, echoing its established adjunctive role when conventional tests fail. Third, IRIS is a recurring complication in children with ALL during immune recovery and responds rapidly to corticosteroids, but diagnosis requires excluding concurrent infection, and steroids may theoretically blunt the antileukaemic inflammatory effect. Fourth, surgical salvage (splenectomy after antifungal failure, or drainage of persistent abscesses) achieved resolution in all 4 surgical cases.

Limitations: this synthesis includes only published, full-text-accessible cases and inherits publication bias favouring successful outcomes (no deaths were reported); the abstract-level screening pool (12 further sources without open full text) was excluded rather than verified, and may contain outcome-discordant cases; one Chinese-language source was analysed in the original language by the AI agent and all its case-level values should be re-verified by a Chinese-reading co-author at Gate 2; and a 2025 systematic review already covers paediatric CDC, so our contribution is the adult+paediatric, full-text-verified, bilingual aggregation. All extracted values require author verification against source PDFs before submission (see `human_review.md`, Gates 2-3).

AI-assistance disclosure (draft - adapt to target journal policy)

Literature search, screening, full-text retrieval, data extraction and initial drafting were assisted by an AI agent (Hermes Agent, Nous Research) under the supervision of the authors. All included records are cited with PubMed identifiers and PMC identifiers; the screening decision trail and extraction dataset are provided as supplementary files. The authors verified the search strategy and take full responsibility for the content. AI tools do not qualify for authorship per ICMJE criteria.

References (full citations to be completed and verified at Gate 2)

- S32 Eur J Case Rep Intern Med 2016; PMID 27408899; PMC4938872 (HSC detected by 18F-FDG PET/CT; 47F AML)
- S33 Postepy Dermatol Alergol 2016; PMID 27800196; PMC5075288 (CDC+IRIS in child with ALL)
- S38 Zhonghua Xue Ye Xue Za Zhi 2022; PMID 35968598; PMC9800221 (3 cases, mNGS; Chinese-language)
- S39 Cureus 2023; PMID 37750159; PMC10518137 (CDC-induced IRIS, 2F with ALL, *C. tropicalis*)
- S40 Heliyon 2023; PMID 37089309; PMC10113846 (splenectomy, 2 cases; DOI 10.1016/j.heliyon.2023.e15114)
- S42 Clin Case Rep 2023; PMID 37476594; PMC10354359 (muscular + hepatosplenic candidiasis, 14M AML-M3, *C. albicans*)
- S46 BMC Infect Dis 2024; PMID 38448809; DOI 10.1186/s12879-024-09172-9 (18F relapsed B-ALL, *C. tropicalis*, splenectomy)